

Chemistry (Salters)

Advanced GCE **A2 7887**

Advanced Subsidiary GCE **AS 3887**

Mark Schemes for the Units

January 2010

3887/7887/MS/R/10J

OCR (Oxford Cambridge and RSA) is a leading UK awarding body, providing a wide range of qualifications to meet the needs of pupils of all ages and abilities. OCR qualifications include AS/A Levels, Diplomas, GCSEs, OCR Nationals, Functional Skills, Key Skills, Entry Level qualifications, NVQs and vocational qualifications in areas such as IT, business, languages, teaching/training, administration and secretarial skills.

It is also responsible for developing new specifications to meet national requirements and the needs of students and teachers. OCR is a not-for-profit organisation; any surplus made is invested back into the establishment to help towards the development of qualifications and support which keep pace with the changing needs of today's society.

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which marks were awarded by Examiners. It does not indicate the details of the discussions which took place at an Examiners' meeting before marking commenced.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes should be read in conjunction with the published question papers and the Report on the Examination.

OCR will not enter into any discussion or correspondence in connection with this mark scheme.

© OCR 2010

Any enquiries about publications should be addressed to:

OCR Publications
PO Box 5050
Annesley
NOTTINGHAM
NG15 0DL

Telephone: 0870 770 6622
Facsimile: 01223 552610
E-mail: publications@ocr.org.uk

CONTENTS




Advanced GCE Chemistry (Salters) (7887)

Advanced Subsidiary GCE Chemistry (Salters) (3887)

MARK SCHEME ON THE UNITS


Unit/Content	Page
2849 Chemistry of Materials	1
2854 Chemistry by Design	9
Grade Thresholds	19





2849 Chemistry of Materials

Question	Expected answers	Marks
1 (a) (i)	 <p>(1) ; allow without the C within the ring.</p>	1
1 (a) (ii)	 <p>all bonds must be shown (1) ;</p>	1
1 (b) (i)	 <p>one of these oxygens only</p> <p>ester linkage correct (1) ; rest correct (1) ; ignore brackets allow ecf from 1(a).</p>	2


Question	Expected answers	Marks
1 (b) (ii)	(below T_g) chains do not have enough energy (may describe in terms of vibration or motion of chains) (1) ; to move over / slide over one another (1) ; force applied to change shape of polymer will cause 'frozen' chains to break / AW (1) ;	3
1 (b) (iii)	intermolecular forces / bonds between chains are greater (1) ; because chains are able to get closer (because of the flat ring system) / so more energy is needed to separate the chains(1) ;	2
1 (c)	burning / combustion (1) ; energy produced can be used / reducing landfill (1) ; or recycling AW (1) ; oil resources saved AW /r educing landfill (1) ; do NOT allow cracking, but allow reducing landfill.	2
1 (d) (i)	$K_C = \frac{[C] \times [H_2O]^2}{[B] \times [C_2H_5OH]^2}$ [Products] / [Reactants] (1) ; indices correct (1) ;	2
1 (d) (ii)	equilibrium position moves in exothermic direction / right since forward reaction is exothermic AW (1) ; K_c , increases (1) ; <i>ecf here for second mark.</i>	2
Total mark		15

Question	Expected answers	Marks
2 (a) (i)	+5 (1) ; accept 5+	1
2 (a) (ii)	H ⁺ / HCl in hydrogen electrode beaker (1) ; both electrodes made from Pt (1) ; electrode dipping into a solution VO ₂ ⁺ and VO ²⁺ containing H ⁺ ions (1) ; conditions given as 1 mol dm ⁻³ concentrations and 298K / 1 atm (1) ; salt bridge and circuit correct (1) ;	5
2 (b) (i)	1.26 V (1) ;	1
2 (b) (ii)	V ²⁺ / V ³⁺ (<i>may give more detail of half-cells</i>) because it has the more negative / less positive electrode potential / AW / in terms of reducing agent / oxidizing agent or electron transfer (1) ;	1
2 (c) (i)	V ²⁺ → V ³⁺ + e ⁻ (1) ;	1
2 (c) (ii)	V ²⁺ + VO ₂ ⁺ + 2H ⁺ → V ³⁺ + VO ²⁺ + H ₂ O correct reactants and products (1) ; equation given balanced correctly (1) ;	2
2 (d) (iii)	specific frequencies of visible radiation / light absorbed (by ions in solution) (1) ; rest of visible light transmitted (as green colour) AW (1) ;	2
Total mark		13

Question	Expected answers	Marks						
3 (a)	<p>1 mark for points in bold, then any 2 from 3:</p> <p>dissolve the sample in the minimum amount / AW (1) ;</p> <p>of hot ethanol (1) ;</p> <p>filter (off any solid impurities) (1) ;</p> <p>leave (solution / filtrate) to cool/to form crystals (1) ;</p> <p>filter off crystals / decant solution (1) ;</p> <p>wash crystals (1) ;</p> <p>and dry (1) ;</p> <p>QWC see separate sheet (1) ;</p>	<p>6</p> <p>1</p>						
3 (b) (i)	<p>broad peak / absorbance around 3100 cm^{-1} indicates O-H (in carboxylic acid) (1) ;</p> <p>strong peak / absorbance around 1720 cm^{-1} indicates C=O (in carboxylic acid) (1) ;</p> <p>hence -COOH / carboxylic acid (1) ;</p> <p><i>the first two marks are for identifying the two important peaks, however much detail is given. These may be shown on the spectrum.</i></p>	3						
3 (b) (ii)	<div></div> <p>correct molecular formula (1) ;</p> <p>correct structure (1) ; do not allow OH</p>	2						
3 (b) (iii)	<p>M_r of acetaminophen = 179.0 (1) ;</p> <p>mass of pure acetaminophen in sample = 0.010×179.0 i.e $\text{mol} \times M_r$ ecf M_r (1) ;</p> <p>percentage = $(1.790 / 3.00) \times 100 = 59.7\%$ ecf (1) ;</p>	3						
3 (c) (i)	<p>iron(III) chloride in solution is yellow</p> <p>accept brown / yellow or brown + orange / red (1) ;</p> <table><tr><th>compound</th><th>observations on adding aqueous Iron(III) chloride</th></tr><tr><td>acetaminophen</td><td>turns purple/violet (1);</td></tr><tr><td>phenacetin</td><td>remains yellow/brown/colour does not change ecf (1).</td></tr></table>	compound	observations on adding aqueous Iron(III) chloride	acetaminophen	turns purple/violet (1);	phenacetin	remains yellow/brown/colour does not change ecf (1).	3
compound	observations on adding aqueous Iron(III) chloride							
acetaminophen	turns purple/violet (1);							
phenacetin	remains yellow/brown/colour does not change ecf (1).							

Question	Expected answers	Marks															
3 (c) (ii)	<table border="1"> <thead> <tr> <th>chemical shifts for acetaminophen</th><th>type of proton</th><th>relative intensity</th></tr> </thead> <tbody> <tr> <td>4.5 -10.0 (1) ; <i>only one peak otherwise no marks</i></td><td>  / phenolic OH </td><td></td></tr> </tbody> </table> <table border="1"> <thead> <tr> <th>chemical shifts for phenacetin</th><th>type of proton</th><th>relative intensity</th></tr> </thead> <tbody> <tr> <td>3.6</td><td>—O—CH₂—R</td><td>2</td></tr> <tr> <td>0.8-1.2</td><td>R—CH₃</td><td>3 (1) ; <i>for relative intensities</i></td></tr> </tbody> </table> <p>1 mark for each set of shift & proton type correct (2)</p>	chemical shifts for acetaminophen	type of proton	relative intensity	4.5 -10.0 (1) ; <i>only one peak otherwise no marks</i>	 / phenolic OH		chemical shifts for phenacetin	type of proton	relative intensity	3.6	—O—CH ₂ —R	2	0.8-1.2	R—CH ₃	3 (1) ; <i>for relative intensities</i>	5
chemical shifts for acetaminophen	type of proton	relative intensity															
4.5 -10.0 (1) ; <i>only one peak otherwise no marks</i>	 / phenolic OH																
chemical shifts for phenacetin	type of proton	relative intensity															
3.6	—O—CH ₂ —R	2															
0.8-1.2	R—CH ₃	3 (1) ; <i>for relative intensities</i>															
3 (d) (i)	phenol / hydroxyl (1) ;	1															
3 (d) (ii)	 negative ion formed by proton loss (1) ; correct structure (1) ; allow 1 mark if amide bond is hydrolysed but phenol has lost proton	2															
Total mark		26															

Question	Expected answers	Marks															
4 (a)	corrosion resistance / stainless / hard / lustrous AW / high strength / hard / AW (1) ;	1															
4 (b) (i)	magnesium sulphide / MgS (1) ;	1															
4 (b) (ii)	coolant (oxygen blow is very exothermic) (1) ;	1															
4 (b) (iii)	any two from: carbon, phosphorus, manganese, silicon (1) ;	1															
4 (c)	<table border="1"><thead><tr><th>statement</th><th>true</th><th>false</th></tr></thead><tbody><tr><td>Iron is reduced</td><td></td><td>✓</td></tr><tr><td>Iron(II) ions are oxidized</td><td>✓</td><td></td></tr><tr><td>Electrons move through the water in the cell that is set up</td><td></td><td>✓</td></tr><tr><td>Iron(II) ions form a brown precipitate in the presence of hydroxide ions</td><td></td><td>✓</td></tr></tbody></table>	statement	true	false	Iron is reduced		✓	Iron(II) ions are oxidized	✓		Electrons move through the water in the cell that is set up		✓	Iron(II) ions form a brown precipitate in the presence of hydroxide ions		✓	2
statement	true	false															
Iron is reduced		✓															
Iron(II) ions are oxidized	✓																
Electrons move through the water in the cell that is set up		✓															
Iron(II) ions form a brown precipitate in the presence of hydroxide ions		✓															
4 (d) (i)	<div style="display: flex; justify-content: space-around; align-items: flex-start;"><div style="text-align: center;">3d Ni <table border="1" style="display: inline-table;"><tr><td>↑↓</td><td>↑↓</td><td>↑↓</td><td>↑</td><td>↑</td></tr></table></div><div style="text-align: center;">4s <table border="1" style="display: inline-table;"><tr><td>↑↓</td></tr></table></div></div> <div style="display: flex; justify-content: space-around; align-items: flex-start; margin-top: 10px;"><div style="text-align: center;">Ni²⁺ <table border="1" style="display: inline-table;"><tr><td>↑↓</td><td>↑↓</td><td>↑↓</td><td>↑</td><td>↑</td></tr></table></div><div style="text-align: center;"><table border="1" style="display: inline-table;"><tr><td> </td></tr></table></div></div> <p>correct number of electrons in Ni (1) ; correct arrangement for Ni²⁺ (1) ;</p>	↑↓	↑↓	↑↓	↑	↑	↑↓	↑↓	↑↓	↑↓	↑	↑		2			
↑↓	↑↓	↑↓	↑	↑													
↑↓																	
↑↓	↑↓	↑↓	↑	↑													
4 (d) (ii)	3Ni + 2NO ₃ ⁻ + 8H ⁺ → 3Ni ²⁺ + 2NO + 4H ₂ O correct reactants and products (1) ; balanced (1) ;	2															
4 (e) (i)	[Ni(H ₂ O) ₆] ²⁺ = 6 (1) ; [Ni(dimethylglyoxime) ₂] ²⁺ = 4 (1) ;	2															
4 (e) (ii)	[Ni(H ₂ O) ₆] ²⁺ = octahedral (1) ; [Ni(dimethylglyoxime) ₂] ²⁺ = tetrahedral / square planar (1) ;	2															
Total mark		14															

Question	Expected answers	Marks
5 (a) (i)	lone pair of electrons on N (1) ; can accept proton/hydrogen ion/H ⁺ (1) ;	2
5 (a) (ii)	water (1) ;	1
5 (a) (iii)	 <p>one mark for both hydrogen bonds (1) ; one mark for both lone pairs (1) ; partial charges correct (1) ; If only one interaction shown but all three components are correct then give 2 marks.</p>	3
5 (a) (iv)	<u>double</u> helix (1) ;	1
5 (b) (i)	any two from the following four points smaller chain length / M_r (1) ; different bases (1) ; do not allow 'complementary bases'; RNA has single chain (1) ; different sugar in chain (1) ;	2
5 (b) (ii)	any two from <u>hydrogen bonds</u> between DNA strands break (1) ; DNA divides / unwinds / uncoils / chains separate (1) ; each strand acts as a template for new strand AW (1) ;	2
Total mark		11

Question	Expected answers	Marks
6 (a)	pentyl (1) ; ethanoate (1) ;	2
6 (b) (i)	$K_c = \frac{[\text{P}].[water]}{[acid].[alcohol]}$ (1) ; at equilibrium $[\text{P}] = [water]$ (1) ; $[\text{P}]^2 = 4.15 \times (1.06) \times (1.06)$ (1) ; $[\text{P}] = 2.16 \text{ mol dm}^{-3}$ answer must be to 3 sfs (1) ; ecf for incorrect equation.	4
6 (b) (ii)	product / compound P lost (1) ; concentrations of reactants will be less (1) ; K_c is unchanged (1) ; does not change with concentration / only changes with temperature / ratio of concentrations remain the same AW (1) ; allow 1 mark for loss of volatile component causing an increase in temperature.	4
6 (b) (iii)	<u>conc.</u> sulphuric acid (1) ;	1
Total mark		11


2854 Chemistry by Design

Question	Expected Answers	Marks
1 (a) (i)	ether (allow ethyl ether / ethoxy / alkoxy) (1) ;	1
1 (a) (ii)	(secondary) amide (1) ;	1
1 (a) (iii)	(primary) amine (1) ;	1
1 (b) (i)	105 – 110 (1) ;	1
1 (b) (ii)	120 (1) ;	1
1 (c)	iron(III) (chloride) (1) ; pink / purple / violet / mauve colour (1) ; <i>mark independently</i>	2
1 (d) (i)	hydrolysis (1);	1
1 (d) (ii)	(named/formula) strong alkali / (named / formula) moderately concentrated / dilute mineral acid (1) ; reflux (1) ; allow concentrated HCl any additional reagents CON first mark award second mark, provided alkali or acid incl conc sulphuric mentioned.	2
1 (e)	IR any one similar bond (1) ; any one different bond (1) ; IGNORE groups (eg 'amide') , except for arene / benzene <i>Mark first bond / absorbance range pair, ignore others</i> Absorbance ranges of both (correct) bond OR group given (1) ; <i>IR similar:</i> C–O (1050-1300); C–H (2850-2950/3000-3100); arene (several peaks in range 1450 – 1650) <i>IR difference:</i> C=O (1630-1700) in phenacetin; N–H (3500 in phenacetin OR ; 3300-3500 in phenetidine) NMR phenacetin: 6 peaks: 3:1:2:2:2:3 { allow 5 peaks: 3:1:4:2:3}(1) ; allow any order – count up if individual values given paraphenetidine (<i>mark first answer and ignore others</i>) 5 peaks / one less peak (allow one less than an incorrect answer for phenacetin) / 2 replaces 3:1 / 0.5 – 6 in spectrum / 5-12 or 2.2 not in spectrum (1) ;	5



Question	Expected Answers	Marks
1 (f) (i)	all of structure indicated, apart from H or O–H (1) ;	1
1 (f) (ii)	part of molecule / group (1) ; that / bonds / fits to receptor / active site (1) ; responsible for pain killing / AW (1) ;	3
1 (g)	idea of shape of <u>active site</u> or molecule fitting / bonding / binding into <u>active site</u> (1) ; mention of ether / $\text{–OC}_2\text{H}_5$ or phenol / –OH group (in terms of fitting / not fitting / bonding) (1) ;	2
	Total mark	21


Question	Expected Answers	Marks
2 (d)	not attached to any particular pair of atoms / spread over several / all C atoms (1) ; each C has one electron (not involved in bonding to C or H) (1) ; arranged in <u>rings</u> above and below ring /molecule [<i>allow diagram</i>] (1) ;	3
2 (e) (i)	positive ion / particle / molecule with partial positive charge attracted to a negatively charged / electron dense region / carbon (1) ; reacts by accepting a pair of electrons (to form a covalent bond) (1) ; (a substitution is where) one group / atom is exchanged for another (1) ;	3
2 (e) (ii)	<i>any three from four</i> polarises Br ₂ / forms Br ^{δ+} / allow forms Br ⁺ (1) ; Br ⁺ /Br ^{δ+} attacks/ accepts electrons from (benzene) / is the electrophile (1) ; Provides a <u>route/ pathway</u> of lower activation enthalpy/energy (1) ; FeBr ₃ restored at end (1) C <i>mark independently</i>	3
2 (e) (iii)	C ₆ H ₆ + Br ₂ → C ₆ H ₅ Br + HBr (1) for LHS (allow skeletal representation of benzene [or with –H]) ; (1) for RHS (depends on first marking point) ;	2
	Total mark	31

Question	Expected Answers	Marks
3 (d) (iii)	$\Delta S^{\circ}_{\text{surr}} = 552000/298 = (+)1852$ (1) ; <i>stated or implied (allow 1.85(2) if kJ units shown)</i> $\Delta S^{\circ}_{\text{tot}} = -7 + 1852^* = +1845$ (1) ; <i>correct expression and evaluation(with sign)</i> allow ecf from 3d(i) and first mpt [-5/-5.2/-5.15 scores 1 overall] *allow rounded to 1850 giving +1843	2
3 (e) (i)	<i>any two from:</i> more moles of gas / increase in volume/ pressure (1) ; exothermic / gas becomes hotter /(reaction) provides energy / AW (1) ; low activation energy / enthalpy or does not need much energy to decompose (1) ;	2
3 (e) (ii)	(N ₂ O decomposes to give) oxygen (AW) (1) ; because of the temperature of/ energy supplied by <u>splint</u> / AW / (allow N ₂ O decomposes at low temperatures) (1) ;	2
	Total mark	24

Question	Expected Answers	Marks
4 (a) (i)	water / dilute acid / brine / [any] hydrocarbon/ natural gas / petrol (1) ;	1
4 (a) (ii)	when burnt / used as fuels (1) ; SO ₂ / sulphuric acid produced (1) ; <i>mark separately</i> acid rain / an effect of acid rain (1) ; <i>depends on either of first two marking points being scored</i>	3
4 (b) (i)	2-aminoethanol/ 2-aminoethan-1-ol / 1-aminoethan-2-ol (1) ; allow 2-hydroxyethylamine	1
4 (b) (ii)	–NH ₃ ⁺ / S ²⁻ (1) ;	1
4 (c) (i)	incompletely ionised / incompletely dissociated / in equilibrium (with water) (1) ; ignore references to ability to donate protons	1
4 (c) (ii)	[H ⁺] [HS ⁻] / [H ₂ S] (1) ;	1
4 (c) (iii)	[H ⁺] = $\sqrt{(0.1 \times 8.9 \times 10^{-8})}$ (1) ; (= 9.43 x 10 ⁻⁵ (mol dm ⁻³)) pH = – log[H ⁺] (stated or implied) (1) ; = 4.03 (1) ; allow 4.0 or 4.025 or ecf from first mpt.	3
4 (d) (i)	[Hg ²⁺] [S ²⁻] charges on ions correct (wherever stated) (1) ; indication of concentration of Hg ion times concentration of S ion (1) ;	2
4 (d) (ii)	[Hg ²⁺] = $\sqrt{(4.0 \times 10^{-53})}$ (= 6.32 x 10 ⁻²⁷ (mol dm ⁻³)) (1) ; 'ans to 1st mpt' x 6.0 x 10 ²³ and evaluated (= 3.8 x 10 ⁻³ (ions dm ⁻³)) (1) ; <i>if first mpt not apparent, allow some number (not 1) x L (and evaluated) for 1 mark</i>	2
4 (d) (iii)	6.32 x 10 ⁻²⁷ (ecf from (d)(ii)) x 233 = 1.5 x 10 ⁻²⁴ (g dm ⁻³) (1) ;	1
4 (e)	 <p>correct O with dative covalent bond (1) ; rest correct (1) ; correct expanded octet with two double bonds scores 1 (ignore shape of molecule)</p>	2

Question	Expected Answers	Marks
4 (f)	$0.01 \times 98/32 = 0.031$ (kg) correct M_r values (1) ; rest of calculation & evaluation (ecf) (1) ; 2sf (provided some calculation shown)(1) ;	3
4 (g)	ionic (1) ; broken: ionic (1) ; hydrogen bonds (1) ; made: ion-dipole (1) ; allow other descriptions eg hydration of ions broken approx equal to made / AW (1) ; ions move to electrodes / ions free to move (1) ;	6
	Total mark	27

Question	Expected Answers	Marks
5 (a)	$(2260 \times 18 / 1000 =) +40.7$ (1) ; allow absence of + sign, allow 40.68 allow answer given in part (a) and not in table.	1
5 (b) (i)	energy absorbed by evaporation (1) ; released on condensation (1) ; <i>mark independently</i>	2
5 (b) (ii)	ratio energy / mass is high / AW (1) ;	1
5 (c) (i)	Either of these, or a mixture  correct pair of hydrogen bonds shown (1) ; (ignore other <i>correct</i> hydrogen bonds; incorrect hydrogen bonds are CON) for both hydrogen bonds: correct partial charges (1) ; lone pair along bond (1) ; straight O–H–O (1) ;	4
5 (c) (ii)	it only forms one / fewer hydrogen bond per molecule (1) ; less energy is needed to break / because it only contains one $\text{H}\delta^+$ / AW (1) ; ignore references to M_r and other imf. ORA throughout	2
5 (d) (i)	 (1) ; ignore incorrect bond angles	1
5 (d) (ii)	permanent (dipole) – permanent dipole (1) ; do not allow abbreviations no hydrogen attached to the oxygen atom / electronegative atom / electron withdrawing group (1) ;	2

Question	Expected Answers	Marks
5 (e) (i)	 (1) ;	1
5 (e) (ii)	<p>any three from:</p> <p>M_r / no. of electrons of hexane is larger / hexane is long(er) (1) ;</p> <p>hexane is less branched (ora) (1) ;</p> <p>more / stronger instantaneous dipole-induced dipole forces (1) ;</p> <p>compensate for permanent dipole-permanent dipole / pd-pd stronger than id-id for molecules of similar size / AW (1) ;</p> <p>allow abbreviations to imf names here.</p> <p>If 'id-id' given as answer to (d)(ii), allow one (in total) mark for ecf here</p>	3
	Total mark	17

Grade Thresholds

Advanced GCE Chemistry (Salters) (3887/7887)
January 2010 Examination Series

Unit Threshold Marks

Unit		Maximum Mark	a	b	c	d	e	u
2849	Raw	90	72	64	56	49	42	0
	UMS	90	72	63	54	45	36	0
2854	Raw	120	80	72	64	56	49	0
	UMS	120	96	84	72	60	48	0
2855	Raw	90	76	68	60	52	44	0
	UMS	90	72	63	54	45	36	0

Specification Aggregation Results

Overall threshold marks in UMS (ie after conversion of raw marks to uniform marks)

	Maximum Mark	A	B	C	D	E	U
3887	300	240	210	180	150	120	0
7887	600	480	420	360	300	240	0

The cumulative percentage of candidates awarded each grade was as follows:

	A	B	C	D	E	U	Total Number of Candidates
3887	15.4	46.2	53.8	61.5	92.3	100.00	14
7887	15.7	42.5	76.1	91.5	97.5	100.0	332

346 candidates aggregated this series

For a description of how UMS marks are calculated see:

http://www.ocr.org.uk/learners/ums_results.html

Statistics are correct at the time of publication.

OCR (Oxford Cambridge and RSA Examinations)
1 Hills Road
Cambridge
CB1 2EU

OCR Customer Contact Centre

14 – 19 Qualifications (General)

Telephone: 01223 553998

Facsimile: 01223 552627

Email: general.qualifications@ocr.org.uk

www.ocr.org.uk

For staff training purposes and as part of our quality assurance programme your call may be recorded or monitored

Oxford Cambridge and RSA Examinations
is a Company Limited by Guarantee
Registered in England
Registered Office; 1 Hills Road, Cambridge, CB1 2EU
Registered Company Number: 3484466
OCR is an exempt Charity



OCR (Oxford Cambridge and RSA Examinations)
Head office
Telephone: 01223 552552
Facsimile: 01223 552553