

A



Physics Equations Sheet

GCSE Physics (8463)

FOR USE IN JUNE 2022 ONLY

[Turn over]

HT = Higher Tier only equations

| | |
|---|--------------------------------|
| kinetic energy = $0.5 \times \text{mass} \times (\text{speed})^2$ | $E_k = \frac{1}{2} m v^2$ |
| elastic potential energy = $0.5 \times \text{spring constant} \times (\text{extension})^2$ | $E_e = \frac{1}{2} k e^2$ |
| gravitational potential energy = mass \times gravitational field strength \times height | $E_p = m g h$ |
| Change in thermal energy = mass \times specific heat capacity \times temperature change | $\Delta E = m c \Delta \theta$ |
| power = $\frac{\text{energy transferred}}{\text{time}}$ | $P = \frac{E}{t}$ |

| | |
|--|-------------------------------------|
| power = $\frac{\text{work done}}{\text{time}}$ | $P = \frac{W}{t}$ |
| efficiency = $\frac{\text{useful output energy transfer}}{\text{total input energy transfer}}$ | |
| efficiency = $\frac{\text{useful power output}}{\text{total power input}}$ | |
| charge flow = current × time | $Q = I t$ |
| potential difference = current × resistance | $V = I R$ |
| power = potential difference × current | $P = V I$ |

| | |
|---|---|
| power = (current)² × resistance | $P = I^2 R$ |
| energy transferred = power × time | $E = P t$ |
| energy transferred = charge flow × potential difference | $E = Q V$ |
| density = $\frac{\text{mass}}{\text{volume}}$ | $\rho = \frac{m}{V}$ |
| Thermal energy for a change of state = mass × specific latent heat | $E = m L$ |
| For gases: pressure × volume = constant | $p V = \text{constant}$ |
| weight = mass × gravitational field strength | $W = m g$ |
| work done = force × distance (along the line of action of the force) | $W = F s$ |

| | |
|---|--------------------------|
| force = spring constant × extension | $F = k e$ |
| moment of a force = force × distance (normal to direction of force) | $M = F d$ |
| pressure = $\frac{\text{force normal to a surface}}{\text{area of that surface}}$ | $p = \frac{F}{A}$ |
| Pressure due to column of liquid = height of column × density of liquid × gravitational field strength | $p = h \rho g$ |
| distance travelled = speed × time | $s = v t$ |
| acceleration = $\frac{\text{change in velocity}}{\text{time taken}}$ | $a = \frac{\Delta v}{t}$ |

HT

[Turn over]

| | |
|--|-------------------------------------|
| <p>HT</p> <p>Force on a conductor (at right angles to a magnetic field) carrying a current = magnetic flux density × current × length</p> | $F = B I l$ |
| <p>HT</p> <p><u>potential difference across primary coil</u> <u>potential difference across secondary coil</u> = <u>number of turns in primary coil</u> <u>number of turns in secondary coil</u></p> | $\frac{V_p}{V_s} = \frac{n_p}{n_s}$ |
| <p>HT</p> <p>potential difference across primary coil × current in primary coil = potential difference across secondary coil × current in secondary coil</p> | $V_p I_p = V_s I_s$ |

BLANK PAGE

**Physics Equations Sheet - GCSE
Physics (8463)**

FOR USE IN JUNE 2022 ONLY

IB/M/CD/Jun22/8463/INS/E1