AQA
Surname
Other Names
Centre Number
Candidate Number
Candidate Signature
A-level
BIOLOGY
Paper 2
7402/2
Thursday 13 June 2019 Morning

Time allowed: 2 hours

At the top of the page, write your surname and other names, your centre number,

your candidate number and add your signature.



For this paper you must have:

- a ruler with millimetre measurements
- a scientific calculator.

INSTRUCTIONS

- Use black ink or black ball-point pen.
- Answer ALL questions.
- You must answer the questions in the spaces provided. Do NOT write on blank pages.
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.




INFORMATION

- The marks for the questions are shown in brackets.
- The maximum mark for this paper is 91.

DO NOT TURN OVER UNTIL TOLD TO DO SO



Answer ALL questions in the spaces provided.



Succession occurs in natural ecosystems. Describe and explain how succession occurs. [4 marks]



5



FIGURE 1, on the opposite page, shows percentages of energy transferred from sunlight to a zebra in a grassland ecosystem.

01.2


Use FIGURE 1 to calculate the percentage of sunlight energy that would be transferred into the faeces and urine of a zebra. Give your answer to 3 significant figures. [1 mark]

%

Answer =



FIGURE 1



7

Respiration

New tissue

Faeces and urine



BLANK PAGE

8






In this ecosystem the net productivity of the vegetation is 24 525 kJ m⁻² year⁻¹

Use this information and FIGURE 1, on page 7, to calculate the energy stored in new tissues of the zebra in kJ m⁻² year ⁻¹ [2 marks]









02

Sickle cell disease (SCD) is a group of inherited disorders. People with SCD have sickle-shaped red blood cells. A single base substitution mutation can cause one type of SCD. This mutation causes a change in the structure of the beta polypeptide chains in haemoglobin.

02.1

Explain how a single base substitution causes a change in the structure of this polypeptide.

Do NOT include details of transcription and translation in your answer. [3 marks]








Haematopoietic stem cell transplantation (HSCT) is a long-term treatment for SCD.

In HSCT, the patient receives stem cells from the bone marrow of a person who does not have SCD. The donor is often the patient's brother or sister. Before the treatment starts, the patient's faulty bone marrow cells have to be destroyed.

02.2

Use this information to explain how HSCT is an effective long-term treatment for SCD. [3 marks]








A new long-term treatment for SCD involves the use of gene therapy.

FIGURE 2, on the opposite page, shows some of the stages involved in this treatment in a child with SCD.



15

FIGURE 2




modified stem cells injected back into the child



BLANK PAGE





Some scientists have concluded that this method of gene therapy will be a more effective long-term treatment for SCD than HSCT. Use all the information provided to evaluate this conclusion. [3 marks]





A student investigated the effects of indoleacetic acid (IAA) on the growth of oat seedlings (young plants).

ი ი removed the shoot tip from each seedling and cut out a 10 mm length of shoot

lengths of shoot into each of 5 Petri dishes

 added to each Petri dish an identical volume of 5% olution

added to each Petri dish 40 cm³ of a different concentration of IAA solution left the Petri dishes at 20 °C in the dark with their lids



A student inves (IAA) on the grd (IAA) on the grd The student: removed the a 10 mm leng a 10 mm leng a 10 mm leng placed 10 len e added to eacl glucose solut added to eacl glucose solut edded to eacl concentration eft the Petri (on for 5 days

removed the shoots after 5 days and measured them

determined the mean change in length of shoot at each concentration of IAA.

TABLE 1 shows her results.

TABLE 1

tration etri dish nillion	10-5	10-3	10-1	-	10
ge in noot / mm	0.0	0.1	1.3	2.4	3.1




[Turn over]

IAA concen added to Pe / parts per n Mean chang length of sh

removed t determine each conc TABLE 1 sho

BLANK PAGE






Explain why the student removed the shoot tip from each seedling. [2 marks]








Explain why the student added glucose solution to each Petri dish. [2 marks]






Explain why the lids were kept on the Petri dishes. [2 marks]








Describe and explain the results shown in TABLE 1, on page 19, and suggest how the results might have differed if lengths of ROOT had been used. [3 marks]






The student produced the different concentrations of IAA using a stock 1 g dm⁻³ solution of IAA (1 g dm⁻³ = 1 part per thousand) and distilled water.

Complete TABLE 2 with the volumes of stock IAA solution and distilled water required to produce 40 cm³ of 10 ppm (parts per million) IAA solution. [1 mark]

TABLE 2

Concentration of IAA solution / parts per million	Volume of distilled water / cm ³
10	





0 4

Scientists investigated the effect of a decrease in pH on muscle contraction. The scientists did the investigation with four different preparations of isolated muscle tissue: A, B, C and D.

- A mouse muscle fibres at typical pH of mouse muscle tissue (control 1).
- B mouse muscle fibres at 0.5 pH units below typical pH.
- C rabbit muscle fibres at typical pH of rabbit muscle tissue (control 2).
- D rabbit muscle fibres at 0.5 pH units below typical pH.

They measured the force of muscle contraction of the muscle fibres at 12 °C,

22 °C and 32 °C


FIGURE 3 shows the results the scientists obtained for B and D compared with the appropriate control.



27

FIGURE 3

Force of muscle contraction / percentage of appropriate control



10203040Temperature / °C



BLANK PAGE





A student looked at the results and concluded that a decrease in pH does cause a decrease in the force of muscle contraction.

Use FIGURE 3, on page 27, to evaluate this conclusion. [4 marks]



30



BLANK PAGE





Another group of scientists suggested that a decrease in the force of muscle contraction is caused by an increase in the concentration of inorganic phosphate, Pi, in muscle tissues.

Their hypothesis is that an increase in the concentration of Pi prevents the release of calcium ions within muscle tissues.

Explain how a decrease in the concentration of calcium ions within muscle tissues could cause a decrease in the force of muscle contraction. [3 marks]



33	





In muscles, pyruvate is converted to lactate during prolonged exercise.

Explain why converting pyruvate to lactate allows the continued production of ATP by anaerobic respiration. [2 marks]



[Turn over]





Describe the role of glucagon in gluconeogenesis.

Do NOT include in your answer details on the second messenger model of glucagon action. [2 marks]





The gene that codes for glucagon is 9.531 kilobases in length. The DNA helix makes one complete turn every 10 base pairs. Every complete turn is 3.4 nm in length.

Use this information to calculate the length in micrometres (µm) of the gene for glucagon. Give your answer to 3 significant figures. [2 marks]







38

Metformin is a drug commonly used to treat type II diabetes. Metformin's ability to lower the blood glucose concentration involves a number of mechanisms including:

- increasing a cell's sensitivity to insulin
- inhibiting adenylate cyclase.

0 5.3

Explain how increasing a cell's sensitivity to insulin will lower the blood glucose concentration. [2 marks]







Explain how inhibiting adenylate cyclase may help to lower the blood glucose concentration. [3 marks]





BLANK PAGE



allele for grey body, G, and a recessive allele for black a gene for body colour has a dominant


A gene for eye colour has a dominant allele for red eyes, R, and a recessive allele for white eyes, r, and is located on the X CHROMOSOME.

FIGURE 4, on the opposite page, shows the phenotypes over four generations.

In fruit flies, body, g. 90

of fruit flies














ece of evidence from FIGURE 4 to show that grey body colour is dominant. [1 mark]








evidence from FIGURE 4, to show	body colour is NOT on the X	[2 marks]
piece of e	for body	








red-eyed male Black-bodied, × white-eyed female Grey-bodied,

×



A heterozyg

expected in

Phenotypes of parents:

Genotypes of parents:





of offspring


notypes



[Turn over]

Ratio of phei

Phenotypes



A population of fruit flies contained 64% grey-bodied flies. Use the Hardy–Weinberg equation to calculate the percentage of flies heterozygous for gene G. [2 marks]






%



BLANK PAGE






In photosynthesis, which chemicals are needed for the light-dependent reaction? Tick (✓) ONE box. [1 mark]

Reduced NADP, ADP, Pi, water and oxygen.

NADP, ATP and water.

Reduced NADP, ATP, water and carbon dioxide.



NADP, ADP, Pi and water.





Describe what happens during photoionisation in the light-dependent reaction. [2 marks]



A student obtained a solution of pigments from the leaves of a plant. Then the student used paper chromatography to separate the pigments.


FIGURE 5, on the opposite page, shows the chromatogram produced.

07.3

Explain why the student marked the origin using a pencil rather than using ink. [1 mark]



FIGURE 5





BLANK PAGE





Describe the method the student used to separate the pigments after the solution of pigments had been applied to the origin. [2 marks]



07.5

Calculating the R_f values of the pigments can help to identify each pigment. An R_f value compares the distance the pigment has moved from the origin with the distance the solvent front has moved from the origin.

distance pigment has moved from the origin distance solvent front has moved from the origin

The distance each pigment has moved is measured from the middle of each spot.

Pigment A has an R_f value of 0.95



Use FIGURE 5, on page 55, to calculate the R_f value of pigment C. [1 mark]

R_f value of pigment C =





The pigments in leaves are different colours. Suggest and explain the advantage of having different coloured pigments in leaves. [1 mark]







What is a DNA probe? [2 marks]



DNA probes are used to detect specific base sequences of DNA.

The process is shown in FIGURE 6.

FIGURE 6

A DNA sample is broken down into smaller fragments

The DNA fragments are separated by electrophoresis and are transferred to a nylon membrane

The DNA fragments are treated to form single strands and DNA probes are added

DNA probes allow specific base sequences of DNA to be detected





Describe how the DNA is broken down into smaller fragments. [2 marks]








The DNA on the nylon membrane is treated to form single strands. Explain why. [1 mark]

A scientist used DNA probes and electrophoresis to screen four volunteers for five different viral DNA fragments.

FIGURE 7, on the opposite page, shows the results the scientist obtained. The lanes numbered 2 to 5 represent the four volunteers



FIGURE 7





BLANK PAGE





Lane 1 of FIGURE 7, on page 65, enabled the size of the different viral fragments to be determined.

Suggest and explain how. [2 marks]



The lengths of the viral DNA fragments were:

- 600 base pairs
- 250 base pairs
- 535 base pairs
- 300 base pairs
- 500 base pairs.

08.5

Which volunteers had at least one of the viral DNA fragments with 250 base pairs or 535 base pairs? [1 mark]





BLANK PAGE



09

The sundew is a small flowering plant, growing in wet habitats such as bogs and marshes. The soil in bogs and marshes is acidic and has very low concentrations of some nutrients. The sundew can trap and digest insects.


09.1

Describe how you could estimate the size of a population of sundews in a small marsh. [5 marks]



71





Suggest and explain how digesting insects helps the sundew to grow in soil with very low concentrations of some nutrients. [2 marks]



BLANK PAGE



10

Guillain–Barré syndrome is a rare disease in which the immune system damages the myelin sheath of neurones. Myelin sheath damage can cause a range of symptoms, for 5 example numbness, muscular weakness and muscular paralysis. Sometimes, neurones of the autonomic nervous system are affected, causing heart rate 10 irregularities.

Huntington's disease is a disorder caused when a protein called huntingtin damages the brain. Huntingtin is produced because of a 15 dominant, mutant allele.

The first successful drug trial to
reduce concentrations of huntingtin
in the human brain involved
46 patients. The patients received20
20
the drug for 4 months. The



concentration of huntingtin was
reduced in all the patients. The drug
was injected at the base of the spine
into the cerebrospinal fluid bathing25the brain and spinal cord. The drug
contains single-stranded DNA
molecules. These single-stranded
molecules inhibit the mRNA needed
to produce huntingtin.30

Symptoms of Huntington's disease can start at any time, but usually develop between 30 and 50 years of age. The likelihood and age when symptoms start are linked to the 35 number of CAG base sequence repeats in the gene for Huntington's disease. However, recent studies have suggested that epigenetics may also affect the age when 40

may also affect the age when symptoms first start.





Damage to the myelin sheath of neurones can cause muscular paralysis (lines 4–7).

Explain how. [3 marks]







Sometimes Guillain–Barré syndrome causes heart rate irregularities (lines 8–11).

Suggest and explain why. [3 marks]








The first successful drug trial to reduce concentrations of huntingtin in the brain used single-stranded DNA molecules (lines 26–28).

Suggest and explain how this drug could cause a reduction in the concentration of the protein huntingtin. [3 marks]









Scientists from the first successful drug trial to reduce concentrations of huntingtin (lines 17–21) reported that the drug is not a cure for Huntington's disease.

Suggest TWO reasons why the drug should NOT be considered a cure. Do NOT include repeats of the drug trial in your answer. [2 marks]

1





Suggest TWO reasons why people had the drug injected into the cerebrospinal fluid (lines 23–26) rather than taking a pill containing the drug. [2 marks]

-

1

2





Suggest and explain ONE way epigenetics may affect the age when symptoms of Huntington's disease start. [2 marks]

END OF QUESTIONS





BLANK PAGE



BLANK PAGE

For Examiner's Use	
Question	Mark
1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
TOTAL	

Copyright information

For confidentiality purposes, from the November 2015 examination series, acknowledgements of third-party copyright material are published in a separate booklet rather than including them on the examination paper or support materials. This booklet is published after each examination series and is available for free download from www.aqa.org.uk after the live examination series.

Permission to reproduce all copyright material has been applied for. In some cases,

efforts to contact copyright-holders may have been unsuccessful and AQA will be happy to rectify any omissions of acknowledgements. If you have any queries please contact the Copyright Team, AQA, Stag Hill House, Guildford, GU2 7XJ.

Copyright © 2019 AQA and its licensors. All rights reserved.

IB/M/CD/Jun19/7402/2/E2



