

ADVANCED SUBSIDIARY (AS)
<b>General Certificate of Education</b>
January 2014

Centre Number							
71							
Cano	didate Number						

# **Biology**

Assessment Unit AS 1

assessing

Molecules and Cells

[AB111]





#### TIME

1 hour 30 minutes.

#### **INSTRUCTIONS TO CANDIDATES**

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

Write your answers in the spaces provided in this question paper.

There is an extra lined page at the end of the paper if required.

Answer all eight questions.

You are provided with **Photograph 1.3** for use with Question 3 in this paper. Do not write your answers on this photograph.

#### **INFORMATION FOR CANDIDATES**

The total mark for this paper is 75.

Section A carries 60 marks. Section B carries 15 marks.

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

You are reminded of the need for good English and clear presentation in your answers.

Use accurate scientific terminology in all answers.

You should spend approximately **20 minutes** on Section B.

You are expected to answer Section B in continuous prose.

**Quality of written communication** will be assessed in Section B, and awarded a maximum of 2 marks.

For Examiner's use only				
Question Number	Marks			
1				
2				
3				
4				
5				
6				
7				
8				

Total	
Marks	

## **BLANK PAGE**

Section A	Examiner O
Read the following passage about the structure of DNA and chromosomes, and write the most appropriate word in each blank space to complete the account.	Marks Ren
A DNA molecule consists of many repeating units called	
which are joined by	
reactions to form the sugar-phosphate backbone. The double helix	
consists of two single strands held together	
by bonds between	
nitrogenous bases. Chromosomes are formed when the DNA coils round	
proteins called [5]	

2 The technique of chromatography can be used to separate, and subsequently identify, substances in solution. When setting up a chromatogram, certain procedures are followed.

Examin	er Only
Marks	Remark

(a) Describe **two** procedures, relevant to setting up a chromatogram, that would ensure valid results.

1			


2			
			ı

A chromatogram was prepared to identify four carbohydrates.

The following carbohydrate solutions were applied to the origin:

- Glucose
- Fructose
- Maltose treated with α-glucosidase (hydrolyses glycosidic bonds)
- Sucrose treated with  $\alpha$ -glucosidase (hydrolyses glycosidic bonds)

One solution was added to each of the four lanes on the origin. The resulting chromatogram is shown below.



(b)	Calculate the R <sub>f</sub> value for the carbohydrate in <b>Lane 1</b> . (Show your working.)		Examin Marks	er Only Remark
	Working./			
		[2]		
		[2]		
(c)	Lanes 1 and 2 contain the monosaccharides. Lanes 3 and 4 contain the hydrolysed disaccharides.	1		
	Identify the carbohydrate added to each of the lanes 1–4.			
	1			
	2			
	3			
	4	[3]		
(d)	A starch solution was also treated with $\alpha$ -glucosidase, so that the starch was fully broken down. Give the full name of the monosaccharide which would be present in the resulting solution.			
		[1]		

		raph 1.3 shows six photomicrographs of animal cells at different of mitosis. These are labelled A–F.	Ent Examiner Only  Marks Remark
(a)	(i)	Identify the stages shown in photomicrographs <b>A–D</b> .	
		A	
		B	
		c	
		D	[4]
	(ii)	Identify the structures labelled <b>X</b> and describe fully their role i process of mitosis.	n the
			_ [3]
	(iii)	The photomicrographs <b>A–F</b> do not show the stages of mitosis the correct sequence. Rearrange the letters <b>A–F</b> to show the correct sequence of the stages.	s in
			_ [2]
(b)	(i)	Identify the process that is beginning to happen in <b>E</b> .	
			_ [1]
	(ii)	Outline how this process would be different in a plant cell.	
			[1]

**6** 

## **BLANK PAGE**


(Questions continue overleaf)

4 Mature seeds contain an embryo plant and a store of energy-rich food. In some seeds, such as soybean, the main energy store is lipid. When seeds are planted in springtime they absorb water, which activates enzymes such as lipase.

Examiner Only

Marks Remark

The graph below shows the relative activity of soybean lipase at various temperatures.



(	(a)	) (	i)	Describe	precisely	the	trends	evident	in	this	graph.

			[2

	Using your understanding of enzyme action, explain the trends you have identified.	Examin Marks	er On Rem
	you have identified.		
	[3]		
TI	Assessment and the second seco		
	temperature range for soybean lipase activity, as shown in the		
	oh, is different to that for mammalian lipase. Suggest a reason for difference in temperature range and suggest how this is an		
	antage to the growth of the soybean plants.		
aave	antage to the growth of the soybean plants.		
	[0]		
	[2]		
			f.

The earliest method of producing a genetic fingerprint involved an analysis of RFLPs. In this method, DNA is cut into pieces using enzymes, such as *Eco*RI and *Bam*HI, and the resulting fragments are separated using electrophoresis.

Examiner Only			
Marks	Remark		

(a) (i)	What term is used to describe enzymes such as <i>EcoRI</i> and
	BamHI?


[1]

(ii)	DNA is the substrate of both EcoRI and BamHI. However, the
	active site of the two enzymes is slightly different. Explain the
	effect of this difference.

I	7	
ı	4	

One use of genetic fingerprinting is in paternity investigations. In dog breeding, it can be beneficial to have evidence of a dog's parentage.

One breeder claimed that a litter of puppies had been fathered by an award-winning show-dog. A buyer had cause to doubt this, based on the development of an inherited illness in a puppy that she had bought, and so sought evidence of parentage from the breeder. Genetic fingerprints of the puppy, its mother and the show-dog were produced. The results are shown below.



(a)	what can you conclude about the puppy's parentage from the genetic		ner Only
	fingerprints? Explain your answer.	Marks	Remark
	[2]		
(c)	The breeder insisted that the puppy was from a litter born following a		
` '	mating between the mother and the show-dog. Comment on the		
	relative validity (quality) of the genetic evidence compared to the		
	breeder's evidence.		
	breeder's evidence.		
	reı		
	[3]		
(d)	Enzymes such as <i>EcoRI</i> and <i>BamHI</i> were originally discovered in		
	bacteria, where they are thought to offer protection from bacteriophages.		
	Suggest how these enzymes could protect bacterial cells from infection.		
	[4]		
	[1]		


6 Proteins can be described as organic macromole	cules
--	-------

Examin	er Only
Marks	Remark

1	a`	<b>)</b> Explai	n the tern	n 'organic	macromolecule'.
١	~	, Expidi		n organio	madi di nolo dalo i



**(b)** Gluten is a type of protein found in grains such as wheat, barley and rye. It is a 'composite' protein, made up of two simpler proteins called glutenin and gliadin. The following diagram represents the primary structure of part of a gliadin molecule.



(i) Identify the building blocks of proteins, as represented by circles in the diagram.

\_\_\_\_\_\_[1]

(ii) Identify the type of bond labelled  ${\bf X}.$ 

\_\_\_\_\_[1]

(c) Some people are intolerant of gluten in the diet. This is the basis of **Examiner Only** Marks Remar the condition known as coeliac disease. One of the proteins in gluten, gliadin, triggers an immune response in sufferers which results in damage to the lining of the ileum. This damage is often patchy and does not affect the whole ileum. Sections of the ileum mucosal layer from a normal individual and from a coeliac sufferer are represented in the diagram below. villi coeliac sufferer normal (i) Describe the effect of coeliac disease on the structure of the ileum. \_\_\_\_\_ [1] (ii) Coeliac sufferers are often nutrient-deficient. With reference to the damage caused to the ileum, suggest an explanation for this. \_ [2] (iii) When investigating the ileum of a potential sufferer for signs of coeliac disease, several biopsies (tissue samples) are taken, each from a different region of the ileum. Suggest why.

\_ [1]

## **BLANK PAGE**

		out to investigate the from a sample of live	•	poison	Examiner Only  Marks Remark
centrifu spins a depend	ge. Ground-up tissue t high speeds. The c ling on their size. The	d from the rest of the e is placed in a tube, ell contents are sepa e smaller the organel the other cell contents.	which the centrifurated into 'fraction le, the higher the	ige then ns',	
slo	Circle the name of the organelle fraction which would require the slowest spinning speed in order to separate it from the rest of the cell contents.				
	mitochondria	ribosomes	nuclei	[1]	
abo to t	ove. These mitochon	ria was obtained usin dria were placed in a rith the same water p	buffer which was	sisotonic	
(i)	Explain fully the pu	rpose of the isotonic	buffer.		
				[2]	
Tw	o test tubes ( <b>A</b> and <b>E</b>	<b>3</b> ) were then prepare	d as follows:		
•	1 cm <sup>3</sup> of cyanide so	ria in buffer solution volution was added to buffer was added to	ube <b>A</b>	h tube	
(ii)	Suggest why an ex detailed above.	tra 1 cm <sup>3</sup> of buffer wa	is added to tube <b>E</b>	<b>3</b> , as	
				[1]	

The concentration of oxygen in each tube was then investigated using an oxygen probe. Readings were taken from each tube initially and then every minute for 10 minutes. Oxygen concentration was measured in micromoles ( $\mu$ M).

Examiner Only

Marks Remark

In tube **A**, the concentration of oxygen initially was 520. Subsequent readings were 511, 505, 500, 497, 495 and then five readings of 493.

In tube **B**, the concentration of oxygen initially was 505. Subsequent readings were 475, 444, 415, 386, 355, 324, 304, 297, 292 and 290.

(iii) Construct a table of these results in the space below.

Your table should have a caption and should include appropriate column headings, units and all the data.

[4]

(iv)	Describe the results of the experiment.		Examin Marks	er Only Remark
		[2]		
(v)	Suggest the effect of cyanide on cell function.			
		[4]		
		ניו		
(vi)	Between 7 and 10 minutes, the rate of oxygen consumption in tube <b>B</b> slows significantly. Suggest a reason for this.			
		[1]		

## **Section B**

Examiner Only

Marks Remark

Quality of written communication is awarded a maximum of 2 marks in this section.

8	The cell membrane consists of a phospholipid bilayer with various protein embedded in it. This structure enables different substances to travel through the membrane by either simple diffusion, facilitated diffusion or active transport.				
	(a)	Describe the similarities and differences between simple diffusion, facilitated diffusion and active transport.	[6]		
	(b)	Large and small molecules, as well as ions, must be able to travel through the cell membrane. Explain why the different methods of transport named in part (a) are necessary to allow each of these substances to pass through and also how they may allow the membrane to be selective.	[7]		
	Qua	ality of written communication	[2]		
	(a)	Describe the similarities and differences between simple diffusion, facilitated diffusion and active transport.			

		Marks	Remark
(b)	Large and small molecules, as well as ions, must be able to travel through the cell membrane. Explain why the different methods of transport named in part (a) are necessary to allow each of these substances to pass through and also how they may allow the membrane to be selective.		

	Examin	
	Marks	Remark

Extra lined page	Exar	miner Only	
,	Marks		
	_		
	-		
	-		
	-		
	-		
	-		
	_		
	-		
	-		
	-		
	_		
	-		
	_		
	-		
	-		
THIS IS THE END OF THE QUESTION PAPER			

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA will be happy to rectify any omissions of acknowledgement in future if notified.

# GCE Biology Advanced Subsidiary (AS)

# Assessment Unit AS 1: Molecules and Cells January 2014

Photograph 1.3 (For use with question 3)

