

ADVANCED SUBSIDIARY (AS) General Certificate of Education January 2014

71	
Cano	didate Number

Centre Number

Biology

Assessment Unit AS 2

assessing
Organisms and Biodiversity

[AB121]





TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number in the spaces provided at the top of this page.

Write your answers in the spaces provided in this question paper. There is an extra lined page at the end of the paper if required. Answer **all eight** questions.

INFORMATION FOR CANDIDATES

The total mark for this paper is 75.

Section A carries 60 marks. Section B carries 15 marks. Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question. You are reminded of the need for good English and clear presentation in your answers.

Use accurate scientific terminology in all answers. You should spend approximately **20 minutes** on Section B. You are expected to answer Section B in continuous prose. **Quality of written communication** will be assessed in Section B, and awarded a maximum of 2 marks.

Total	
Marks	

For Examiner's

use only

Marks

Question

Number

2

3

4

5

6


7



BLANK PAGE

Section A	Examiner Marks F
Read the following passage on classification, and write the most appropriate word in the blank spaces to complete the account.	
All organisms can be grouped into one of five kingdoms namely Anima	alia,
Plantae, Fungi, and	
Organisms in four of the kingdoms have cells containing membrane-bo	ound
organelles, and are therefore classified as B	Both
plant and fungal cells are surrounded by a cell wall which, in the kingde	om
Fungi, is composed of The cells in kingdom	
Plantae possess chlorophyll to enable the mo	ode
of nutrition, which is typical of this kingdom.	[5]

2 (a) The oxygen dissociation curve of human haemoglobin at two different pH values is shown in the graph below.



(i) State how haemoglobin's affinity for oxygen is affected by a decrease in pH.

______[1]

(ii) Explain how a decrease in pH is brought about in respiring tissue.

______[1]


(iii) Identify **one** other change, which occurs in respiring tissue, that affects haemoglobin saturation in the same way as a decrease in pH.

______[1]

(b)		e oxygen dissociation curve for myoglobin differs from that of emoglobin.	Examiner Only Marks Remark
	(i)	Sketch the dissociation curve for myoglobin on the following graph, which already shows the dissociation curve for haemoglobin.	
		Haemoglobin Haemoglobin	
		Partial pressure of oxygen/kPa [1]	
	(ii)	The bottlenose dolphin can dive and remain submerged for up to ten minutes. The level of myoglobin in the muscle tissue of diving mammals, such as the bottlenose dolphin, is higher than that of similarly-sized terrestrial (land) mammals. Suggest a reason for this.	
		[2]	

3 A phloem sieve tube element in the vascular bundle of a plant stem, with its associated companion cell, are represented in the diagram below.






(a) On the diagram:

- clearly indicate the position of a sieve plate by labelling with the letter S
- use an arrow, or arrows, to clearly indicate the direction of translocation of organic solutes [2]

(b) The percentage composition of the sieve tube contents in barley plants is shown in the pie chart below.





(i)	Suggest the	identity	of	carbohy	drate A	A.


______[1]

(ii) The amino acids in the sieve tubes have been synthesised by the barley, using nitrate ions which have been taken in from the soil.

Suggest why there are no nitrate ions located in the sieve tubes.

_____ [1]

4 (a) The graph below shows pressure changes which take place in the left side of the heart during one complete cardiac cycle.



(i) State the cause of the increase in ventricular pressure which occurs at 0.1s.

______[1]

(ii) State the effect of the closure of the semi-lunar valves on blood flow, and give the time at which this occurs in the graph above.


Effect of semi-lunar valves closure _____

Time at which this occurs _____ [2]

State how the r	maximum pressı	ure in the riaht v	/entricle diffe	s and	
account for this	difference.	are in the right	CHUIOIC GIIICI	o dila	
				[2]	
				[~]	

(b) It is considered that a diet which is rich in fat can lead to the development of cardiovascular diseases such as coronary heart disease (CHD).

The amount of all types of fat consumed per person and the percentage of total deaths due to coronary heart disease (CHD) were investigated in four countries. The results are shown in the graph below.



			Examin	er Only
(i)	From the graph, identify one piece of evidence in favour of the proposed link between fat consumption and deaths from CHD and one piece of evidence against the existence of such a link.		Marks	Remark
	Evidence in favour:	-		
	Evidence against:	-		
	[2]			

	[1]	

5 A bog in Co. Down is designated as an ASSI. In this bog, the percentage cover of plant species was estimated in 10 randomly placed quadrats. The results are summarised in the table below.

Examiner Only		
Marks	Remark	

Plant species	Total % cover of each species in 10 quadrats (n _i)	n _i (n _i -1)
Bog moss	990	979110
Bog cotton	445	197 580
Ling heather	220	48 180
Cross-leaved heather	330	108 570
Cranberry	240	57 360
Lichen	45	1 980
Bog asphodel	235	54 990
	N = 2505	$\sum n_i(n_i - 1) = 1447770$

(a)	(i)	State what is represented by the letters ASSI.
-----	-----	--

(ii) Calculate the value for Simpson's Index (D) for this bog.

The formula for calculating D is presented as

$$D = \frac{\sum n_i(n_i - 1)}{N(N - 1)}$$

where $n_i =$ the total % cover of each individual species N = the total % cover of organisms of all species (Show your working.)

Angwor	[3]
Allswei	L4.

(b)	(i)	In another bog, located in Co. Armagh, plant species were also sampled and the Simpson's Index (D) was calculated as 0.24.)	Examin Marks	er Only Remark
		Compare the biodiversity of the two bogs and suggest an explanation for the value of the Simpson's Index obtained in the Co. Armagh bog.	e		
			[2]		
	(ii)	Simpson's Index is a measurement of species diversity.			
		Distinguish between the terms 'species diversity' and 'genetic diversity'.			
			[2]		
(c)	foui rela	o distinct species, ling heather and cross-leaved heather, were not in both bogland locations. If these two species were closely sted, state how their DNA and protein molecules would display ilarities.			
	DN	Α			
	Pro	tein molecules			
			[2]		

(d)	Bog cotton was also found in both bogland locations. Bog cotton possesses a hydrophytic adaptation in the form of aerenchyma tissue. This tissue enables the passage of oxygen to the roots from the leaves.	Examin Marks	er Only Remark
	Explain fully why aerenchyma tissue is a useful adaptation for plants in bogland habitats.		
	[3]		

BLANK PAGE

(Questions continue overleaf)

6 An investigation was carried out to show how light intensity affects the rate of carbon dioxide uptake in two different plant species. The results are shown in the table below.

Examin	er Only
Marks	Remark


Light intensity/ Wm ⁻²	Rate of carbon dioxide uptake/ arbitrary units		
WIII -	Species A	Species B	
0	-2.0	-2.0	
20	1.6	-0.3	
40	3.3	1.4	
120	4.8	8.2	
160	5.1	9.3	
240	5.1	9.3	

(a)	Using the most appropriate graphical technique, plot the above data	₹.
	(Use the graph paper opposite.)	[4]

Ìί	Jant species B is grown commercially using artificial light sources. Jsing the graph, determine the light intensity value at which species B will grow most economically. Explain your answer.
_	
_	[2]

(c)	For each plant species, there is a particular light intensity value which
	results in no net exchange of carbon dioxide.

State the term used to describe this value and explain how it arises.	
	_
	_
	_
21)1




(d)	(i)	In its natural habitat, plant species A can be found growing under dense tree cover, but plant species B is confined to bright forest clearings.	Examin Marks	er Only Remark
		Using all the information provided, explain how plant species A is adapted for growing in shady conditions.		
		[2]		
	(ii)	The chloroplasts in plant species A are particularly large. This is an adaptation for the low levels of light intensity that occur in its natural habitat. Suggest how directional selection may have given rise to this adaptation in species A .		
		[_O]		

BLANK PAGE

(Questions continue overleaf)

7 The photograph below shows a male and female stick insect. Adult females lack wings and are larger than the males. Adult males are winged and must release sufficient energy from respiration to facilitate flight.

Examiner Only	
Marks	Remark



© Pascal Goetgheluck / Science Photo Library

Insects do not use blood to transport oxygen and therefore oxygen needs to be carried directly from the atmosphere to the respiring tissues.

There are surface pores along the sides of the stick insect, which enable air to move in and out. These surface pores are connected internally to a series of fine tubes (resembling bronchioles) to allow the passage of oxygen to the respiring cells.

(a)	Explain how the body shape of a stick insect assists the diffusion of oxygen which has entered the fine tubes from the atmosphere.		
	[2]		

(b)	(i)	A student hypothesises that the density of surface pores on the male would be greater than that of the female. Suggest the reasoning behind this hypothesis.	Examine Marks	r Only Remark
		[2]		
	(ii)	Another student puts forward a different hypothesis, proposing that the density of surface pores on the male would not differ greatly from that of the female. Suggest the reasoning behind this alternative hypothesis.		
		[2]		
(c)	the	stick insects, the junction between the ends of the fine tubes and respiring tissue is filled with a small amount of fluid. Explain the ction of the fluid in this particular location.		
		[2]		

(d) The diagram below shows a respirometer which is being used to monitor the exchange of gases between a stick insect and its environment. Tube A contains the stick insect and tube B is acting as a thermobarometer (compensates for changes in temperature and pressure).






Image taken from www.practicalbiology.org and reproduced with permission from the Nuffield Foundation and the Society of Biology.

Explain how the apparatus shown would be used to determine the rate of oxygen uptake by the stick insect.			

[4]

Section B

Examiner Only

Marks Remark

Quality of written communication is awarded a maximum of 2 marks in this section.

n intensive agriculture, farmers maximise both the area productivity of the land which they farm. However, such been associated with a reduction in biodiversity.	
Describe the range of practices used in intensive agricunow they can reduce biodiversity on farms.	lture, and explain [13]
Quality of written communication	[2]

i	Marks	Remark

	Examiner Only	
	Marks	Remark
.		
.		
.		
.		
.		
.		
.		
.		

Extra lined page	Examii Marks	ner Only Remark
	Warks	Remark
	-	
	-	
	_	
	_	
	•	
	-	
	_	
	_	
	_	

Permission to reproduce all copyright material has been applied for. In some cases, efforts to contact copyright holders may have been unsuccessful and CCEA will be happy to rectify any omissions of acknowledgement in future if notified.