



**ADVANCED  
General Certificate of Education  
January 2010**

---

**Mathematics**  
**Assessment Unit F2**  
*assessing*  
**Module FP2: Further Pure Mathematics 2**

**[AMF21]**



**WEDNESDAY 3 FEBRUARY, MORNING**

---

**TIME**

1 hour 30 minutes.

**INSTRUCTIONS TO CANDIDATES**

Write your Centre Number and Candidate Number on the Answer Booklet provided.  
Answer **all six** questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

You are permitted to use a graphic or scientific calculator in this paper.

**INFORMATION FOR CANDIDATES**

The total mark for this paper is 75

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

A copy of the **Mathematical Formulae and Tables booklet** is provided.

Throughout the paper the logarithmic notation used is  $\ln z$  where it is noted that  
 $\ln z \equiv \log_e z$

**Answer all six questions.**

**Show clearly the full development of your answers.**

**Answers should be given to three significant figures unless otherwise stated.**

- 1 Find, in radians, the general solution of the equation

$$4 \sin x \cos x + 2 \cos x - 2 \sin x - 1 = 0$$

[7]

- 2 Use mathematical induction to prove that

$$1^3 + 2^3 + \dots + n^3 = \frac{1}{4}n^2(n+1)^2$$

for all positive integers  $n$ .

[7]

- 3 (i) Express in partial fractions

$$\frac{1+x}{(x^2+1)(1-x)}$$

[6]

- (ii) Hence, or otherwise, find the general solution to the differential equation

$$\frac{dy}{dx} - \frac{2x}{x^2+1}y = \frac{1+x}{1-x}$$

[8]

- (iii) Given that  $y = 3$  when  $x = 0$ , find the particular solution.

[2]

- 4 (a) Illustrate on an Argand diagram the roots of the equation

$$z^8 - (3e^{i\frac{\pi}{10}})^8 = 0$$

[3]

- (b) Let  $z = \cos \theta + i \sin \theta$  be a complex number.

- (i) Show that

$$\frac{1}{2}(z + z^{-1}) = \cos \theta$$

[2]

- (ii) Hence, find an expression for  $\cos^6 \theta$  in the form

$$\cos^6 \theta = a \cos 6\theta + b \cos 4\theta + c \cos 2\theta + d$$

where  $a, b, c$  and  $d$  are rational numbers which are to be determined.

[6]

- (iii) Hence find

$$\int \cos^6 \theta \, d\theta$$

[2]

- 5 (i) Use Maclaurin's theorem to derive the series expansion for  $\cos x$  up to and including the term in  $x^4$

[5]

- (ii) Hence show that the series expansion of

$$(1 - x^2)^{-\frac{1}{2}} \cos kx$$

is  $1 + (\frac{1}{2} - \frac{1}{2}k^2)x^2 + (\frac{1}{24}k^4 - \frac{1}{4}k^2 + \frac{3}{8})x^4 + \dots$

where  $k$  is a real number.

[8]

- (iii) If this series expansion takes the form  $1 + px^4 + \dots$ , find the values of  $k$  and  $p$ .

[3]




Fig. 1

**Fig. 1** above shows part of an ellipse centred at the origin O.

The point  $(12, 0)$  is a focus, and the line  $x = \frac{169}{12}$  is a directrix of the ellipse.

- (i) Show that the equation of the ellipse is

$$\frac{x^2}{169} + \frac{y^2}{25} = 1 \quad [5]$$

P is a point on the ellipse with y coordinate 3

- (ii) Show that the equation of the tangent to the ellipse at P is

$$39y + 20x = 325 \quad [8]$$

T is the point where the tangent at P cuts the x-axis and N is the foot of the perpendicular from P to the x-axis.

- (iii) Show that  $OT \times ON$  equals the square of the length of the semi-major axis. [3]