



**ADVANCED
General Certificate of Education
January 2013**

**Mathematics
Assessment Unit C4
assessing
Module C4: Core Mathematics 4
[AMC41]**



FRIDAY 25 JANUARY, AFTERNOON

TIME

1 hour 30 minutes.

INSTRUCTIONS TO CANDIDATES

Write your Centre Number and Candidate Number on the Answer Booklet provided.
Answer **all eight** questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

You are permitted to use a graphic or scientific calculator in this paper.

INFORMATION FOR CANDIDATES

The total mark for this paper is 75

Figures in brackets printed down the right-hand side of pages indicate the marks awarded to each question or part question.

A copy of the **Mathematical Formulae and Tables booklet** is provided.

Throughout the paper the logarithmic notation used is $\ln z$ where it is noted that $\ln z \equiv \log_e z$

Answer all eight questions.

Show clearly the full development of your answers.

Answers should be given to three significant figures unless otherwise stated.

- 1** Find the coordinates of the point on the curve given parametrically by

$$x = t^2 + 1 \quad y = 2t^3$$

at which the gradient is 6

[6]

- 2** The vertices of a triangle ABC are: A (2, 4, 5)
B (3, 3, -1)
C (0, 1, 4)

Find:

(i) \vec{BA} and \vec{BC}

[3]

(ii) $|\vec{BA}|$ and $|\vec{BC}|$

[3]

(iii) the angle ABC.

[3]

- 3 (i)** Rewrite

$$5 \sin x - 12 \cos x$$

in the form $R \sin(x - \alpha)$ where R is an integer and $0^\circ \leq \alpha \leq 90^\circ$

[3]

(ii) Hence solve the equation

$$5 \sin x - 12 \cos x = 7 \quad 0^\circ \leq x \leq 360^\circ$$

[4]

4 (i) Given that

$$(x + y)^3 = 2x$$

use implicit differentiation to show that

$$\frac{dy}{dx} = \frac{2}{3(x+y)^2} - 1$$

[4]

(ii) Hence find the equation of the tangent to the curve

$$(x + y)^3 = 2x$$

at the point (4, -2) in the form $ax + by + c = 0$, where a , b and c are integers.

[4]

5 (a) Find

$$\int x \cos 2x \, dx$$

[6]

(b) Use the substitution $u = \cot x$ to find

$$\int (1 + \cot^2 x) e^{\cot x} \, dx$$

[7]

6 A function f is defined by

$$f(x) = x^2 - 6x + 5 \quad x \in \mathbb{R}$$

(i) Express $f(x)$ in the form

$$(x - a)^2 - b$$

[3]

(ii) Hence state the range of the function $f(x)$.

[1]

(iii) State the largest possible domain so that the function $f(x)$ is increasing.

[1]

(iv) Find the inverse function $f^{-1}(x)$, for the 1 – 1 function in part (iii), stating its domain. [5]

7 A radioactive isotope decays at a rate proportional to the amount, A , present at time t years.

- (i) Form a differential equation to model this.

[2]

The half-life of the isotope is 250 years.

- (ii) Find the percentage of the original amount that remains after 100 years.

[10]

8 Fig. 1 below shows part of the graph of

$$y = \sin x + \cos x$$




Fig. 1

A glass lampshade is formed by rotating the area enclosed by the curve, the positive x -axis and the y -axis through 2π radians about the x -axis.

Find the **exact** volume enclosed by the lampshade.

[10]