CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2013 series

9184 BIOLOGY (US)

9184/35   Paper 35 (Advanced Practical Skills 1),
maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

1 (a) (i)

<table>
<tr><td rowspan="4" style="writing-mode: vertical-lr">MMO decisions 2</td><td>mp1</td><td>(shows correct use of 20 % glucose)
adds 6, 8, and 10 cm^3 of 20 % glucose to prepare correct final concentration of glucose;</td></tr>
</table>

volume of 20 % glucose solution / cm^3	final percentage concentration of glucose
6	6
8	8
10	10

mp2 — (shows the correct use of distilled water)
adds 14, 12 and 10 cm^3 of (distilled) water/W to prepare correct final concentration of glucose;

volume of distilled water / cm^3	final percentage concentration of glucose
14	6
12	8
10	10

(ii) [1]

MMO decisions 1	1	sec (onds) or s;

(iii)			
MMO decisions 1	1	volume of **S** or **PM** or **U1** or **U2** or (glucose) solution OR 5 $\underline{cm}^{3}$ **S** or 2 cm^{3} **PM** or 10 cm^{3} **U1 / U2** / solutions ;	
(iv)			[4]
ACE improvements 1	mp1	use <u>syringe</u> ;	
MMO collection 1	mp2	records start and end-point times for **U1** and **U2**;	
MMO decisions 1	mp3	start time for **U1** is before **U2**;	
PDO recording 1	mp4	all readings to the same precision;	

	(v)		
ACE interpretation 1	1	time taken for the 6 % glucose solution to reach end-point in <u>sec(onds)</u> or <u>s</u> ;	

	(vi)		[4]
PDO recording 2	mp1	table with all cells drawn	**AND** heading (top or left) <u>percent(age) conc(entration) of glucose</u>;
		Can have • no outer boundary • % • test tube/ additional columns or rows • notes outside the area	**Do not give mark if** • units in cells of headed column/row • other units e.g. mol dm^{-3} • no units
	mp2	(heading) <u>time</u> (/) s or sec(onds);	
MMO collection 2	mp3	for 6 concentrations including **U1** and **U2** records <u>only</u> processed results as <u>whole numbers</u>;	
	mp4	highest concentration of glucose solution is shortest time compared to 6 %, 8 % and 10 % glucose solution;	

(b)				
ACE interpretation 1	1	glucose tolerance test solution;		

[1]

(c) (i)

			cause of error	WITH idea of error
ACE interpretation max 1				
		mp1	(dependent) change to colourless/end-point	difficult to judge see or identify or determine or is subjective may be different;
		mp2	(standardised) measuring the potassium manganate (**PM**)	difficult to read the syringe due to darkness of **PM**;
		mp3	(standardised) mixing of **S** and glucose	not the same or varies or different;
		mp4	(idea of) reaction	too quick or describes more concentrated solution goes too quickly;

(ii) [max 2]

		mp1	(independent variable) (concentration of glucose) *idea of* use more or different or wider/narrower range of concentrations;
ACE improvements max 2	max 2	mp2	repeat more than once/replicates to obtain three readings;
		mp3	use a colorimeter or *idea of* individual timing/ set each one up separately;

[Total: 17]

2 (a) (i)

<table>
<tr>
<td rowspan="4" style="writing-mode: vertical-lr;">MMO collection 2</td>
<td>mp1</td>
<td colspan="4">(collects correct values for each tissue, J, K, L and M)</td>
</tr>
<tr>
<td></td>
<td>J</td>
<td>M</td>
<td>N</td>
<td>P</td>
</tr>
<tr>
<td></td>
<td>Ignore</td>
<td>4–6</td>
<td>26–28</td>
<td>4–6;</td>
</tr>
<tr>
<td>mp2</td>
<td colspan="4">any four values which add up to 43;</td>
</tr>
</table>

(ii) [3]

Ignore any labels / label lines / brackets

PDO Layout 1	mp1	suitable plan diagram	**AND** clear, sharp, unbroken lines;
		Do not give mark if • any line outside the two given drawn lines left and right • any line off the grid • any shading • any ruled line	**Do not give mark if** • less than 3 lines **or if any line** • any part of the line 1 mm or thicker • any feathery or broken or dashed line or gap • any tail or overlap

PDO recording 1	mp2	any line completing the top edge of the vascular bundle between the two drawn lines;

MMO decision 1	mp3	(draws correct proportions and shape of layers) width of layers **N** is at least double the combined width of **J** and **M** measured along line;

(iii) [2]

PDO display 2	mp1	shows counting of all the 1 cm × 1 cm squares half or more only within the completed outline on Fig. 2.2;	
	mp2	number clearly linked to bundle **and** number clearly linked to xylem	**AND** larger whole number to smaller whole number or as fraction larger number over smaller number;

		(b)	
PDO layout 1	mp1	suitable drawing;	
		Do not give mark if • any shading • draw over the print of the question • any ruled lines • less than 40 mm at widest distance across largest cell • less than six enclosed areas **or if any outer lines (of enclosure)** • are thicker than 1 mm • are feathery or broken / dashed or have gaps • have tails or overlaps	
MMO collection 3	mp2	only <u>6</u> complete cells drawn **AND** as two groups of 3 touching cells;	
	mp3	the largest dimension of the biggest cell from near the centre of the stem is at least 3 times the smallest dimension of the smallest cell from the corner *;*	
	mp4	in one group of three cells, all cells must be drawn with double lines all the way round **AND** where two pairs of cells touch there must be 3 lines (representing the middle lamella);	
MMO decision 1	mp5	<u>one</u> correct label <u>cell wall</u>, with <u>one</u> label line which must touch outermost line of a cell or finish between the two cell wall lines;	

(c)

PDO recording 1	1	organise as a table with only three columns or rows separated by lines (no cells needed)	**AND** headed <u>L1</u> and <u>Fig. 2.3</u>	**AND** third column contains features;

Additional guidance: column/row with features can be left, right or in middle and does not require heading <u>J1 Fig. 2.3</u>

<table>
<tr><td rowspan="6">ACE interpretation max 3</td><td rowspan="6">max 3</td><td></td><td>feature</td><td>L1</td><td>Fig.2.3</td></tr>
<tr><td>mp1</td><td>vascular bundles tissue/xylem</td><td>fewer/few separate bundles/ring/in the corners</td><td>many/more ring/separate bundles;</td></tr>
<tr><td>mp2</td><td>hollow centre/pith (some slides)</td><td>present or has or yes</td><td>absent or does not have or no(ne);</td></tr>
<tr><td>mp3</td><td>thickening/collenchyma sclerenchyma/ fibres/described</td><td>present or has or yes or in corner or forms bumps</td><td>absent or does not have or no(ne);</td></tr>
<tr><td>mp4</td><td>Idea of gaps/air spaces</td><td>absent or does not have or no(ne)</td><td>present or has or yes;</td></tr>
<tr><td>mp5</td><td>outer layer(s)/cortex/ epidermis</td><td>thick(er) or idea of regular cells</td><td>thin(er) or idea of irregular cells;</td></tr>
</table>

(d) (i)

if line graph drawn award only mp1

<table>
<tr>
<td rowspan="4">PDO layout 4</td>
<td>mp1</td>
<td>

x-axis

 type of plant tissue
</td>
<td>

AND *y*-axis

concentration of glucose arbitrary units or [glucose] arbitrary units;
</td>
</tr>
<tr>
<td>mp2</td>
<td>even width of blocks</td>
<td>

AND (0 at origin)
1.0 a.u. to 1 cm labelled 2.0, 4.0, 6.0
(ignore 0.0 at the origin or 8.0);
</td>
</tr>
<tr>
<td>mp3</td>
<td colspan="2">

correct plotting of each blocks in the order in the table with a
- horizontal,
- ruled,
- even line, less than 1 mm;

A	2.0
B	6.5
C	4.2
D	5.6
E	3.2

</td>
</tr>
<tr>
<td>mp4</td>
<td>

all blocks separated by a space (could be uneven)
quality – vertical lines
- ruled,
- smooth line less than 1 mm
- meets horizontal line exactly
</td>
<td>

AND
labelled with any clear labels A,B,C,D,E
e.g. must be directly below correct bar or inside bar or shaded with key;
</td>
</tr>
</table>

		(ii)	
ACE interpretation 1	mp1	*idea of* concentration of glucose (inside the cells) is different/not the same (for each type of plant tissue) or gives figure for each plant tissue or compares two different tissues using figures or concentration of glucose is highest in cells in plant tissue B and lowest in plant tissue A;	
ACE conclusion 2	mp2	glucose is absorbed against a concentration or diffusion gradient/glucose did not move out of cells;	
	mp3	<u>active transport</u> or no <u>diffusion</u> or <u>diffusion</u> prevented;	
			[Total: 23]