CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2013 series

9184 BIOLOGY (US)

9184/23

Paper 2 (AS Structured Questions), maximum raw mark 60

www.PapaCambridge.com

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabu
	GCE AS/A LEVEL – May/June 2013	9184
Mark scheme; / R A AW underline	e abbreviations: separates marking points alternative answers for the same point reject accept (for answers correctly cued by the question, or by exi alternative wording (where responses vary more than usual) actual word given must be used by candidate (grammatical)	

Mark scheme abbreviations:

max indicates the maximum number of marks that can be given

or reverse argument ora

marking point (with relevant number) mp

error carried forward ecf

ignore

Page	e 3	Mark Scheme	Syllabu
		GCE AS/A LEVEL – May/June 2013	9184
E	3 = (inter	roplast ; A granum / grana rcellular) air space ; eolus ; A nucleus	Cambridge Con
(b) <i>t</i>	wo mark	rs for correct answer	

 $9(\mu m);;$

award one mark if not to nearest micrometre, 9.3/9.28 or correct measurement incorrectly converted but correct formula used 64/65/66 mm ÷ 7000

[2]

- (c) 1 phospholipid bilayer;
 - A lipid bilayer
 - 2 phospholipids have, phosphate / hydrophilic, heads, and, hydrophobic tails / fatty acid chains;
 - 3 ref. to, labile nature of bilayer structure / phospholipid (molecules) moving (within their monolayer);
 - 4 protein molecules, interspersed/scattered/not a complete layer/AW;
 - 5 many/AW, different/AW (protein molecules);
 - 6 example of type of protein;
 - 7 idea of (most) proteins, moving/not in fixed position;
 - 8 reference to cholesterol;

[max 4]

- (d) 1 ref. to movement, down water potential gradient/from high(er) to low(er);
 - 2 apoplastic/cell wall, pathway from xylem to cell walls of (palisade mesophyll) cells;
 - 3 ref. to osmosis; in context of movement, into cell/through cell surface membrane/through tonoplast,

R osmosis from xylem to vacuole

- 4 symplastic/cytoplasmic, pathway (within cell);
- 5 via plasmodesmata; in context of water arriving from adjacent cell
- 6 ref. to channel proteins/aquaporins;
- 7 solutes/named, in vacuole;

[max 3]

[Total: 12]

Page 4	Mark Scheme	Syllabu	er
	GCE AS/A LEVEL – May/June 2013	9184	900

2 (a) (i) cells have machinery for protein synthesis/AW;

A plant/animal, cells have RER

(assumption that) cells will continue to produce protein at high rate;

large number of/many/AW, ribosomes;

available supply of/AW, amino acids;

ref. to presence of tRNA molecules;

ATP available;

ref. to easier to harvest high levels of protein;

[max 2]

(ii) idea that any added mRNA, has easier access to/can reach, ribosomes/RER; so that the cell's own, DNA/mRNA can be accessed/AW; easier to, harvest/extract, protein products;

[max 1]

- (iii) only the desired protein is produced/AW; ora unwanted protein does not have to be separated from desired protein; idea that inefficient process if translation machinery used to produce other proteins; cell's proteins may, inhibit/affect/hinder/AW, process; [max 1]
- (iv) ref. to ribosome function not altered;
 R ref. to prokaryotic and eukaryotic ribosomes being the same mechanism of translation/described, is the same in all cells;
 e.g. tRNA can respond to introduced mRNA

all types of cells use mRNA for protein synthesis;

mRNA only has one role;

genetic code/codons, are the same in all cells;

A genetic code is universal

mRNA, contains only exons/introns removed, so translation can occur;

(b) different, structure/rRNA, (of ribosomes); (ribosomes), larger/80S, in eukaryotes or smaller/70S in prokaryotes; (some) attached to/AW, (eternal surface of) RER in eukaryotes; ora A only found in cytoplasm in prokaryotes

[max 2]

[max 2]

(c) other organelles/components, damaged or whole cell all organelles intact; some, ribosomes/RER, lost/damaged; idea that cell-free system is disorganised; ora fewer amino acids available; ora no/reduced, respiration; AW other, components/AW, required are, lost/at lower levels; organelles/components, not replaced; ora ref. to difficulty in creating identical conditions to cell environment; may be able to use cells that can replicate (hence continuous production); AVP;

[max 1]

[Total: 9]

Page 5		Mark Scheme	Syllabu
		GCE AS/A LEVEL – May/June 2013	9184
3	 (a) (i) A = denitrification/reduction; B = nitrogen fixation/lightning; C = nitrification/oxidation 		Cambridge
	A ı sa	composition/decay; reference to decomposers protrophs/bacteria/fungi;	COM

A detritivores

digest/breakdown/hydrolyse, organic nitrogen/protein/amino acids/urea;

protease/urease:

deamination;

production of ammonium (ions)/ammonification;

nitrification/ammonium (ions) to nitrate (ions);

accept correct formulae for ammonium ions, nitrite ions and nitrate ions

[max 3]

(b) (i) phosphate

any one relevant;

e.g. part of structure of

AMP/ADP/ATP

nucleotide

nucleic acid/DNA/RNA/polynucleotide

phospholipid

A phospholipid bilayer

phosphorylation/enzyme activation

bone tissue

nitrate

any one relevant;

e.g. (nitrogen for) amino acids/proteins/enzymes/named (e.g. haemoglobin)

AMP/ADP/ATP

nucleotide

nucleic acid/DNA/RNA/polynucleotide named nitrogen base (adenine/cytosine/

thymine/uracil/quanine)

(some) phospholipids

[2]

- growth linked to, increase in cell size/cell number; (ii) 1
 - growth linked to, increase in biomass/reproduction;
 - increases energy available as food for next trophic level;
 - nitrogen is in, amino acids/proteins, for growth;
 - ATP (containing phosphate) required for, transcription/protein synthesis/enzyme synthesis/anabolic reactions/growth;
 - idea that, growth of cells/cell division, requires membrane synthesis; 6
 - nitrogen in membrane proteins/phosphate in membrane phospholipids;
 - (cell division and), DNA, synthesis/replication;
 - idea that more biomass (per unit time) returned to soil;
 - **10** AVP; e.g. ref. to phosphate taken up by plants and passed into food chain; [max 3]

[Total: 11]

			32
	Page 6	Mark Scheme	Syllabu
	_	GCE AS/A LEVEL – May/June 2013	9184
4	(a) any one valid;		Canno

e.g.(first) appearance of (brown) colour use of, colour standards/colour charts use of colorimeter time-lapse photography/video

(b) allow catechol for substrate throughout rate of reaction 0 au, no substrate to act on / AW;

at substrate concentrations lower than 5mM

substrate (concentration) is limiting (factor in rate of reaction);

presence of free active sites/enzyme is in excess;

few collisions between enzyme and substrate;

rate increases with substrate concentration as more, active sites can be occupied/E-S complexes can form;

one data quote to support response

V_{max} reached/rate becomes maximum, at 4.5–5 mM substrate concentration; rate constant/levels out/AW, from 4.5–5 mmol substrate concentration;

at substrate concentrations greater than 5 mM enzyme (concentration) becomes limiting (factor);

all active sites, saturated/occupied;

(so) further increase in substrate concentation does not increase rate;

[max 5]

(c) (i) curve always lower than that with no inhibitor; must be similar shape curve reaches the maximum; A curve heading to maximum

[2]

(ii) PHBA/ inhibitor, similar shape to, substrate/ catechol; (so) binds to active site:

blocks access to substrate/fewer (successful) enzyme-substrate collisions; reduces rate of, reaction/conversion of substrate to product; AVP;

e.g. inhibitor has a greater effect on rate at lower substrate concentrations V_{max} reached at higher substrate concentrations inhibitor forms same interactions with R-groups in active site

[max 2]

(d) enzymes work in a limited pH range / either side of optimum pH rate decreases; (acid so) presence of H⁺ ions, partially denatures / denatures (some), enzymes; further detail; e.g. ref. to breaking ionic or hyrdrogen bonds change of active site shape means substrate can no longer fit; AVP; e.g. ref. to antioxidant effect of, lemon juice / citric acid / vitamin C

[max 2]

[Total: 12]

Pa	age 7	Mark Scheme	Syllabu er
		GCE AS/A LEVEL – May/June 2013	9184
(a)	production replacing repair (o R repair asexual	by increase in cell number); on of genetically identical cells; g (damaged) cells; f tissue); allow 'regeneration' if mp3 and mp4 not awarde cells reproduction; g A vegetative propagation	Syllabut dibbacambhig
(b)	one tick	in each box;	[1]
(c)	chromos spindle f centriole R sides/	nce of chromosomes/condensation of chromatin/AW; comes visible as two, sister chromatids/chromatids joined cormation/spindle fibres form/microtubules assemble/AW s, move to/reach, opposite <u>poles</u> ; ends arance of nucleolus;	
		nbly/breakdown of, nuclear envelope; ar membrane	[max 4]
(d)	uncontro ignore (prophase, will begin again, too soon/immediately; olled/repeated, cell division/mitosis; risk of), tumour formation/cancerous growth onsequences on the timing of the cell cycle;	[max 3]
			[Total: 11]
(a)	result of	mainly) linked to habits during life of person/AW; choices made by person/AW;	
	example e.g. (tob	acco) smoking/exposure to asbestos at work	[max 2]
(b)	transmis	Morbilli/measles sion: aerosol, infection/transmission or droplet, infection/ol route described from infected to uninfected	transmission ; [2]
(c)	(infected), visitors/immigrants/returning residents, from count	ries where measles

(c) (infected), visitors/immigrants/returning residents, from countries where measles occurs;
unvescipated people returning from travel abroad;

unvaccinated people returning from travel abroad;

reduction in vaccination rates;

percentage cover too low;

change in reporting pattern;

existing vaccine no longer, effective/AW;

AVP;

e.g. mutation of virus produces new antigens, vaccines ineffective increase in malnourished children, no immune response to vaccine

[max 1]

[Total: 5]