#### UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced/Advanced Subsidiary Level

# MARK SCHEME for the May/June 2006 question paper

# 9700 BIOLOGY

9700/06

Paper 6

Maximum raw mark 40

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published Report on the Examination.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the Report on the Examination.

The minimum marks in these components needed for various grades were previously published with these mark schemes, but are now instead included in the Report on the Examination for this session.

CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2006 question papers for most IGCSE and GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

www.PapaCambridge.com

<u> </u>	age	1	Mark Scheme Sylla	er
			GCE A/AS LEVEL – May/June 2006 9700	30
			OPTION 1: Mammalian Physiology:	and
(	(a)	nerv to b nerv radi	Mark Scheme  GCE A/AS LEVEL – May/June 2006  OPTION 1: Mammalian Physiology:  t sensed by, retina / cones / rods; we impulses along optic nerve; rain; we impulses to muscles in Iris; al muscles contract; automatic nervous system;	max 4
(	(b)	(i)	as time increases number of correct positives decreases; non-linear relationship / words to that effect; use of correct manipulated figures	max 2
		(ii)	blood alchohol concentration is going down; as liver breaks down alchohol so fewer people will be over the limit / have more than 0.08% blood concentration;	alchoho max 2
		(iii)	many false negatives even at time 0; when all subjects were expected to have more than 0.08% blood concentration; some false positives at all times; could lead to people being wrongly, convicted / words to that effect; ref. figures.	alchoho max 2
(	(c)	(i)	alchohol dehydrogenase; converts (ethanol) to, ethanal / acetaldehyde; aldehyde dehydrogenase; converts, ethanal / acetaldehyde, to, ethanoate / acetate; hydrogens picked up by NAD / NAD is coenzyme / NAD is reduced;	max 3
		(ii)	fatty acids accumulate; fats deposited (in liver) / fatty liver; alchohol, toxic to / kills hepatocytes; fibrous tissue, builds up/replaces hepatocytes; blood supply reduced;	
			cirrhosis;	max 2
				Total: 15
	/ _ <b>\</b>	ono	mark for each correct label;	2

(c) (i) ref. osmosis;

of water / ions;

water potential in lumen is lowered; below that of the, cells / intestine wall;

water moves down water potential gradient; max 3

(ii) difference is 14.2-7.5=6.7; so percentage change is  $(6.7 \div 14.2) \times 100 = 47.2 \%$ ; or  $(6.7 \div 7.5) \times 100 = 89.3\%$ 

2

max 2

(iii) water has left the blood:

Page 2	Mark Scheme	Sylia	er
	GCE A/AS LEVEL – May/June 2006	9700	

plasma proteins do not leave blood; ref. more antibodies:

Total:

3 (a) (i) lens cannot (easily) change shape;

therefore poor, accommodation / focussing at different distances;

lens does not take up rounded shape when tension relaxed / ciliary muscle contracted.

So difficult to focus on near objects;

max 3

(ii) light cannot pass through lens; vision is clouded;

2

(b) Cloudy area of lens / whole lens, is removed; can be replaced with artificial lens / patient wears glasses; detail;

max 2

Total: 7

4 (a) (i) A centrum and B neural spine;

1

(ii) X plus reason
Position of neural spines, position of aorta, position of ribs

1

**(b)** thoracic has longer neural spines than lumbar; ref. to muscle attachment;

thoracic has extra articulating surfaces; for ribs;

lumbar has, larger / thicker, centrum; extra load-bearing / stronger muscles;

lumbar has heavier transverse processes; for attachment of stronger muscles;

max 4

(c) bone / vertebrae, loses calcium; bone / vertebrae, loses, bulk / strength; vertebrae become smaller;

max 2

Page 3	Mark Scheme Sylla er
	GCE A/AS LEVEL – May/June 2006 9700
	OPTION 2: Microorganisms and Biotechnology
(a) (i)	1cm³ of effluent added to 9cm³ sterile water (gives 10 <sup>-1</sup> );
	1cm³ of the first dilution removed and added to 9cm³ sterile water (gives 10 <sup>-2</sup> );
	repeat precedure with second and subsequent dilutions to obtain the range required: 3

# **OPTION 2: Microorganisms and Biotechnology**

repeat procedure with second and subsequent dilutions to obtain the range required; 3

- (ii) 10<sup>-4</sup>; 1
  - 10<sup>-3</sup> too many to count accurately as colonies overlap; 10<sup>-5</sup> too few as sampling errors in dilutions are very great;
- (iii) 55 in 0.1cm<sup>3</sup> = 550 per cm<sup>3</sup> = 5.5 x  $10^2$ ;

dilution is  $10^4 = 5.5 \times 10^6 \, \text{per cm}^3$ allow ecf from (ii)

2

2

- (b) activated sludge / trickling filter / description; aerobic bacteria digest organic matter; aerobic bacteria respire / metabolise / AW, organic matter; insect larvae / protoctista, feed on bacteria insect larvae / protoctista (protozoa), form a layer on surface of stones;
- max 3

(c) saprophytic / putrefying, bacteria digest protein to amino acids;

amino acids deaminated releasing ammonia;

ammonium compounds / urea, acted on by, nitrifying bacteria / named example;

ammonium converted to nitrite;

nitrite converted to nitrate;

max 4

Γ	Page	4	Mark Scheme	Sylla	er
			GCE A/AS LEVEL – May/June 2006	9700	000
2	(a)	(i)	Cut with endonucleases; separated by size using electrophoresis;		DaCambril
		(ii)	Mass of, disorganised / undifferentiated / unspecialised, plant containing DNA from two different sources	ells;	1
		(iii)	To enable selection of the transgenic cells; only the cells with the new DNA can grow in the presence of the	e antibiotic;	2
	(b)	(i)	may help to prevent night blindness; enable local farmers to grow a cash crop; enable the development of rice breeding to improve local crops	;	max 1
		(ii)	May help reduce the risk of cancer;		1
					Total: 8
3	(a)	(i)	Inject antigen into mouse; Extract, blood / spleen, containing lymphocytes; centrifuge to separate lymphocytes;		max 2
		(ii)	It causes the lymphocytes to divide faster than normal; It gives "immortality" / cells survive indefinitely		max 1
	(b)	(i)	Cancer changes the antigens on cells; Antibodies are specific so only cancer cells are affected;		2
		(ii)	The antibiotics are delivered directly to the infected cells;		1
			If present the HIV antibodies in serum bind to HIV antigens;		
			Anti – HIV antibody binds to the HIV antibodies;		
			Enzyme attached to the anti-HIV antibody catalyses reaction product;	on to give a	coloured 3
					[Total 9]

Page	e 5	Mark Scheme	Sylla
		GCE A/AS LEVEL – May/June 2006	9700
(a)	B cell C cyt D DN	embrane; I wall; coplasm; IA / nucleic acid; accept: chromosome rk for 2 correct, rounded up.	Cambridge Com
(b)	(Gran	n positive)	

(a) A membrane;

walls are thicker;

have more, peptidoglycan / murein;

more rigid;

no outer membrane;

no lipid / no polysaccharide;

stain is taken up more easily;

Allow ora. max 3

- (c) Up to 220 slow growth of population after air supplied the population increases rapidly;
- 1 (d) (i) To prevent the entry of the other microorganisms / AW;
  - (ii) The air rising to the top of the fermenter will carry materials from the bottom; 1

Page	e 6	Mark Scheme	Sylla
		GCE A/AS LEVEL – May/June 2006	9700
		OPTION 3 – Growth, Development and Re	eproduction
(a)	(i)	0 to 4 days, decreases; 4-30 days increases; 30-35 days, no change;	S.COM

## **OPTION 3 – Growth, Development and Reproduction**

4-30 days increases; 30-35 days, no change; after 35 days, decreases; ref. figs.;

max 3

(ii) 0 to 4 days

uses up food store;

respiration may be more than photosynthesis;

after 35 days

seeds/fruits/leaves, fall off;

3

(iii) to make results more reliable;

some seeds may, not continue to grow/die; some seedlings grow at different rates so gives a better pattern;

max 2

(b) (i) amount of water may vary in seedling; evaporation of water from soil may vary, affecting results;

max 1

(ii) not enough seed/only 60 seeds, as only single plant; destroys plants with each reading/AW; AVP;

max 2

(c) 'in terms of short day plant'

plants flower when long, dark/night;

short day plant;

phytochrome;

long night converts P<sub>FR</sub>/P730 to P<sub>R</sub>/P660;

AVP; e.g. low P<sub>FR</sub> allows flowering, P<sub>R</sub>/P660 and P<sub>FR</sub>/P730, during day is P<sub>FR</sub>, P<sub>FR</sub> is active form, Inhibits flowering in SDP max 4

Γ	Page 7	Mark Scheme	Sylla
	_	GCE A/AS LEVEL – May/June 2006	9700
2	regi incr GH stim inhi	erior (lobe of) pituitary, produces growth hormone/GH; ulates growth of all parts of the body; eases rate of, cell growth/cell division/protein synthesis; release controlled by hypothalamus/AW; nulated by growth hormone releasing factor/GHRF; bited by growth hormone release-inhibiting hormone/GHRIH; P; e.g. GH favours use of fat, so body less fat and more muscle:	Cambridge com

2 (a) Anterior (lobe of) pituitary, produces growth hormone/GH; regulates growth of all parts of the body; increases rate of, cell growth/cell division/protein synthesis; GH release controlled by hypothalamus/AW; stimulated by growth hormone releasing factor/GHRF; inhibited by growth hormone release-inhibiting hormone/GHRIH; AVP; e.g. GH favours use of fat, so body less fat and more muscle; no feedback inhibition;

max 3

**(b) (i)** 0 to 2 years, Y more rapid increase in growth/height than Z; Y reaches puberty/growth spurt at 10,Z at 12 yrs;

Y adult height taller/figs, than Z; Y reaches final height sooner/ora; ref. comparative figs.;

max 3

(ii) deficient diet during pregnancy, lower birth weight, reduced growth; breast milk deficiencies/AW; lack of protein/adequate diet/named nutrient; different genetic makeup/genotype; different levels of growth hormone; hormonal differences/named hormone level; AVP;

max3

Page 8	Mark Scheme	Sylla
	GCE A/AS LEVEL – May/June 2006	9700
Q ere R pr S ep	ethra, ectile tissue, rostate gland, pididymis; mark for 2 correct, rounded up.	Cambridge.com

**(b) (i)** 92-52, 40 40/92 X 100; 43.(125)%;

2

(ii) increase in oestrogen/female hormone in drinking water, reference sperm production/affects male hormones/testosterone;

(c)

spermatogenesis	oogenesis	
Continuous after puberty,	In cycles after puberty;	
Millions produced,	One/few per cycle;	
Occurs 12-65+,	9 -menopause/40;	
4 sperm , 1 ovum per meiosis;		
No,	Polar bodies;	
All mitotic products used	many mitotic products degenerate/less mitotic replication;	
Complete meiosis on release,	s on release, Completes meiosis after ovulation/AW;	
Primary spermatocyte smaller,	Than primary oocyte/primary oocyte greater growth phase;	
Products need to differentiate	no differentiation of products;	
Requires testosterone,	Requires oestrogen;	

max 4

Page 9	Mark Scheme	Sylia	er
	GCE A/AS LEVEL – May/June 2006	9700	

- 4 (a) (i) asexual;
  - (ii) plantlets could be removed, grown to give large number; parent plant can be used over and over, so cheaper; plantlets could be transported easily as smaller than parent; genetically identical/clone; AVP;

max 2

(b) (i) changes, gene/DNA, base sequence/described e.g. deletion/addition/substitution; codes for protein/enzyme with different amino acid/acids; codes for protein with amino acids missing; enzyme/protein has different, tertiary/active site structure/3D shape; AVP; detail of protein synthesis e.g. change to mRNA, changes t-RNA;

max 3

(ii) idea of less chlorophyll/less photosynthesis, less growth; ora

1

Page 10	Mark Scheme	Sylla	
	GCE A/AS LEVEL – May/June 2006	9700	

## **OPTION 4 - Applications Of Genetics**

1 (a) autosomal / chromosome 7;

recessive (allele);

homozygote sufferer;

heterozygote carrier;

correct statement re inheritance; [e.g. 1in 4 from two carrier parents]

max 4

**(b) (i)** move towards anode;

because negatively charged;

rate of movement inversely proportional to, mass/length;

smaller fragments move further / ora;

max 3

(ii) one of two lower bands;

1

(iii) C is heterozygote;

different allele on each homologue;

one normal (100 bp) fragment and mutant (97 bp) fragment;

max 2

(c) (i)  $\Delta$ F508 CFTR not inserted in membrane so no conductance possible;

R117H CFTR different, shape / 3' structure so poorer conductance;

does not fit ions correctly;

effect on Cl greater than HCO<sub>3</sub>;

does not bind ATP correctly;

max 3

(ii)  $(33-5) \times 100$ ;

84.8 (%);

2

Page 11	Mark Scheme	Sylla
	GCE A/AS LEVEL – May/June 2006	9700

2 (a) (i) straight line showing unchanged activity;

(ii) inhibits VKOR; non-competitive; binds to VKOR and alters shape of active site; too little vitamin K produced; vitamin K involved in clotting;

max 2

(b) (i) gene mutation; substitution of base (pair) in DNA; change of triplet code;

max 3

(ii) different, primary structure; different shape / 3' structure; ref. active site; no longer binds warfarin; enzyme not inhibited; no problem with vitamin K metabolism;

so encodes different amino acid;

max 4

	Page	12	Mark Scheme	Sylla er	
			GCE A/AS LEVEL – May/June 2006	9700 Page	
3	(a)	(i)	no chlorophyll; no photosynthesis; no, primary pigment / reaction centre / photosystem;	9700 TODA ET 9700 PARTIES TO ANNUAL PROPERTY OF THE PARTIES TO ANNUAL PROP	10
		(ii)	AaBB yellow AaBb yellow aaBB green aaBb green half marks rounded up	·	2
	(b)	(i)	clear diagram showing: cross over in between two loci of non-sister chromatids giving Al other chromatids unchanged;		2
		(ii)	large number of, parental types / AB and ab; small number of, recombinant types / Ab and aB; more recombinants further loci are apart / ora;	max 2	2
				Total: 8	3
4	(a)		parents isolated; seed parent, emasculated / AW; flower bagged; before and after pollination; pollination by hand;	max :	3
	(b)		annual life cycle / AW; idea several generations needed for selection; for large, sweet and black; backcross to commercial variety; increase contribution of commercial variety / ora; ref. alleles of background genes;	max 4	4