


CANDIDATE

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level



9700/32

May/June 2010

CANDIDATE

NUMBER

2 hours

Candidates answer on the Question Paper.

Additional Materials:

As listed in the Confidential Instructions

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do **not** use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

At the end of the examination, fasten all your work securely together.


The number of marks is given in brackets [] at the end of each question or part question.

You are advised to spend one hour on each question.

For Examiner's Use	
1	
2	
Total	

This document consists of 13 printed pages and 3 blank pages.





www.PapaCambridge.com You are reminded that you have only one hour for each question in the practical examile You should read carefully through the whole of each question and then plan your us the time to make sure that you finish all of the work that you would like to do.

You will gain marks for recording your results according to the instructions.

Yeast cells contain enzymes which catalyse the breakdown of glucose to produce carbon dioxide and water.

The carbon dioxide reacts with water and forms a weak acid.

Bromothymol blue is a pH indicator and changes colour as shown in Table 1.1.

Table 1.1

рН	colour of bromothymol blue
8	blue
7	green
6	yellow

You are required

- to immobilise the yeast cells in sodium alginate beads
- to follow a student's procedure to investigate the independent variable, changing the surface area of the beads.

You are provided with

- 15 cm³ of yeast suspension, labelled Y
- 15 cm³ of 2.0 % sodium alginate solution, labelled S
- 50 cm³ of 1.5% calcium chloride solution, labelled C
- 40 cm³ of 2.0 % glucose solution, labelled G
- 50 cm³ of bromothymol blue, labelled B
- 20 cm³ of sodium hydroxide solution, labelled A
- 1. Put 20 cm³ of **C** into a large test-tube.
- 2. Put 5 cm³ of **S** into a small beaker or container.
- Collect 5 cm³ of Y from below the froth and put it into the same container as S. Mix 3. well.
- 4. Use a 5 cm³ syringe to collect 2 cm³ of the mixture **S** and **Y**.

5. Suspend the 5 cm³ syringe over the large test-tube containing **C** as shown in Fig.




Fig. 1.1

- Gently press down on the plunger of the syringe with your thumb to release a drop into solution C. The drop should form a bead.
- Repeat step 6 to make the number of beads that you think you will need. 7.
- 8. Tip the contents of the large test-tube into a Petri dish or shallow container.

e blunt forceps Company

You will need to calculate the mean surface area of the beads. Use blunt forceps up the beads.

To do this

- decide on the number of beads you will measure
- use the 2mm x 2mm grid to measure each bead
- calculate the surface area of each bead using the formula surface area = $4\pi r^2$ where $\pi = 3.14$, r = radius of a bead
- calculate the mean surface area of the beads.
- (a) (i) Prepare the space below to show your measurements and calculations.

Show all the steps in your calculation of the mean.

mean surface area of the beadsmm² [5]

www.PapaCambridge.com A student suggested that it was possible to investigate the independent variable, area, by changing the number of beads. The maximum number of beads used was Decide the other numbers of beads to use and state the different number of beads you was use.

Carry out the student's procedure.

- 9. Label as many small test-tubes as you will need with the number of beads for each test-tube.
- 10. Put 10 cm³ of solution **G** into each test-tube.
- 11. Put 1 cm³ of **B** into each test-tube. Put the bung in each test-tube in turn and mix.
- 12. If the contents of the test-tube are not blue, add one drop at a time of A to the contents of each test-tube to turn them all the same blue colour.
- 13. Put the required number of beads into each test-tube.
- 14. Put the bung in each test-tube in turn and mix contents. Mix every 2 minutes for 6 minutes.
- 15. Record your observations after each 2 minutes, up to 6 minutes.
 - (ii) Prepare the space below to record your observations.

(iv)

	[1]
Suggest how you would make three improvements to the student's procedure.	
	[3]

A student set up the apparatus as shown in Fig. 1.2 as another way to measure the carbon dioxide produced by immobilised yeast cells over a period of 75 minutes. The student measured the distance the liquid moved in the capillary tubing.




Fig. 1.2

Table 1.2

time /min	total distance moved by the liquid in the capillary tube /mm
0	0
15	3
30	10
45	18
60	19
75	19

Describe and explain the results shown in Table 1.2.
[3]

[Total: 19]

www.PapaCambridge.com

- g. 2.1.

 RahaCambridge.com
- 2 M1 is a slide of a stained transverse section through a plant organ.
 - (a) (i) Draw a large plan diagram of a half of the specimen as shown in Fig. 2.1.




Fig. 2.1

Label the xylem and the cortex.

www.PapaCambridge.Com

- (ii) Make a high-power drawing of
 - a group of three complete touching xylem vessels and
 - a group of three complete touching cortex cells.

Label a cell wall and a lumen.

xylem vessels

cortex cells




Fig. 2.2

The actual length of the line \boldsymbol{X} is 1900 μm .

(b) (i) Calculate the magnification of the specimen shown in Fig. 2.2. Show all the steps in your calculation.

www.PanaCambridge.com (ii) Prepare the space below so that it is suitable for you to record the obs differences between the specimens on slide M1 and in Fig. 2.2.

Record your observations in the space you have prepared.

The apparatus was set up by the student as shown in Fig. 2.3.





Fig. 2.3

The volume of water added to top up the water level was recorded each hour for a 5 hours.

Table 2.1

time /hours	volume of water absorbed /cm³hr ⁻¹
1	0.9
2	1.5
3	2.1
4	2.3
5	1.6

(c) Plot a graph of the data shown in Table 2.1.



[4]

[Total: 21]

BLANK PAGE

www.PapaCambridge.com

BLANK PAGE

www.PapaCambridge.com

16

BLANK PAGE

www.PapaCambridge.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.