WWW. Pals

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the October/November 2011 question paper for the guidance of teachers

9700 BIOLOGY

9700/41

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page	2 Mark Scheme: Teachers' version	Syllabus	
	GCE AS/A LEVEL – October/November 2011	9700	
Mark scher	me abbreviations:	Calif	
;	separates marking points	ST.	
1	alternative answers for the same point		
R	reject	, co	
Α	accept (for answers correctly cued by the question, or by extra guidance)		
AW	alternative wording (where responses vary more than usua	ni)	
<u>underline</u>	actual word given must be used by candidate (grammatical variants excepted)		

Mark scheme abbreviations:

max indicates the maximum number of marks that can be given

or reverse argument ora

marking point (with relevant number) mp

error carried forward ecf

ignore

AVP Alternative valid point (examples given as guidance)

				The state of the s			
	Pa	ige 3	}	Mark Scheme: Teachers' version Syllabus GCE AS/A LEVEL – October/November 2011 9700	er er		
1	(a)	allo	Mark Scheme: Teachers' version GCE AS/A LEVEL – October/November 2011 gllopatric; packs / populations, isolated from each other; inbreeding / no interbreeding;				
	(b)	3.	inbr little	ks / populations, isolated from each other; reeding / no interbreeding; remutation; recording; recording / no interbreeding; recording / no interbreeding / no interbreed	[2 max]		
	(c)		<i>idea</i> hun	culture / buildings / AW; a of wolves dying; ting / trapping / AW; ridisation / infertility / change in (wolf) gene pool / loss of wolf alleles / AW	; [4]		
	(d)		-	one mark for number not rounded up or incorrect answer but correct idea regarding			
					[Total: 9]		
2	(a)	1. 2.		three colours (for positive reactions) / only a small range; measurement of actual concentration / no numerical value measured;	[2]		
	(b)	(i)	per	oxidase;	[1]		
		(ii)	1. 2. 3. 4.	(catalyses breakdown of hydrogen peroxide) to produce oxygen; chromogen, oxidised by / reacts with, (oxygen); produces range of colours; more, peroxide / oxygen produced, = greater change / darker colour;	[2 max]		
		(iii)	1. 2. 3.	to keep out, proteins / enzymes / polymer / named large molecule; R large molecules unqualified to prevent interference (to reactions); to prevent loss of, enzyme / chromogen; so still allowing reaction to occur;	[2 may]		
			4.	so still allowing reaction to occur;	[2 max]		

(c) (i) 1.

in A

in B

B has diabetes and **A** does not;

3. (insulin affects), liver / muscle, cells;

5. increase in use of glucose in respiration;6. (more) glucose converted to glycogen;

7. because cells unresponsive to insulin;

A's, values / peak, lower because he secretes insulin

B's, values / peak, higher because, no / little, insulin;

accept quoted values for lower and higher in mark point 2

4. increase in glucose uptake / increase in permeability of membranes (to glucose);

[4 max]

Page 4	Mark Scheme: Teachers' version	Syllabus	er
	GCE AS/A LEVEL – October/November 2011	9700	20

- (ii) (the concentration of blood glucose), above which some glucose appears in AW;
- (iii) 1. (at first), glucose reabsorbed by proximal convoluted tubule;
 - 2. ref. co transported with Na⁺ / facilitated diffusion / protein carrier;
 - 3. above 180mg (100cm⁻³ glucose in blood) no further reabsorption;
 - 4. because carriers (in PCT) saturated / AW;

[3 max]

[Total: 15]

3 (a)

	male		female	
1	produces sperm	or	produces, oocyte	;
2	division of cytoplasm is equal	or	division of cytoplasm is unequal	;
3	four gametes produced	or	one gamete produced	;
4	no polar bodies	or	polar bodies	;
5	ref. maturation	or	no equivalent maturation stage	;
6	ref. meiosis completed	or	ref. incomplete meiosis	;

[3 max]

- (b) 1. a ductless gland;
 - 2. hormones in the blood;
 - 3. ref. target, organ / tissues;

[2 max]

- (c) 1. (both), reduce / stop, secretion (of FSH and LH);
 - 2. (both) involve negative feedback;
 - 3. to, anterior pituitary / hypothalamus;
 - 4. both are, contraceptives / description;

[3 max]

[Total: 8]

- 4 (a) 1. low oxygen (in water) results in anaerobic respiration;
 - 2. (anaerobic respiration) produces alcohol;
 - 3. rice tolerant to alcohol;
 - 4. (because rice has) high levels of, alcohol dehydrogenase / enzyme that breaks down alcohol;
 - 5. presence of, aerenchyma / described;
 - 6. allows, oxygen / air, to reach roots (from aerial tissues);

[3 max]

- (b) (i) 1. (immersion in water) stimulates production of ethene;
 - 2. (concentration of) ethene produced increased with time (after submergence);
 - 3. very little difference in ethene production between T65 and C9285;
 - 4. use of figures; 2 values of ethene plus 2 values of time for either T65 or C9285

[2 max]

Page 5		<u> </u>	Mark Scheme: Teachers' version Syllabus	er	
			GCE AS/A LEVEL – October/November 2011 9700	No.	
		(ii)	2.	in T65 ethene does not affect internode elongation but in C9285 eth internode elongation; in C9285, greater concentrations of ethene cause greater elongation; use of comparative figures to support mark point 1 or mark point 2; least once	Ori
	(c)	1. 2. 3. 4.	incr GA	genes present in C9285 / SK genes not present in T65; eased production of GA in C9285 / little or no increased production of GA stimulates, stem elongation / AW; P; e.g. T65 has no receptors for ethene	in T65 ; [3 max]
	(d)	(i)		? more important ; ora nivara has mutated <i>SK2</i> and does not have deepwater response	
				nlumaepatula has SK2 but not SK1 and does have deepwater response	e; [2]
		(ii)	1. 2. 3.	(addition / insertion), of a, base / nucleotide, to DNA / to a gene; changes a, sequence of three bases / triplet / codon; ignore ref. to fr (triplet) no longer codes for an amino acid;	rame shift [2 max]
		(iii)	2. 3.	breed deepwater variety with (high-yielding) non-deepwater variety; identify / select, offspring with both deepwater response and high yield breed selected offspring (with both deepwater response and high yield continue for many generations;	
					[Total: 17]
5	(a)	 3. 4. 6. 	delive only ease serie	sed by a single gene; sed by a recessive allele; very of, correct / dominant / normal, allele (could correct the condition) red to get allele into a few cells; e of access to affected area; ous so worth the risk; P; e.g. only targets eye / no surgery needed	; [3 max]
	(b)	1. 2. 3.	corr	s no longer able to cause infections ; ect / dominant / normal, allele (of <i>RPE65</i>) added ; moter added ;	[2 max]
	(c)	2. 3.	rare exp	to safety / not known if the technique might have side effects; condition; ense; ?; e.g. trial to see if delivery method works	[2 max]
					[Total: 7]
6	(a)	(i)	pho	sphorylation ;	[1]

(ii) <u>lysis</u>;

[1]

Page 6	Mark Scheme: Teachers' version	Syllabus	er
	GCE AS/A LEVEL – October/November 2011	9700	100

(iii) dehydrogenation / oxidation; ignore reduction of NAD

(b) provides activation energy / AW; for it to split / AW;

- (c) 1. decarboxylated / carbon dioxide given off;
 - 2. ethanal produced;
 - 3. ethanal reduced;
 - 4. by reduced NAD;
 - 5. to ethanol;
 - 6. dehydrogenase;

[4 max]

[Total: 9]

7

step	reason for step	
obtain copies of gene with sticky ends	the gene codes for the synthesis of insulin	
plasmid (used);	acts as a vector for the transfer of the gene into the host	
use restriction endonuclease enzyme	to produce 'sticky ends' or cut at specific, site / sequence;	
mix vector and gene	gene inserts into, vector / plasmid or forms recombinant DNA / AW;	
	A detail of complementary base pairing	
(use DNA) ligase ;	to seal the sugar-phosphate backbone	
insert, plasmid / vector, into host / E. coli / bacteria;	to obtain transformed host <i>E. coli</i> cells	
screen for, and obtain, successfully transformed cells	so only recombinant host cells cultured / AW;	
ref. batch / continuous, culture or fermenter or bacterial cloning / population growth;	to obtain large amounts of insulin for extraction and purification	

[7]

[Total: 7]

Page 7	Mark Scheme: Teachers' version	Syllabus	er
-	GCE AS/A LEVEL – October/November 2011	9700	100

8 (a) any two from

(cigarette) smoke ;
named air pollutant ;
animal fur / skin flakes / AW ;
perfumes / aerosol / solvents ;
dust / mites ;
pollen / spores ;

[2 max]

- (b) (i) high tensile strength / withstands pulling forces / fibrous / insoluble / forms fibrils / flexible; [1]
 - (ii) 1. inbreeding;
 - 2. little genetic variation / small gene pool / small population;
 - 3. many carried faulty allele / AW;
 - 4. faulty / mutant, allele, could be dominant or recessive;
 - 5. little phenotypic variation;

[3 max]

[Total: 6]

9 (a) dominant

(allele) that always expresses itself (in the phenotype) when present

or

(allele) which influences the phenotype even in the presence of an alternative allele;

gene

length of DNA / sequence of nucleotides, coding for a (specific) polypeptide; A protein [2]

(b)

parental phenotypes	man without TSC	woman with TSC	
parental genotypes	tt	Tt	
gametes	all t	T or t	;
offspring genotypes	Tt	tt	
offspring phenotypes	TSC	normal	;
probability of child having TSC	50% / 0.50 / 1in 2 ;		

[3]

- (c) 1. spontaneous / random / chance;
 - 2. mutation of, gene / allele;
 - 3. AVP; e.g. named mutagen / detail of mutation

[2 max]

[Total: 7]

Page 8	Mark Scheme: Teachers' version	Syllabus	er	
	GCE AS/A LEVEL – October/November 2011	9700	100	
	•	•	AO 4	_

- 10 (a) 1. ground substance / stroma;
 - 2. for, light independent stage / Calvin cycle;
 - 3. contains enzymes / named enzyme e.g. rubisco;
 - 4. also, sugars / lipids / starch / ribosomes / DNA;
 - 5. internal membrane system;
 - 6. for, light dependent stage;
 - 7. fluid-filled sacs / thylakoids;
 - 8. grana are stacks of thylakoids;
 - 9. (grana) hold (photosynthetic) pigments;
 - 10. (grana) have large surface area for (maximum) light absorption;
 - 11. (pigments are arranged in), light harvesting clusters / photosystems;
 - 12. primary pigment / reaction centre / chlorophyll a, surrounded by accessory pigments;
 - 13. (accessory pigments) pass energy to, primary pigment / reaction centre / chlorophyll a;
 - 14. different photosystems absorb light at different wavelengths;
 - 15. membranes hold, ATP synthase / electron carriers;
 - 16. for, photophosphorylation / chemiosmosis;

[9 max]

- (b) 17. grind leaf with solvent;
 - 18. example of solvent; e.g. propanone
 - 19. leaf extract contains mixture of pigments;
 - 20. ref. concentrate extract;
 - 21. further detail; e.g. pencil line drawn / extract placed on chromatography paper / repetitive spotting / drying between spots
 - 22. paper placed (vertically) in jar of (different) solvent;
 - 23. solvent rises up paper;
 - 24. each pigment travels at different speed;
 - 25. pigments separated as they ascend;
 - 26. distance moved by each pigment is unique;
 - 27. Rf value;
 - 28. two dimensional chromatography;
 - 29. better separation of pigments;

[6 max]

[Total: 15]

- **11** (a) 1. axon phospholipid bilayer impermeable to K⁺ / Na⁺;
 - 2. sodium potassium pump;
 - 3. detail of sodium-potassium pump; e.g. transmembrane / globular / ATP binding site
 - 4. active process / ATP used / energy needed;
 - 3 Na⁺ (pumped) out / 2 K⁺ (pumped) in ;
 - 6. K⁺ diffuse out / Na⁺ diffuse in ;
 - 7. through, protein channels transport proteins;
 - 8. more K⁺ channels open than Na⁺ channels;
 - 9. therefore, membrane more permeable to K⁺ or more K⁺ leave than Na⁺ enter (axon);
 - 10. inside relatively more negative than outside;
 - 11. -65mV; **A** -70mV
 - 12. idea of leaking K⁺ responsible for resting potential / AW;
 - 13. electrochemical gradient;
 - 14. voltage-gated channels closed;

[9 max]

Page 9	Mark Scheme: Teachers' version	Syllabus	· Q er
•	GCE AS/A LEVEL – October/November 2011	9700	100

(b) general

- 15. respond to stimuli / AW;
- 16. (some) receptors are the ends of sensory neurones;
- 17. (some) receptors are cells;
- 18. they are energy transducers;
- 19. stimulus causes sodium ion channels to open;
- 20. sodium ions enter cell;
- 21. depolarisation;
- 22. receptor / generator, potential;
- 23. if (receptor potential) greater than threshold then action potential generated / all or nothing principle described;
- 24. increased stimulus strength leads to increased frequency of action potentials;

examples – allow any two below

receptor	form of energy detected
rods / cones	light;
taste buds / olfactory cells	chemical;
Pacinian \ Meissner's, corpuscle	pressure / touch ;
Ruffinis endings	heat;
proprioreceptors	mechanical displacement;
hair cells in semicircular canals	movement;
hairs cells in cochlea	sound;

[max 2] [6 max]

[Total: 15]