

CANDIDATE
NAME

--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

BIOLOGY

9700/21

Paper 2 AS Level Structured Questions

May/June 2019

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **14** printed pages and **2** blank pages.

Answer **all** questions.

- 1 (a) Antibody molecules are proteins that show primary structure, secondary structure, tertiary structure and quaternary structure.

Fig. 1.1 shows a ribbon diagram of an antibody molecule.




Fig. 1.1

Describe how Fig. 1.1 shows the secondary structure **and** tertiary structure of the antibody molecule.

secondary structure ... α -helix and β -pleated structures that loop

tertiary structure ... secondary structures folded onto itself to be globular

[3]

(b) Fig. 1.2 is a transmission electron micrograph of a hybridoma cell.




Fig. 1.2

- (i) The hybridoma cell in Fig. 1.2 synthesises and secretes molecules of a monoclonal antibody.

State the roles of the structures labelled X and Y in the production of antibody molecules in the hybridoma cell.

X site of synthesis of light and heavy polypeptides

Y production of ATP

[2]

- (ii) The hybridoma method for the production of monoclonal antibodies involves a number of stages. One of these stages is the formation of hybridoma cells.

Outline the stage in which hybridoma cells are formed.

Cell membrane fusion by using a fusogen like polyethylene glycol between plasma cells and myeloma cells.


[3]

- (iii) Outline the use of monoclonal antibodies in the treatment of disease.

mAbs do not damage other non-target cells as they bind to specific antigens. We can attach drugs / radioactive isotopes / enzymes to mAbs so drugs can be activated at site of action. mAbs can be used for passive immunity and can stimulate immunity system to kill cancer cells.

[3]

[Total: 11]



- 2 Linoleic acid is an unsaturated fatty acid that is found in some triglycerides and some phospholipids.

Phospholipids are components of cell membranes.

Fig. 2.1 shows a molecule of linoleic acid.




Fig. 2.1

- (a) The composition of cell membranes of plants changes in response to changes in temperature.

At the start of the cold season there is an increase in the proportion of phospholipids with unsaturated fatty acids in the chickpea, *Cicer arietinum*. Chickpea plants that do not make this change do not survive.

Suggest how the increase in the proportion of phospholipids with unsaturated fatty acids helps plants, such as chickpea, survive decreases in temperature.

- Prevents close packing of phospholipids
- increases fluidity
- prevent damage to membranes by preventing freezing
- maintains movement of substance across the membrane

[2]

- (b) (i) State why triglycerides and phospholipids **cannot** be described as polymers.

As they do not have repeating sub-units/
monomers

[1]

- (ii) State two **differences** in the **structure** of triglycerides and phospholipids.

- 1 Triglyceride has 3 fatty acid chains, while
pl has 2 fatty acid chains
- 2 PL has a phosphate group but triglyceride
does not

[2]

- (c) Platelets metabolise linoleic acid to produce a molecule known as thromboxane.

Thromboxane is released by platelets when blood loss occurs. Thromboxane acts on smooth muscle cells in the walls of arteries. This causes arteries to constrict, which reduces blood flow.

Explain why the constriction of arteries following blood loss is an example of cell signalling.

- Thromboxane is a cell signalling molecule
- Target cells are smooth muscle cells
- Thromboxane travels in the blood and binds to
specific receptors on cell - as it is complementary.
- results in activation of enzyme cascade
- response is arteries constricting as smooth
muscle contracts

[4]

[Total: 9]

- 3 Neutrase® is an enzyme that is used to hydrolyse proteins in solution. When the enzyme is mixed with a 2% protein solution the reaction mixture changes from white to colourless.

A student carried out an experiment to find the effect of copper sulfate and potassium sulfate on the activity of Neutrase®.

The student made four reaction mixtures in test-tubes **A** to **D**. Test-tubes **A** to **C** contained equal volumes of protein solution and 0.1 cm^3 of solutions of copper sulfate or potassium sulfate. Test-tube **D** contained the same volume of protein solution and 0.1 cm^3 of water.

0.5 cm^3 of a 1% Neutrase® solution was added to test-tube **A** and immediately placed into a colorimeter. The colorimeter was used to measure the intensity of light that is absorbed by the solution (absorbance) over 100 seconds. The procedure was repeated with the other reaction mixtures, **B**, **C** and **D**.

The results are shown in Fig. 3.1.




Fig. 3.1

- (a) (i) Suggest **and** explain why measuring the absorbance of the reaction mixture over 100 s is a suitable method for determining the activity of Neutrase®.

- To allow time for reaction to complete
- During the reaction, less light is absorbed.
- Neutrase® hydrolyses proteins into smaller molecules.

[2]

(ii) With reference to Fig. 3.1:

- describe the effects of copper sulfate solution and potassium sulfate solution on the activity of Neutrase®
- suggest explanations for the effects that you have described.

- Copper sulfate decreases the activity of Neutrase
- $0.01 \text{ mol dm}^{-3} \text{ CuSO}_4$ has less of an effect on Neutrase than 0.05 mol dm^{-3} of CuSO_4
- K_2SO_4 has little to no effect on activity on Neutrase
- CuSO_4 is an inhibitor of Neutrase
- CuSO_4 binds to Neutrase
- K_2SO_4 is not an inhibitor
- fewer E+S complexes form as of CuSO_4

[5]

(b) Neutrase® can be immobilised in alginate. Immobilised Neutrase® is used in the food industry to produce foods with high nutritional content.

Explain the advantages of using immobilised enzymes, such as Neutrase®, compared with using the same enzymes free in solution.

- Immobilised enzymes are more thermostable/pH resistant
- Longer shelf life
- Can be reused

[2]

[Total: 9]

- 4 Meristematic tissue is found in the growing regions of plants, such as shoot tips. Meristematic cells have a similar role to stem cells in animals.

Fig. 4.1 shows some of the stages in the formation of a mature phloem sieve tube element and companion cells from a meristematic cell.




Fig. 4.1

- (a) Cells **E** and **F** in Fig. 4.1 are daughter cells produced when the meristematic cell divides in the shoot tip.

Explain why it is important that one of the daughter cells (cell **E**) is a meristematic cell.

For continued growth of the shoot tip/tissue and to provide cells that can divide,

[1]

- (b) Complete Table 4.1 to describe the changes that are shown in Fig. 4.1 between stages:

- **F** and **G**
- **G** and **H**
- **H** and **J**.

Table 4.1

stages	description
F and G	cell grows and a vacuole is formed
G and H	cell divides and one of the cell has vacuole
H and J	developing sieve tube grows and loses its nucleus

[3]

- (c) Explain how the structure of a mature sieve tube element is related to its function.

- Elongated cells to form long tubes
- little cytoplasm and fewer organelles to reduce resistance to flow
- Sieve plates have pores so little barrier to flow from cell to cell
- Sieve plates support sieve tube elements.
- Plasmodesmata present between sieve tube and companion cell for ease of loading and unloading.

[4]

(d) Describe the functions of companion cells in transport in the phloem.

It allows sucrose from source to sink and moves sucrose in and out of sieve tubes through the plasmodesmata. It pumps protons/ H^+ out of the cell and provides ATP for sieve tubes. [2]

[Total: 10]

5 (a) The mammalian circulatory system is described as a closed double circulation.

Explain why it is called a *closed* and *double* circulation.

closed as blood flows through the blood vessels

double blood flows through the heart twice in one complete circulation [2]

(b) Fig. 5.1 shows a drawing of an external view of a mammalian heart.

Two cross-sections were made of the heart:

- section 1 was made across the line A–B.
- section 2 was made across the line C–D.

Drawings of the two sections were viewed from above as shown by the arrow on Fig. 5.1. Fig. 5.2 is a drawing of section A–B. Fig. 5.3 is a drawing of section C–D.




Fig. 5.1




Fig. 5.2



Fig. 5.3

- (i) Name structures 1, 2 and 3, as shown in Fig. 5.2.

1 pulmonary vein
 2 semi-lunar/pulmonary valve
 3 right atrium

[3]

- (ii) Explain why the wall of chamber Y is thicker than the wall of chamber X, as shown in Fig. 5.3.

Distance travelled by blood pumped from chamber Y is greater than distance travelled by blood from chamber X so higher pressure is required for blood to flow the longer distance which requires more muscular force.

[3]

- (c) Explain how the contractions of the chambers of the heart are coordinated during one cardiac cycle.

Impulse passes from SAN to atria muscles resulting in both atria contracting together. Atria contract before the ventricles as fibrous/non-conducting tissue prevents impulse from travelling to ventricles. Impulse is delayed at AVN and AVN passes impulse to Purkyně fibres that conduct impulses to muscles of the wall of ventricles. Then ventricles contract together from the bottom upwards.

[4]

[Total: 12]

6 (a) The DNA in the nucleus is known as nuclear DNA.

- (i) In the cells of the grasshopper, *Chorthippus brunneus*, 20% of the nucleotides in nuclear DNA contain thymine.

Calculate the percentage of nucleotides in the nuclear DNA of *C. brunneus* that contain guanine **and** explain your answer in terms of the structure of DNA.

percentage 30%.

explanation Bases are complimentary so hydrogen bonds between A and T or C and G.

$$A + T = 2(20\%) = 40\%$$

$$C + G = 100\% - 40\% = 60\%$$

$$G = \frac{60\%}{2} = 30\%$$

[3]

- (ii) State another location, **other than** the nucleus, where DNA occurs in cells of *C. brunneus*.

mitochondrion

[1]

(b) Fig. 6.1 is a diagram of a molecule of tRNA.

The region labelled **R** shows detail of part of the tRNA molecule.




Fig. 6.1

(i) Complete Fig. 6.1 by writing the sequence of bases in the region labelled **R**. [1]

(ii) State the name of region **Q** and explain the role of region **Q** in translation.

name *anti-codon*

explanation *Q binds to codon on mRNA and this specificity ensures correct primary structure of polypeptide.*


[3]


(iii) State the function of region **P**.

It is the site of attachment of specific amino acid.

[1]

[Total: 9]





Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.