

**MARK SCHEME for the October/November 2011 question paper
for the guidance of teachers**

9701 CHEMISTRY

9701/43

Paper 4 (A2 Structured Questions), maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – October/November 2011	9701

- 1 (a) Cr^{3+} : $1s^2 2s^2 2p^6 3s^2 3p^6 3d^3$
 Mn^{2+} : $1s^2 2s^2 2p^6 3s^2 3p^6 3d^5$
- (b) (i) Any two from
- H^+ is on the oxidant/L.H. side of each of the $\frac{1}{2}$ -equations, *or* H^+ is a reactant
 - (increasing $[\text{H}^+]$) will make E^\ominus more positive
 - (increasing $[\text{H}^+]$) will drive the reaction over to the R.H./reductant side *or* forward direction
- [1] + [1]
- (ii) KMnO_4 : Purple/violet to colourless (allow **very** pale pink) [1]
 $\text{K}_2\text{Cr}_2\text{O}_7$ Orange to green [1]
[4]
- (c) (i) $\text{MnO}_2 + \text{SO}_2 \longrightarrow \text{MnSO}_4$ (*or* $\text{Mn}^{2+} + \text{SO}_4^{2-}$) [1]
manganese changes/is reduced from +4 to +2 [1]
sulfur changes/is oxidised from +4 to +6 [1]
- (ii) **No effect**, because H^+ does not appear in the overall equation *or* its effect on the $\text{MnO}_2/\text{Mn}^{2+}$ change is cancelled out by its effect on the $\text{SO}_2/\text{SO}_4^{2-}$ change [1]
[4]
- (d) (i) $\text{MnO}_2 + 4\text{H}^+ + \text{Sn}^{2+} \longrightarrow \text{Mn}^{2+} + 2\text{H}_2\text{O} + \text{Sn}^{4+}$ [1]
- (ii) $n(\text{MnO}_4^-) = 0.02 \times 18.1/1000 = 3.62 \times 10^{-4} \text{ mol}$ [1]
 $n(\text{Sn}^{2+}) = 3.62 \times 10^{-4} \times 5/2 = 9.05 \times 10^{-4} \text{ mol}$ [1]
 $n(\text{Sn}^{2+})$ that reacted with $\text{MnO}_2 = (20 - 9.05) \times 10^{-4} = 1.095 \times 10^{-3} \text{ mol}$ [1]
reaction is 1:1, so this is also $n(\text{MnO}_2)$
mass of $\text{MnO}_2 = 1.095 \times 10^{-3} \times (54.9+16+16) = 0.0952 \text{ g}$ [1]
 \Rightarrow **95% – 96%**; 2 or more s.f. [1]
[6]

[Total: 16]

Page 3	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – October/November 2011	9701

- 2 (a) (i) A molecule/ion/species with a lone pair (of electrons) or electron pair donor...
.... that bonds to a metal ion/transition element.... [2]
- (ii) ...by means of a dative/coordinate (covalent) bond [1]
- (b) (i) straight line from (0, 0.01) to point at (350, 0.0028) with all points on the line [1]
- (ii) order w.r.t. $\text{Cr}(\text{CO})_6$ is 1 **and** order w.r.t. PR_3 is zero [1]
because (a) $\text{Cr}(\text{CO})_6$ graph has a constant half-life (which is 700 s)
or construction lines on graph showing this [1]
because (b) PR_3 graph is a straight line (of constant slope) or line shows a constant rate
of reaction or no change in rate or shows a linear decrease [1]
- (iii) rate = $k[\text{Cr}(\text{CO})_6]$ [1]
 $k = (0.9 - 1.1) \times 10^{-3} \text{ (s}^{-1}\text{)}$ (one or more s.f.) [1]
either $\text{rate}_0 = 0.01/1020 = 9.8 \times 10^{-6} \text{ mol sec}^{-1}$ when $[\text{Cr}(\text{CO})_6] = 0.01 \text{ mol dm}^{-3}$
so $k = 9.8 \times 10^{-6}/0.01 = 9.8 \times 10^{-4}$
or $t_{1/2} \approx 700 \text{ sec}$
 $k = 0.693/700 = 9.9 \times 10^{-4}$
- (iv) (units of k are) sec^{-1} [1]
- (v) N.B. the chosen mechanism must be consistent with the rate equation in (iii). Thus:
either if rate = $k[\text{Cr}(\text{CO})_6]$
mechanism **B** is consistent [1]
because it's the only mechanism that does NOT involve PR_3 in its slow/rate-determining
step or only $\text{Cr}(\text{CO})_6$ is involved in slow step or $[\text{PR}_3]$ does not affect the rate [1]
- or
if rate = $k[\text{Cr}(\text{CO})_6][\text{PR}_3]$, then
mechanism **A** or **C** or **D** is consistent [1]
because both reactants are involved in slow step [1]
[9]


[Total: 11]

Page 4	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – October/November 2011	9701

- 3 (a) (i) E is $\text{CH}_3\text{CH}(\text{NH}_2)\text{CN}$
 (ii) $\text{C}_6\text{H}_5\text{CH}_2\text{CHO}$


- (b) (i) a polymer/polypeptide of amino acids, (joined by peptide bonds)
 (allow 'chain' of amino acids' but not 'sequence': the idea of 'many' has to be conveyed) [1]

(ii)



peptide bond shown in full (C=O) in an ala-ala fragment in a chain
 two repeat units [1]


Allow peptide bond shown in full (C=O) in a dipeptide ala-ala for 1 mark



[3]


- (c) (i) HCl or H_2SO_4 or NaOH or H^+ or OH^- reagents [1]
 + heat and $\text{H}_2\text{O}/\text{aq}$ (allow H_3O^+).
 If T is quoted, $80\text{ }^\circ\text{C} < T < 120\text{ }^\circ\text{C}$. NOT warm. conditions [1]

(ii)









(if a structural formula, it must have all H atoms) allow protonated or deprotonated versions [1] + [1]

[max 3]



(ii)

compound	zwitterion
	
	
	

[3]

[4]

(e) (i) A buffer is a solution whose pH stays **fairly** constant *or* which maintains **roughly** the same pH *or* which resists/minimises changes in pH [1]
when **small/moderate** amounts of acid/ H^+ or alkali/ OH^- are added [1]

(ii) $\text{NH}_2\text{CH}(\text{CH}_3)\text{CO}_2\text{H} + \text{H}(\text{Cl}) \longrightarrow {}^+\text{NH}_3\text{CH}(\text{CH}_3)\text{CO}_2\text{H} (+ \text{Cl}^-)$ [1]

(iii) blood contain HCO_3^- (*or* in an equation) [1]
which absorbs H^+ *or* equ $\text{H}^+ + \text{HCO}_3^- \longrightarrow \text{H}_2\text{CO}_3 (\text{H}_2\text{O} + \text{CO}_2)$ [1]
or absorbs OH^- *or* equ $\text{OH}^- + \text{HCO}_3^- \longrightarrow \text{CO}_3^{2-} + \text{H}_2\text{O}$ [1]

(iv) $[\text{CH}_3\text{CO}_2\text{Na}] = 0.05$ $[\text{CH}_3\text{CO}_2\text{H}] = 0.075$ [1]
 $\text{pH} = 4.76 + \log (0.05/0.075) = 4.58$ *or* 4.6 [1]

[7]

[Total: 19]

Page 6	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – October/November 2011	9701



- (b) (down the group) nitrates become **more stable** or require a higher temperature to decompose [1]
 as size/radius of (cat)ion increases or charge density of ion decreases [1]
 so polarisation/distortion of **anion/nitrate** decreases [1]
 [3]



- (ii) radius of Li ion/ Li^+ is less than that of Na ion/ Na^+ (or polarising power of M^+ is greater) [1]

- (iii) Brown/orange fumes/gas would be evolved or glowing splint relights [1]
 Since the nitrate is likely to be thermally unstable or decomposes (just like the carbonate) or the balanced equation: $2\text{LiNO}_3 \longrightarrow \text{Li}_2\text{O} + 2\text{NO}_2 + \frac{1}{2}\text{O}_2$ [1]
 [4]

[Total: 8]

Page 7	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – October/November 2011	9701

- 5 (a) Alkanes are non-polar *or* have no dipole *or* C–H bonds are strong *or* C and H have similar electronegativities

(b) (i) (free) radical substitution *or* substitution by homolytic fission [1]

(ii) initiation: $\text{Cl}_2 \longrightarrow 2\text{Cl}^\bullet$ [1]

propagation: $\text{Cl}^\bullet + \text{C}_2\text{H}_6 \longrightarrow \text{C}_2\text{H}_5^\bullet + \text{HCl}$

$\text{C}_2\text{H}_5^\bullet + \text{Cl}_2 \longrightarrow \text{C}_2\text{H}_5\text{Cl} + \text{Cl}^\bullet$ [1]

termination: $\text{C}_2\text{H}_5^\bullet + \text{Cl}^\bullet \longrightarrow \text{C}_2\text{H}_5\text{Cl}$

or $\text{Cl}^\bullet + \text{Cl}^\bullet \longrightarrow \text{Cl}_2$ etc [1]

all 3 names [1]

(iii)

structural formula of by-product	formed by
CH₂Cl–CH₂Cl (or isomer)	further substitution
CH₃CH₂CH₂CH₃	(termination of 2 ×) C₂H₅[•]
CH₃CH₂CH₂CH₂Cl (or isomer)	substitution of C₄H₁₀ by-product

[3]

accept in the “formed by” column the formulae of radicals that will produce the compound in the “by-product” column, or the reagents, e.g. $\text{C}_4\text{H}_9^\bullet + \text{Cl}_2$ *or* $\text{C}_4\text{H}_9^\bullet + \text{Cl}^\bullet$ *or* $\text{C}_4\text{H}_{10} + \text{Cl}_2$ (giving $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{Cl}$).

do not allow anything more Cl-substituted than dichlorobutane.

N.B. $\text{C}_2\text{H}_5\text{Cl}$ is the **major** product, not a **by-product**, so do not allow $\text{C}_2\text{H}_5\text{Cl}$.


(iv) J/K = **2.3 : 1** *or* 7:3 *or* 21:9 [2]

(reason: straightforward relative rate suggests 21:1, but there are 9 primary to 1 tertiary, so divide this ratio by 9. $21/9 = 2.33$)

allow [1] mark if J/K ratio is given as 21:1;

[10]

(c)



4 isomers 4 × [1]

2 chiral atoms identified correctly, even in incorrect structures

[1] + [1]

[max 5]

[Total: 16]

Page 8	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – October/November 2011	9701

- 6 (a) (i) **K**, because it is the (only) one to contain nitrogen *or* it's an amino acid *or* because it contains CO₂H *or* NH groups
- (ii) molecule: **J**, polymer: RNA (**not** DNA)
or molecule: **L**, polymer: starch, cellulose, glycogen *or* polysaccharide (**not** carbohydrate) [2]
- (b) (i) Covalent bonding [1]
(ii) Hydrogen bonding [1]
(iii) Ionic/electrovalent bonding *or* disulphide/–S–S– bonding *or* van der Waals' forces [1]
[3]
- (c) (i) Enzymes [1]
(ii) • change in pH
• increase in T (NOT decrease; T > 40 °C *or* "too high" are OK)
• addition of heavy metal ions *or* specific, e.g. Hg²⁺, Ag⁺, Pb²⁺ etc.
any two bullet points [1] + [1]
- change in pH disrupts ionic bonds
or metal ions disrupt ionic bonds
or metal ions disrupt –S–S– bonds
or heating disrupts hydrogen bonds
any one [1]
- This changes: the 3D structure *or* shape of the enzyme *or* the active site [1]
[max 4]
[Total: 9]

7 (a)

structural information	analytical technique
three-dimensional arrangement of atoms and bonds in a molecule	X-ray crystallography/diffraction
chemical environment of protons in a molecule	NMR (spectroscopy) only
identity of amino acids present in a polypeptide	Electrophoresis / chromatography / mass spectrometry

[1] + [1] + [1]
[3]

(b) (i) **paper chromatography;**

The components **partition** between the solvent/moving phase and the water/liquid stationary phase *or* separation relies on different solubilities (of components) in the moving solvent and the stationary water phase. [1]

(ii) **thin-layer chromatography.**

Separation depends on the differential **adsorption** of the components onto the solid particles/phase *or* Al_2O_3 *or* SiO_2 . [1]
[2]

(c) (i) No. of carbon atoms present = $\frac{0.2 \times 100}{5.9 \times 1.1} = 3.08$ hence 3 carbons [1]


(ii) Bromine [1]

(iii) **One** bromine is present as there is only an M+2 peak / no M+4 peak *or* the M and M+2 peaks are of similar height [1]

(iv) *NMR spectrum shows a single hydrogen split by many adjacent protons and 6 protons in an identical chemical environment. This suggests...*
two $-CH_3$ groups and a lone proton attached to the central carbon atom [1]

Empirical formula of **N** is C_3H_7Br [1]

Hence **N** is $(CH_3)_2CHBr$ *or*



[1]
[6]

[Total: 11]

Page 10	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – October/November 2011	9701

- 8 (a) (i) Soluble form would be most effective
- (ii) **Q**, since the 'mini-pills'/granules/powder have a larger surface area
or **P**, because it has no protective casing [1]
- (iii) The gel coat stops it being broken down while passing through the upper part of the digestive system/stomach
or the gel coat is stable to stomach acid. [3]
- (b) The drug is taken quickly/directly to the target
or more accurate dosing can be achieved [1]
- When the drug is taken by mouth it has to pass through the stomach/intestine wall to get into the bloodstream. or some is digested/lost to the system [1]
- [2]
- (c) (i) condensation (polymerisation) [1]
- (ii) hydrogen bonds or van der Waals' [1]
- (iii) It would change the overall shape of the (drug) molecule
The 'fit' into the active site would be less effective [1] + [1]
- (iv) Hydrolysis [1]
- [5]
- [Total: 10]