CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the October/November 2012 series

9701 CHEMISTRY

9701/36

Paper 3 (Advanced Practical Skills 2), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2012 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

www.PapaCambridge.com

| | | | V . | |
|--------|--|---------|-----|--|
| Page 2 | Mark Scheme | Syllabu | er | |
| | GCE AS/A LEVEL – October/November 2012 | 9701 | 100 | |

| Question | Section | Indicative material | Mai | Mbr. |
|----------|------------------|--|-----|-----------|
| 1 (a) | PDO Layout | I Records initial and final burette readings and titre for rough. Tabulates and records initial and final burette readings and volume of FB 2 run from burette for all accurate titrations. Not awarded if 50/50.0/50.00 shown more than once. | 1 | Inbridge. |
| | PDO Recording | II Appropriate headings and units: initial/final (burette) reading/volume, reading or volume at start/finish/beginning/end, volume used/volume added/FB 2 used/titre. Units are /cm³, (cm³) or volume in cm³. | 1 | |
| | PDO Recording | III All accurate burette readings recorded to 0.05 cm ³ (this includes 0.00). Two (minimum) accurate titrations needed. | 1 | |
| | MMO Quality | Difference between candidate's mean titre and Supervisor's is calculated. Award IV, V and VI for $\delta \leq 0.20\mathrm{cm}^3$ Award IV and V for $0.20\mathrm{cm}^3 < \delta \leq 0.40\mathrm{cm}^3$ Award IV for $0.40\mathrm{cm}^3 < \delta \leq 0.60\mathrm{cm}^3$ Spread penalty Titres (selected by Examiner) differ by $\geq 0.5\mathrm{cm}^3$ or only 1 accurate titration penalty -1. This mark is deducted from those awarded in IV to VI but no negative marks. | 3 | [6] |
| (b) | PDO Display | Calculation of mean Candidate must average two (or more) accurate titres that are within 0.20 cm³ of another. Working must be shown or ticks must be put next to the two (or more) accurate readings selected. The mean should normally be quoted to 2 decimal places rounded to the nearest 0.01. Example: 26.667 must be rounded to 26.67. Do not award this mark if: | 1 | |
| | | any selected titre is not within 0.20 cm³ of any other selected titre; the rough titre was used to calculate the mean; the candidate carried out only 1 accurate titration; burette readings were incorrectly subtracted to obtain any of the accurate titre values. Note: the candidate's mean will sometimes be marked as correct even if it is different from the mean calculated by the Examiner for the purpose of assessing accuracy. | | [1] |

| | • | May . |
|--------|--|-------------|
| Page 3 | Mark Scheme | Syllabu. er |
| | GCE AS/A LEVEL – October/November 2012 | 9701 |

| | T | | . Č | |
|---------|-----------------------|--|-----|-----------|
| (c) (i) | ACE Interpretation | 0.1 × answer to (b) calculated correctly (3 to 4 significant 1000 figures) | | Inbridge. |
| (ii) | | Answer to (c)(i) (at least 2 significant figures) | 1 | 20 |
| (iii) | | Answer to (c)(ii) × 1000 calculated correctly 25.0 (3 to 4 significant figures) | 1 | [3] |
| (d) | ACE | Any two from | | |
| | Improvements | Larger quantity of starch – no effect because starch is the indicator or wtte . | 1 | |
| | | Larger volume of KI – no effect because KI is already in excess or will have an effect because KI not (already) in excess. | 1 | |
| | | Filter before titration – effective because easier to see end-point / colour change or not effective because iodine/chemicals stay on the filter paper. | | [2] |
| | | | [To | otal: 12] |
| 2 (a) | MMO Collection | Mass of zinc used between 2.1 and 2.3 g. Subtraction must be correct from unambiguous weighings. | 1 | |
| | PDO Recording | Table completed and all temperatures recorded to 0.0 or 0.5°C. Must include at least one ending in 0.0 and one ending in 0.5. | 1 | [2] |
| (b) (i) | PDO Layout | Axes labelled temperature or T/°C or (°C) or temperature in °C (<i>y</i> -axis) and time (<i>x</i> -axis) or t/ minutes etc. Linear scales chosen so that graph occupies at least half | | |
| | | the available length for both axes. This includes the 5 °C extension. | 1 | |
| (ii) | PDO Layout | Plotting accurate (within ½ small square and in correct square). Must plot all readings taken – minimum 10. | 1 | |
| | MMO Collection | 2 straight lines drawn – one before 3 minutes and one after maximum temperature. | 1 | |
| | PDO Layout | 3 appropriate lines drawn – including extrapolations (not falling). | 1 | [4] |

| | | • | May . |
|---|--------|--|-------------|
| | Page 4 | Mark Scheme | Syllabu. er |
| ſ | | GCE AS/A LEVEL – October/November 2012 | 9701 |

| | | correct to nearest 0.5°C, decimal places not needed for 0.0. Allow ΔT at $3\frac{1}{2}$ min, even if not max, provided some indication on graph. | 1 | ambridge. |
|-------|-----------------------|--|---|-----------|
| (ii) | PDO Display | Correctly calculates 50 × 4.3 × candidate's ΔT (from (i) unless value from graph was not using maximum vertical). Must be max. | 1 | |
| (iii) | ACE Interpretation | Moles Zn = candidate's mass of zinc 65.4 (working must be shown and answer correct to significant figures shown) | 1 | |
| | | Moles $Cu^{2+} = \frac{50 \times 1.1}{1000} = 0.055$ (working must be shown and answer correct to significant figures shown) | 1 | |
| (iv) | | (c)(ii) (correctly calculated, ecf possible) 1000 × moles of Zn (from (c)(iii)) | 1 | |
| | ACE Conclusion | Sign negative and answer to 2–4 significant figures (stand alone). | 1 | [6] |
| (d) | ACE Interpretation | Error in one temperature reading = 0.5 (°C) | 1 | |
| | morpretation | Maximum % error = $\frac{1.0 \times 100}{12.0}$ = 8.3(3) % (ecf 2 × error) | 1 | |
| | | Expression or correct answer. | | [2] |

| | • | my | |
|--------|--|---------|------|
| Page 5 | Mark Scheme | Syllabu | er |
| | GCE AS/A LEVEL – October/November 2012 | 9701 | 100- |

| (a) (i | | Solution/ FB 5/liquid goes from green to blue. | 1 | 3 |
|--------|----------------------|--|-------|----------|
| | Collection | Green precipitate and insoluble in excess. | 1 | ambridge |
| | | Unqualified white precipitate, insoluble in acid. | 1 | |
| (ii) | ACE Conclusions | FB 5 is a sulfate/nickel sulfate/NiSO ₄ /SO ₄ ²⁻ . (allow conclusion even if green ppt in (i)) | 1 | [4] |
| (b) (i | | Solution/FB 6/liquid goes from yellow to orange. | 1 | |
| | Collection | (Solution) goes green/blue (allow grey-green or blue-green). | 1 | |
| | | Yellow precipitate. | 1 | |
| (ii |) ACE Conclusions | Anion in FB 6 is $CrO_4^{2^-}$ / chromate Cation in FB 7 is H ⁺ / hydrogen Cation in FB 8 is Ba ²⁺ / barium or Pb ²⁺ / lead. | 1 1 1 | |
| | Considerations | Add named (aqueous) chloride – white ppt Pb ²⁺ (not Ba ²⁺) or | | |
| (iii |) MMO Decisions | Add (aqueous) NaOH – white ppt Pb ²⁺ (not Ba ²⁺) or Add (aqueous) ammonia - white ppt Pb ²⁺ (not Ba ²⁺) or | | |
| | | Add (aqueous) KI – yellow ppt Pb ²⁺ (not Ba ²⁺) or Add (aqueous) named chromate (only allow if FB 6 not | | |
| | | identified as chromate) – yellow ppt both Ba ²⁺ and Pb ²⁺ with indication that Ba ²⁺ is paler. | 1 | |
| | | Allow ecf from candidate's FB 8. | | [7] |
| (c) | ACE Conclusions | Prediction must follow identities of FB 5 and FB 8 . (If these correct then should be white ppt.) If candidate's ions in (b)(ii) and (a)(ii) would give two different results, both must be specified. | 1 | [1] |
| (d) | ACE Conclusions | (Ethanol/it was) oxidised/ an aldehyde was formed/ a carboxylic acid was formed (not redox). | 1 | [1] |
| (e) | MMO Decisions | Uses a (named) carbonate or a (named) reactive metal to produce effervescence/ positive test for the gas. Named indicator with correct colour. NaOH with temperature increase. | | |
| | | (no ecf possible) | 1 | [1] |