

**MARK SCHEME for the October/November 2008 question paper**

**9709/07**

**9709 MATHEMATICS**

Paper 7, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2008 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme	Syllabus	Pa
	GCE A/AS LEVEL – October/November 2008	9709	07

**Mark Scheme Notes**

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol  $\surd$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking  $g$  equal to 9.8 or 9.81 instead of 10.

<b>Page 3</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Pa</b>
	<b>GCE A/AS LEVEL – October/November 2008</b>	<b>9709</b>	<b>07</b>

The following abbreviations may be used in a mark scheme or used on the scripts:

- AEF Any Equivalent Form (of answer is equally acceptable)
- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- CWO Correct Working Only - often written by a 'fortuitous' answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

**Penalties**

- MR -1 A penalty of MR -1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through  $\sqrt{\phantom{x}}$ " marks. MR is not applied when the candidate misreads his own figures - this is regarded as an error in accuracy. An MR -2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA -1 This is deducted from A or B marks in the case of premature approximation. The PA -1 penalty is usually discussed at the meeting.

<b>Page 4</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Pa</b>
	<b>GCE A/AS LEVEL – October/November 2008</b>	<b>9709</b>	<b>07</b>

<b>1 (i)</b> Not all totals have the same probability e.g. $P(7) = 6/36, P(4) = 3/36$	B1 B1	<b>[2]</b>	Or equivalent Any example to correctly justify their statement above
	-----		
<b>(ii)</b> Any valid method e.g. using pieces of paper, calculator, random number tables	B1 B1	<b>[2]</b>	Valid idea Method of choosing – full description
	-----		
<b>2</b> $H_0: \mu = 42$ $H_1: \mu \neq 42$  $\text{Test statistic } z = \frac{35 - 42}{(15.7 / \sqrt{3})}$ $= -0.772$  $  -0.772   < 1.645$  Teacher's estimate can be accepted.  <b>OR:</b> $42 \pm 1.645(15.7/\sqrt{3})$ $(27.1, 56.9)$ $27.1 < 35 < 56.9$ Teacher's estimate can be accepted.	B1 M1 A1 M1 A1ft	<b>[5]</b>	Correct $H_0$ <b>and</b> $H_1$ Standardising attempt, must have $\sqrt{3}$ used correctly Correct test statistic ( $\pm$ ) Correct comparison. $\pm 1.645$ seen or $\pm 1.64$ or $\pm 1.65$ must compare + with + or – with – (or 1.282 if one-tail test being followed) Correct conclusion. (ft) No contradictions.
	M1 A1 M1 A1ft		Correct comparison Correct conclusion
<b>3</b> $2T \sim N(72, 2 \times 1.6^2)$ $10C \sim N(73, 10 \times 0.4^2)$ $2T - 10C \sim N(-1, 2 \times 1.6^2 + 10 \times 0.4^2)$ $\sim N(-1, 6.72)$  $P((2T - 10C) > 0) = 1 - \Phi\left(\frac{0 - (-1)}{\sqrt{6.72}}\right)$ $= 1 - \Phi(0.3857)$ $= 1 - 0.650$ $= 0.350$	B1 B1 M1 M1 A1	<b>[5]</b>	Correct mean $\pm(72-73)$ $2 \times 1.6^2 + 10 \times 0.4^2$ or 6.72 seen Consideration of $2T - 10C$ and standardising, no cc, sq root Correct side ( $<0.5$ ) – consistent with their working Correct answer
	-----		
<b>4 (i)</b> $\bar{x} = 4.27$  $s^2 = \frac{1}{129} \left( 2371.3 - \frac{555.1^2}{130} \right)$ $= 0.00793$	B1 M1 A1	<b>[3]</b>	Correct mean Substituting in formula from tables (or equiv) Correct variance
	-----		
<b>(ii)</b> $CI = 4.27 \pm 2.17 \times \frac{0.08905}{\sqrt{130}}$ $= (4.25, 4.29)$	B1 M1 A1	<b>[3]</b>	Correct $z$ used (2.169–2.171) Correct form of expression $\sqrt{130}$ in denominator Correct answer (cwo)
	-----		
<b>(iii)</b> 9	B1	<b>[1]</b>	c.a.o

<b>Page 5</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Pa</b>
	<b>GCE A/AS LEVEL – October/November 2008</b>	<b>9709</b>	<b>07</b>

<b>5</b>	(i) $P(\text{Type I error}) = P(1 \text{ or more}) = 0.0202$	M1 A1 [2]	Identifying correct probability Correct answer (condone Poisson approx)
	(ii) $P(\text{Type II error}) = P(0) \text{ under } H_1 = (1 - 0.0024)^{12} = 0.972$	B1 M1 A1 [3]	Identifying correct probability Attempt to find their prob using 0.0024 Correct final answer (condone Poisson approx)
	(iii) Poisson approximation $\lambda = 0.288$ $P(2) = e^{-0.288} \left( \frac{0.288^2}{2} \right) = 0.0311$	B1 M1 A1 [3]	For $0.0024 \times 120$ in a Poisson expression Poisson expression for $P(2)$ , any mean Correct answer <b>SR</b> Use of Binomial giving final answer of 0.0310 scores B1
	<b>6</b>	(i) $\lambda = 1.15$	B1 [1]
	(ii) $P(0) \times P(>0) = e^{-1.15} \times (1 - e^{-1.15}) = 0.3166 \times 0.6833 = 0.216$	M1 A1 [2]	Multiplying two Poisson probs meant to be no goals in first half and something in second half Correct answer
	(iii) $\lambda = \frac{60}{90} \times 2.3 = 1.53(3)$ $P(\text{at least } 1) = 1 - P(0) = 1 - e^{-1.533} = 0.784$	B1 M1 A1 [3]	New mean Attempt at finding $1 - P(0)$ with new mean Correct answer (cwo)
	(iv) $\lambda = 4.1$ $P(\text{at least } 3) = 1 - P(0, 1, 2) = 1 - e^{-4.1} \left( 1 + 4.1 + \frac{4.1^2}{2} \right) = 1 - 0.224 = 0.776$	B1 M1 A1 [3]	New mean (or 6 correct combinations 0,0 1,0 2,0 etc) Using Poisson with new mean (or combinations) to find $P(\geq 3)$ condone 1 end error Correct answer

Page 6	Mark Scheme	Syllabus	Pa
	GCE A/AS LEVEL – October/November 2008	9709	07

<p>7 (i) <math>\int_1^4 kt^{0.5} dt = 1</math></p> $\left[ \frac{2kt^{1.5}}{3} \right]_1^4 = 1$ $\frac{16k}{3} - \frac{2k}{3} = 1$ $k = 3/14 \text{ AG}$	<p>M1</p> <p>A1</p> <p>A1 [3]</p>	<p>Equating to 1 and attempting to integrate (ignore limits)</p> <p>Correct integration ignore limits</p> <p>Correct answer legitimately obtained</p>
<p>(ii) mean time = <math>\int_1^4 kt^{1.5} dt</math></p> $= \left[ \frac{2}{5} kt^{2.5} \right]_1^4 = \left[ \frac{64k}{5} - \frac{2k}{5} \right]$ <p>= 2.66 hours</p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1 [4]</p>	<p>Attempting to evaluate <math>\int_1^4 kt^{1.5} dt</math> (ignore limits)</p> <p>Correct integration and correct limits</p> <p>Substituting correct limits in their integration (need not be correct)</p> <p>Correct answer</p>
<p>(iii) <math>\int_1^m kt^{0.5} dt = 0.5</math></p> $\frac{m^{1.5}}{7} - \frac{1}{7} = 0.5$ <p><math>m = 2.73</math> hours</p>	<p>M1</p> <p>M1</p> <p>A1 [3]</p>	<p>Attempt to evaluate <math>\int_1^m kt^{0.5} dt</math> (accept <math>k</math> missing)</p> <p>Attempt to solve an equation in <math>m</math>, = 0.5</p> <p>Correct answer (aef)</p>
<p>(iv) <math>\int_{2.657}^{2.726} kt^{0.5} dt = \left[ \frac{2.726^{1.5}}{7} - \frac{2.657^{1.5}}{7} \right]</math></p> <p>= 0.0243</p>	<p>M1</p> <p>A1 [2]</p>	<p>Attempt to integrate using their mean and median as limits</p> <p>Correct answer accept between 0.0241 and 0.0257</p>