



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
General Certificate of Education  
Advanced Subsidiary Level and Advanced Level

www.PapaCambridge.com

**MATHEMATICS**

**9709/61**

Paper 6 Probability & Statistics 1 (S1)

**May/June 2010**

**1 hour 15 minutes**

Additional Materials: Answer Booklet/Paper  
Graph Paper  
List of Formulae (MF9)

\* 1 1 1 5 4 8 6 5 5 \*

**READ THESE INSTRUCTIONS FIRST**

If you have been given an Answer Booklet, follow the instructions on the front cover of the Booklet.  
Write your Centre number, candidate number and name on all the work you hand in.  
Write in dark blue or black pen.  
You may use a soft pencil for any diagrams or graphs.  
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** the questions.  
Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place in the case of angles in degrees, unless a different level of accuracy is specified in the question.  
The use of an electronic calculator is expected, where appropriate.  
You are reminded of the need for clear presentation in your answers.

At the end of the examination, fasten all your work securely together.  
The number of marks is given in brackets [ ] at the end of each question or part question.  
The total number of marks for this paper is 50.  
Questions carrying smaller numbers of marks are printed earlier in the paper, and questions carrying larger numbers of marks later in the paper.

This document consists of **3** printed pages and **1** blank page.

- 1 The probability distribution of the discrete random variable  $X$  is shown in the table below.

$x$	-3	-1	0	4
$P(X = x)$	$a$	$b$	0.15	0.4

Given that  $E(X) = 0.75$ , find the values of  $a$  and  $b$ . [4]

- 2 The numbers of people travelling on a certain bus at different times of the day are as follows.

17	5	2	23	16	31	8
22	14	25	35	17	27	12
6	23	19	21	23	8	26

- (i) Draw a stem-and-leaf diagram to illustrate the information given above. [3]
- (ii) Find the median, the lower quartile, the upper quartile and the interquartile range. [3]
- (iii) State, in this case, which of the median and mode is preferable as a measure of central tendency, and why. [1]
- 3 The random variable  $X$  is the length of time in minutes that Jannon takes to mend a bicycle puncture.  $X$  has a normal distribution with mean  $\mu$  and variance  $\sigma^2$ . It is given that  $P(X > 30.0) = 0.1480$  and  $P(X > 20.9) = 0.6228$ . Find  $\mu$  and  $\sigma$ . [5]


- 4 The numbers of rides taken by two students, Fei and Graeme, at a fairground are shown in the following table.

	Roller coaster	Water slide	Revolving drum
Fei	4	2	0
Graeme	1	3	6

- (i) The mean cost of Fei's rides is \$2.50 and the standard deviation of the costs of Fei's rides is \$0. Explain how you can tell that the roller coaster and the water slide each cost \$2.50 per ride. [2]
- (ii) The mean cost of Graeme's rides is \$3.76. Find the standard deviation of the costs of Graeme's rides. [5]

- 5 In the holidays Martin spends 25% of the day playing computer games. Martin's friend phones him once a day at a randomly chosen time.
- Find the probability that, in one holiday period of 8 days, there are exactly 2 days on which Martin is playing computer games when his friend phones. [4]
  - Another holiday period lasts for 12 days. State with a reason whether it is appropriate to use a normal approximation to find the probability that there are fewer than 7 days on which Martin is playing computer games when his friend phones. [1]
  - Find the probability that there are at least 13 days of a 40-day holiday period on which Martin is playing computer games when his friend phones. [5]
- 6
- Find the number of different ways that a set of 10 different mugs can be shared between Lucy and Monica if each receives an odd number of mugs. [3]
  - Another set consists of 6 plastic mugs each of a different design and 3 china mugs each of a different design. Find in how many ways these 9 mugs can be arranged in a row if the china mugs are all separated from each other. [3]
  - Another set consists of 3 identical red mugs, 4 identical blue mugs and 7 identical yellow mugs. These 14 mugs are placed in a row. Find how many different arrangements of the colours are possible if the red mugs are kept together. [3]
- 7 In a television quiz show Peter answers questions one after another, stopping as soon as a question is answered wrongly.
- The probability that Peter gives the correct answer himself to any question is 0.7.
  - The probability that Peter gives a wrong answer himself to any question is 0.1.
  - The probability that Peter decides to ask for help for any question is 0.2.

On the first occasion that Peter decides to ask for help he asks the audience. The probability that the audience gives the correct answer to any question is 0.95. This information is shown in the tree diagram below.



- Show that the probability that the first question is answered correctly is 0.89. [1]

On the second occasion that Peter decides to ask for help he phones a friend. The probability that his friend gives the correct answer to any question is 0.65.

- Find the probability that the first two questions are both answered correctly. [6]
- Given that the first two questions were both answered correctly, find the probability that Peter asked the audience. [3]

