WWW. Palls

#### UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

**GCE Advanced Subsidiary Level and GCE Advanced Level** 

# MARK SCHEME for the May/June 2011 question paper for the guidance of teachers

# 9709 MATHEMATICS

9709/61

Paper 6, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Mbridge: com

Page 2	Mark Scheme: Teachers' version	Syllabus	
	GCE AS/A LEVEL – May/June 2011	9709	

## **Mark Scheme Notes**

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
  B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus	· A er
	GCE AS/A LEVEL – May/June 2011	9709	100-

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form	of answer is equally	/ acceptable)
-----	---------------------	----------------------	---------------

- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- CWO Correct Working Only often written by a 'fortuitous' answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

### **Penalties**

- MR −1 A penalty of MR −1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through √" marks. MR is not applied when the candidate misreads his own figures this is regarded as an error in accuracy. An MR −2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

		www
Page 4	Mark Scheme: Teachers' version	Syllabus
	GCE AS/A LEVEL – May/June 2011	9709

				3
P(2,	= 2.7  p = 0.15	B1 M1		Correct value for <i>p</i> Summing 3 binomial probs o.e  Correct unsimplified answer Correct answer
$+\frac{18}{18}$ = 0.	$C_4(0.15)^4(0.85)^{14}$	A1 A1	[4]	Correct unsimplified answer Correct answer
2 P(pe	encil case   find) =	M1		Attempt to use cond prob formula, must be quotient
$P(\mathfrak{p}$	pencilcase and find) = $0.7 \times 1$	A1		Correct num of a fraction
	$P(\text{find}) = \frac{-0.7 + 0.3 \times 0.2}{-0.7 + 0.3 \times 0.2}$	A1		Correct denominator
= 0.	921	A1	[4]	Correct answer
3 (i)	P(any other number) = $9/70$ P(X < 2) = $27/70 + 1/10$	B1		9/70 Seen
	= 34/70 (17/35) (0.486)	B1ft	[2]	Ft their probs if < 1
(ii)	E(X) = 108/70 (54/35) (1.543)	M1		Valid attempt at E(X) (needn't be accurate)
	$Var(X) = ((-2)^2 + + 5^2) \times 9 / 70 - (54/35)^2$	M1		Using a variance formula correctly with mean <sup>2</sup> subtracted numerically, no extra
	= 5.33	A1	[3]	division Correct final answer
(iii)	a=1	B1	[1]	
4 (i)	Options 5 bat 5 bl 1 Wk in	M1		Multiplying three combinations together
<b>\</b> /	$^{10}\text{C}_5 \times ^{9}\text{C}_5 \times ^{2}\text{C}_1 = 63504 \text{ ways}$ or 5 bat 4 bl 2 Wk in	M1		Summing more than one sensible option
	$^{10}\text{C}_5 \times ^{9}\text{C}_4 \times ^{2}\text{C}_2 = 31752 \text{ ways}$ or 6 bat 4 bl 1 Wk in			
	or 6 bat 4 bi 1 Wk in ${}^{10}C_6 \times {}^9C_4 \times {}^2C_1 = 52920$ ways	A1		Two options correct unsimplified
	Total = $148176 (148000)$	A1	[4]	Correct final answer
(ii)	$\frac{11!}{5!4!2!} = 6930$	B1	[1]	Correct answer evaluated
(iii)	Omit a pen $\frac{10!}{4!4!2!} = 3150$	M1		Summing three options
	Omit a diary $\frac{10!}{5!3!2!} = 2520$	B1		One option correct
	Omit a notebook $\frac{10!}{5!4!} = 1260$			
	Total = 6930	A1	[3]	Correct final answer
		1		L

		www.
Page 5	Mark Scheme: Teachers' version	Syllabus er
	GCE AS/A LEVEL – May/June 2011	9709

		<del></del>	72
5 (a)	$z > \frac{2\mu - \mu}{\sigma} = \frac{\mu}{\sigma} = \frac{7\sigma^2}{3\sigma}$	M1	Standardising attempt resulting in $z > \text{some } \mu/\sigma$ Substituting to eliminate $\mu$ or $\sigma$
	$\frac{7\sigma}{3} = 1.272$	M1 B1	Substituting to eliminate $\mu$ or $\sigma$ 1.272 seen
	$\sigma = 0.545$ $\mu = 0.693$	A1 <b>[4]</b>	Both answers correct
(b)	P(X < a + 33) = 0.75 z = 0.674	M1 A1	Using 0.75 oe ± 0.674 seen
	$\frac{a+33-33}{\sqrt{21}} = 0.674$	M1	Standardising, no cc, must have sq rt
	a = 3.09	A1 [4]	Correct answer
6 (i)	<u> </u>	M1	Sensible attempt at graph using u.c.b.
		M1 (Indpt)	2500 seen in median attempt on a CF graph Can be implied
	pupils		
	Median 270	A1 [3]	Correct answer + or – 5
(ii)	20% less than	M1	Using 20%
(232)	160	A1 [2]	Correct answer + or – 5
	2100 - 1600 = 500 $(50.5 \times 200 + 125.5 \times 600 + 175.5 \times 800 +$	B1 [1] M1	Using an attempt at mid-points
(11)	225.5 × 500 + 300.5 × 2000 + 400.5 × 600 +	M1	Using an attempt at frequencies
	525.5 × 300) / 5000 = 268	A1 A1 <b>[4]</b>	Correct mid-points or frequencies Correct answer only
		[T]	
7 (a)	(i) P(at least one 3) = 1 – P(no 3s) = $1 - (5/6)^9$	M1	Using 1 – none
7 (a)	$= 1 - (5/6)^9$ = 0.806	A1 [2]	
7 (a)	$=1-(5/6)^9$		Using 1 – none Correct answer  Equation or inequality involving <i>n</i> and 0.9 Solving attempt of sensible equation, can
7 (a)	$= 1 - (5/6)^{9}$ $= 0.806$ (ii) P(at least 1 three) = 1 - (5/6)^{n} 1 - (5/6)^{n} > 0.9	A1 [2] B1 M1	Using 1 – none Correct answer  Equation or inequality involving <i>n</i> and 0.9
	$= 1 - (5/6)^{9}$ $= 0.806$ (ii) P(at least 1 three) = 1 - (5/6)^{n} $1 - (5/6)^{n} > 0.9$ $n > 12.6$	A1 [2] B1 M1 M1	Using 1 – none Correct answer  Equation or inequality involving <i>n</i> and 0.9 Solving attempt of sensible equation, can be trial
	$= 1 - (5/6)^{9}$ $= 0.806$ (ii) P(at least 1 three) = 1 - (5/6)^{n} 1 - (5/6)^{n} > 0.9 n > 12.6 n = 13 P(R wins his 1 <sup>st</sup> ball) = P(GY)	A1 [2] B1 M1 M1 A1 [4]	Using 1 – none Correct answer  Equation or inequality involving <i>n</i> and 0.9 Solving attempt of sensible equation, can be trial Correct answer