

Cambridge International AS Level

MATHEMATICS

Paper 2 Pure Mathematics 2

MARK SCHEME

Maximum Mark: 50

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the February/March 2024 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

Cambridge International AS Level - Mark Scheme

PUBLISHED

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptions for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

Mathematics Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- 3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- 6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

PUBLISHED

Mark Scheme Notes

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- **B** Mark for a correct result or statement independent of method marks.
- **DM** or **DB** When a part of a question has two or more 'method' steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly, when there are several B marks allocated. The notation DM or DB is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
 - FT Implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only.
- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT above).
- For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures or would be correct to 3 significant figures if rounded (1 decimal place for angles in degrees).
- The total number of marks available for each question is shown at the bottom of the Marks column.
- Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
- Square brackets [] around text or numbers show extra information not needed for the mark to be awarded.

PUBLISHED

Abbreviations

AEF/OE Any Equivalent Form (of answer is equally acceptable) / Or Equivalent

AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)

CAO Correct Answer Only (emphasising that no 'follow through' from a previous error is allowed)

CWO Correct Working Only

ISW Ignore Subsequent Working

SOI Seen Or Implied

SC Special Case (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the

light of a particular circumstance)

WWW Without Wrong Working

AWRT Answer Which Rounds To

Question	Answer	Marks	Guidance
1	Apply logarithms to both sides and apply power law at least once	*M1	
	Obtain $(4x+3)\ln 3 = (2x+7)\ln 5$	A1	Or equivalent with x not in a power.
	Attempt solution of linear equation	DM1	
	Obtain 6.78	A1	Or greater accuracy.
		4	

Question	Answer	Marks	Guidance
2(a)	Draw V-shaped graph with vertex on positive x-axis	B1	8- 4- 2- 0 2 4 6
	State $(\frac{7}{3}, 0)$ and $(0, 7)$	B1	Allow if only $\frac{7}{3}$ and 7 shown on relevant axes.
		2	
2(b)	State or imply that gradient of left-hand part of graph is -3	B1	
	State $-3 < k < 0$	B1	Using < and not ≤.
		2	

Question	Answer	Marks	Guidance
3(a)	Substitute $x = \frac{1}{2}$, equate to zero and attempt solution	M1	
	Obtain $a=31$	A1	
	Divide by $2x-1$ at least as far as $3x^2 + mx$	M1	Or equivalent (for example by inspection,).
	Obtain $3x^2 + 17x + 10$	A1	
	Obtain $(2x-1)(3x+2)(x+5)$	A1	
		5	
3(b)	Attempt solution of $\sin \theta = k$ where k is valid constant from answer to part (a)	M1	
	Obtain -11.5	A1	Or greater accuracy.
		2	

Question	Answer	Marks	Guidance
4(a)	Use y-values $\sqrt{2}$, $\sqrt{1+e}$, $\sqrt{1+e^2}$, $\sqrt{1+e^3}$	B1	Or decimal equivalents
	Use correct formula, or equivalent, with $h=2$	M1	
	Obtain 15.7	A1	Or greater accuracy.
		3	

Question	Answer	Marks	Guidance
4(b)	State or imply that volume is $\pi \int (1 + e^{0.5x}) dx$	B1	Implied if π appears only at the end.
	Integrate to obtain form $k_1x + k_2e^{0.5x}$	M1	Where $k_1 k_2 \neq 0$ with or without π .
	Obtain $\pi(x + 2e^{0.5x})$ or $x + 2e^{0.5x}$	A1	
	Obtain $4\pi + 2\pi e^3$	A1	Or exact equivalent.
		4	

Question	Answer	Marks	Guidance
5(a)	Differentiate using quotient rule	*M1	OE
	Obtain $\frac{(x+2)3x^2 - x^3}{(x+2)^2}$	A1	OE
	Equate first derivative to 6 and simplify at least as far as $x^3 =$	DM1	
	Confirm $x = \sqrt[3]{12x + 12}$	A1	Answer given – necessary detail needed.
		4	
5(b)	Consider sign of $x - \sqrt[3]{12x + 12}$, or equivalent, for 3.8 and 4.0	M1	
	Obtain -0.06 and 0.08, or equivalents, and justify conclusion	A1	Answer given – necessary detail needed.
		2	

Question	Answer	Marks	Guidance
5(c)	Use iterative process correctly at least once	M1	
	Obtain final answer 3.88	A1	Answer required to exactly 3 sf.
	Show sufficient iterations to 5 sf to justify answer or show sign change in the interval [3.875, 3.885]	A1	
		3	

Question	Answer	Marks	Guidance
6(a)	Use product rule to find $\frac{dy}{dt}$	M1	Or chain rule if expression expanded before differentiation.
	Obtain $\frac{dy}{dt} = \frac{2\ln t}{t} - \frac{1}{t}$	A1	
	Divide attempt at $\frac{dy}{dt}$ by attempt at $\frac{dx}{dt}$	M1	
	Confirm $\frac{dy}{dx} = \frac{4\ln t - 2}{\sqrt{t}}$	A1	Answer given – necessary detail needed.
		4	
6(b)	State or imply $t = e^3$ at point B	B1	
	Obtain gradient $10e^{-\frac{3}{2}}$ or exact equivalent	B1	
		2	

Question	Answer	Marks	Guidance
6(c)	Equate first derivative to zero and obtain value for $\ln t$ or t	M1	
	Obtain x-coordinate $1 + e^{\frac{1}{4}}$	A1	Or exact equivalent.
	Obtain y-coordinate $-\frac{25}{4}$	A1	Or equivalent.
		3	

Question	Answer	Marks	Guidance
7(a)	Use correct identity for $\sin 2\theta$ or for $\cot \theta$	B1	
	Express left-hand side in terms of $\sin \theta$ and $\cos \theta$	M1	
	Attempt to express $k_1 \cos^2 \theta + k_2 \sin^2 \theta$ in terms of $\cos 2\theta$ only	M1	
	Confirm $a+b+(a-b)\cos 2\theta$	A1	Answer given – necessary detail needed.
		4	
7(b)	Obtain integrand $8+2\cos 2\theta$	B1	
	Integrate to obtain $8\theta + \sin 2\theta$	B1	
	Apply limits correctly to obtain $\frac{2}{3}\pi + \frac{1}{2}\sqrt{3} - \frac{1}{2}$	B1	Or exact equivalent.
		3	

Question	Answer	Marks	Guidance
7(c)	Use identity to obtain $5\cos\frac{2}{3}\alpha = -2$	B1	
	Obtain $\alpha = 2.97$	B1	Or greater accuracy.
	Obtain $\alpha = -2.97$	B1	Or greater accuracy; and no others between $-\pi$ and π .
		3	