

UNIVERSITY of CAMBRIDGE International Examinations

SYLLABUS

Cambridge International AS and A Level Mathematics **9709** Munan Banacambridge Cal Interna AS & A Le

For examination in June and November 2014



University of Cambridge International Examinations retains the copyright on all its publications. Registered Centres are permitted to copy material from this booklet for their own internal use. However, we cannot give permission to Centres to photocopy any material that is acknowledged to a third party even for internal use within a Centre.

© University of Cambridge International Examinations 2011

Contents

	ontents
С	ontents
1	
1.	Introduction 2 1.1 Why choose Cambridge? 2 1.2 Why choose Cambridge International AS and A Level? 2 1.3 Why choose Cambridge International AS and A Level Mathematics? 2 1.4 Cambridge AICE (Advanced International Certificate of Education) Diploma 1.5 How can I find out more?
2.	Assessment at a glance
3.	Syllabus aims and objectives
4.	Curriculum content
5.	Resource list
6.	List of formulae and tables of the normal distribution
7.	Mathematical notation
8.	Additional information

Introduction 1.

Why choose Cambridge? 1.1

www.papacambridge.com University of Cambridge International Examinations is the world's largest provider of international education programmes and qualifications for 5 to 19 year olds. We are part of the University of Cambridge, trusted for excellence in education. Our qualifications are recognised by the world's universities and employers.

Recognition

A Cambridge International AS or A Level is recognised around the world by schools, universities and employers. The qualifications are accepted as proof of academic ability for entry to universities worldwide, though some courses do require specific subjects.

Cambridge International A Levels typically take two years to complete and offer a flexible course of study that gives students the freedom to select subjects that are right for them. Cambridge International AS Levels often represent the first half of an A Level course but may also be taken as a freestanding qualification. They are accepted in all UK universities and carry half the weighting of an A Level. University course credit and advanced standing is often available for Cambridge International A/AS Levels in countries such as the USA and Canada.

Learn more at www.cie.org.uk/recognition

Excellence in education

We understand education. We work with over 9000 schools in over 160 countries who offer our programmes and gualifications. Understanding learners' needs around the world means listening carefully to our community of schools, and we are pleased that 98 % of Cambridge schools say they would recommend us to other schools.

Our mission is to provide excellence in education, and our vision is that Cambridge learners become confident, responsible, innovative and engaged.

Cambridge programmes and qualifications help Cambridge learners to become:

- confident in working with information and ideas their own and those of others
- responsible for themselves, responsive to and respectful of others
- innovative and equipped for new and future challenges
- engaged intellectually and socially, ready to make a difference.

Support in the classroom

We provide a world-class support service for Cambridge teachers and exams officers. We offer a wide range of teacher materials to Cambridge schools, plus teacher training (online and face-to-face), expert advice and learner-support materials. Exams officers can trust in reliable, efficient administration of exams entry and excellent, personal support from our customer services. Learn more at www.cie.org.uk/teachers

Not-for-profit, part of the University of Cambridge

We are a part of Cambridge Assessment, a department of the University of Cambridge and a not-for-profit organisation.

Ma invest constantly in research and development to improve our programmes and qualifications

1.2 Why choose Cambridge International AS and A Level?

www.papaCambridge.com Cambridge International AS and A Levels have a proven reputation for preparing students well for univers employment and life. They help develop the in-depth subject knowledge and understanding which are so important to universities and employers.

You can offer almost any combination of 55 subjects. Students can specialise or study a range of subjects, ensuring breadth. Giving students the power to choose helps motivate them throughout their studies.

Cambridge International AS and A Level gives you building blocks to build an individualised curriculum that develops your learners' knowledge, understanding and skills in:

- in-depth subject content
- independent thinking •
- applying knowledge and understanding to new as well as familiar situations •
- handling and evaluating different types of information sources
- thinking logically and presenting ordered and coherent arguments ٠
- making judgements, recommendations and decisions ٠
- presenting reasoned explanations, understanding implications and communicating them clearly and logically
- working and communicating in English.

The syllabuses are international in outlook, but retain a local relevance. They have been created specifically for an international student body with content to suit a wide variety of schools and avoid cultural bias.

1.3 Why choose Cambridge International AS and A Level Mathematics?

Cambridge International AS and A Level Mathematics is accepted by universities and employers as proof of mathematical knowledge and understanding. Successful candidates gain lifelong skills, including:

- a deeper understanding of mathematical principles;
- the further development of mathematical skills including the use of applications of mathematics in the context of everyday situations and in other subjects that they may be studying;
- the ability to analyse problems logically, recognising when and how a situation may be represented mathematically;
- the use of mathematics as a means of communication;
- a solid foundation for further study.

The syllabus allows Centres flexibility to choose from three different routes to AS Level Mathematics - Pure Mathematics only or Pure Mathematics and Mechanics or Pure Mathematics and Probability and Statistics. Centres can choose from three different routes to Cambridge International A Level Mathematics depending on the choice of Mechanics, or Probability and Statistics, or both, in the broad area of 'applications'.

Cambridge AICE (Advanced International Certificate of 1.4 Education) Diploma

www.papacambridge.com Cambridge AICE (Advanced International Certificate of Education) Diploma is the group award of Cambridge International AS and A Level.

Cambridge AICE Diploma involves the selection of subjects from three curriculum groups - Mathematics and Science; Languages; Arts and Humanities.

A Cambridge International A Level counts as a double-credit qualification and a Cambridge International AS Level as a single-credit qualification within the Cambridge AICE Diploma award framework.

To be considered for an AICE Diploma, a candidate must earn the equivalent of six credits by passing a combination of examinations at either double credit or single credit, with at least one course coming from each of the three curriculum areas.

The AICE Diploma is comprised of examinations administered in May/June and October/November series each year.

Mathematics (9709) falls into Group 1, Mathematics and Sciences.

Learn more about the AICE Diploma at http://www.cie.org.uk/qualifications/academic/uppersec/aice

1.5 How can I find out more?

If you are already a Cambridge school

You can make entries for this qualification through your usual channels. If you have any questions, please contact us at international@cie.org.uk

If you are not yet a Cambridge school

Learn about the benefits of becoming a Cambridge school at www.cie.org.uk/startcambridge. Email us at international@cie.org.uk to find out how your organisation can become a Cambridge school.

2. Assessment at a glance

The 7 units in the scheme cover the following subject areas:

- Pure Mathematics (units P1, P2 and P3);
- Mechanics (units M1 and M2);
- Probability and Statistics (units S1 and S2).

Centres and candidates may:

• take all four Advanced (A) Level components in the same examination series for the full Cambridge International A Level;

www.papaCambridge.com

- follow a staged assessment route to the Cambridge International A Level by taking two Advanced Subsidiary (AS) papers (P1 & M1 or P1 & S1) in an earlier examination series;
- take the Advanced Subsidiary (AS) qualification only.

Cambridge International AS Level candidates take:

Paper 1: Pure Mathematics 1 (P1)

1¾ hours

About 10 shorter and longer questions

75 marks weighted at 60% of total

plus one of the following papers:

	Paper 6: Probability and Statistics 1 (S1)
t 7 shorter and longer ions	1¼ hours About 7 shorter and longer questions 50 marks weighted at 40%
	Durs t 7 shorter and longer ions arks weighted at 40% al

Cambridge International A Level candidates take:

nent at a glance	MMM. Dabac
Cambridge International A Level candid	lates take:
Paper 1: Pure Mathematics 1 (P1)	Paper 3 Pure Mathematics 3 (P3)
1¾ hours	1¾ hours
About 10 shorter and longer questions	About 10 shorter and longer questions
75 marks weighted at 30% of total	75 marks weighted at 30% of total

plus **one** of the following combinations of two papers:

Paper 4: Mechanics 1 (M1)	Paper 6: Probability and Statistics 1 (S1)
1¼ hours	1¼ hours
About 7 shorter and longer questions	About 7 shorter and longer questions
50 marks weighted at 20% of total	50 marks weighted at 20% of total

or

Paper 4: Mechanics 1 (M1)	Paper 5: Mechanics 2 (M2)
1¼ hours	1¼ hours
About 7 shorter and longer questions	About 7 shorter and longer questions
50 marks weighted at 20% of total	50 marks weighted at 20% of total

or

Paper 6: Probability and Statistics 1 (S1)	Paper 7: Probability and Statistics 2 (S2)
1¼ hours	1¼ hours
About 7 shorter and longer questions	About 7 shorter and longer questions
50 marks weighted at 20% of total	50 marks weighted at 20% of total

Question papers

www.papaCambridge.com There is no choice of questions in any of the question papers and questions will be arranged approximate in order of increasing mark allocations.

It is expected that candidates will have a calculator with standard 'scientific' functions available for use for all papers in the examination. Computers, and calculators capable of algebraic manipulation, are not permitted.

A list of formulae and tables of the normal distribution (MF9) is supplied for the use of candidates in the examination. Details of the items in this list are given for reference in Section 6.

Relationships between units

Units P2, M2, S2 are sequential to units P1, M1, S1 respectively, and the later unit in each subject area may not be used for certification unless the corresponding earlier unit is being (or has already been) used.

Unit P3 is also sequential to unit P1, and may not be used for certification unless P1 is being (or has already been) used. The subject content of unit P2 is a subset of the subject content of unit P3; otherwise, the subject content for different units does not overlap, although later units in each subject area assume knowledge of the earlier units.

Availability

This syllabus is examined in the May/June examination series and the October/November examination series.

This syllabus is available to private candidates.

Centres in the UK that receive government funding are advised to consult the Cambridge website **www.cie.org.uk** for the latest information before beginning to teach this syllabus.

Combining this with other syllabuses

Candidates can combine this syllabus in an examination series with any other Cambridge syllabus, except:

• syllabuses with the same title at the same level

Syllabus aims and objectives 3.

3.1 Aims

www.papaCambridge.com The aims of the syllabus are the same for all students. These are set out below and describe the educational purposes of any course based on the Mathematics units for the Cambridge International AS and A Level examinations. The aims are not listed in order of priority.

The aims are to enable candidates to:

- develop their mathematical knowledge and skills in a way which encourages confidence and provides satisfaction and enjoyment;
- develop an understanding of mathematical principles and an appreciation of mathematics as a logical and coherent subject;
- acquire a range of mathematical skills, particularly those which will enable them to use applications of mathematics in the context of everyday situations and of other subjects they may be studying;
- develop the ability to analyse problems logically, recognise when and how a situation may be represented mathematically, identify and interpret relevant factors and, where necessary, select an appropriate mathematical method to solve the problem;
- use mathematics as a means of communication with emphasis on the use of clear expression;
- acquire the mathematical background necessary for further study in this or related subjects.

Assessment objectives 3.2

The abilities assessed in the examinations cover a single area: technique with application.

The examination will test the ability of candidates to:

- understand relevant mathematical concepts, terminology and notation; .
- recall accurately and use successfully appropriate manipulative techniques;
- recognise the appropriate mathematical procedure for a given situation; .
- apply combinations of mathematical skills and techniques in solving problems;
- present mathematical work, and communicate conclusions, in a clear and logical way.

Curriculum content 4.

www.papacambridge.com The mathematical content for each unit in the scheme is detailed below. The order in which topics are listed is not intended to imply anything about the order in which they might be taught.

As well as demonstrating skill in the appropriate techniques, candidates will be expected to apply their knowledge in the solution of problems. Individual questions set may involve ideas and methods from more than one section of the relevant content list.

For all units, knowledge of the content of Cambridge O Level/Cambridge IGCSE Mathematics is assumed. Candidates will be expected to be familiar with scientific notation for the expression of compound units, e.g. 5 m s⁻¹ for 5 metres per second.

Unit P1: Pure Math	ematics 1 (Paper 1)
	Candidates should be able to:
1. Quadratics	 carry out the process of completing the square for a quadratic polynomial ax² + bx + c, and use this form, e.g. to locate the vertex of the graph of y = ax² + bx + c or to sketch the graph; find the discriminant of a quadratic polynomial ax² + bx + c and use the discriminant, e.g. to determine the number of real roots of the equation ax² + bx + c = 0; solve quadratic equations, and linear and quadratic inequalities, in one unknown; solve by substitution a pair of simultaneous equations of which one is linear and one is quadratic; recognise and solve equations in <i>x</i> which are quadratic in some function of <i>x</i>, e.g. x⁴ - 5x² + 4 = 0.
2. Functions	 understand the terms function, domain, range, one-one function, inverse function and composition of functions; identify the range of a given function in simple cases, and find the composition of two given functions; determine whether or not a given function is one-one, and find the inverse of a one-one function in simple cases; illustrate in graphical terms the relation between a one-one function and its inverse.

n content	MANN Papac
3. Coordinate geometry	 find the length, gradient and mid-point of a line segment, given the coordinates of the end-points; find the equation of a straight line given sufficient information (e.g. the coordinates of two points on it, or one point on it and its gradient); understand and use the relationships between the gradients of parallel and perpendicular lines; interpret and use linear equations, particularly the forms y = mx + c and y - y₁ = m(x - x₁); understand the relationship between a graph and its associated algebraic equation, and use the relationship between points of intersection of graphs and solutions of equations (including, in simple cases, the correspondence between a line being tangent to a curve and a repeated root of an equation).
4. Circular measure	 understand the definition of a radian, and use the relationship between radians and degrees; use the formulae s = rθ and A = ¹/₂r²θ in solving problems concerning the arc length and sector area of a circle.
5. Trigonometry	 sketch and use graphs of the sine, cosine and tangent functions (for angles of any size, and using either degrees or radians); use the exact values of the sine, cosine and tangent of 30°, 45°, 60°, and related angles, e.g. cos 150° = -1/2 √3; use the notations sin⁻¹x, cos⁻¹x, tan⁻¹x to denote the principal values of the inverse trigonometric relations; use the identities sin θ/cos θ = tan θ and sin² θ + cos² θ = 1; find all the solutions of simple trigonometrical equations lying in a specified interval (general forms of solution are not included).
6. Vectors	 use standard notations for vectors, i.e. \$\begin{pmatrix} x \begin{pmatrix} y \begin{pmatrix} x \begin{pmatrix} x \begin{pmatrix} y \begin{pmatrix} x \begin{pmatrix}

	• use the expansion of $(a + b)^n$, where <i>n</i> is a positive integer (knowledge of the greatest term and properties of the coefficients are not required, but the notations $\binom{n}{r}$ and <i>n</i> ! should be known);
7. Series	 use the expansion of (a + b)ⁿ, where n is a positive integer (knowledge of the greatest term and properties of the coefficients are not required, but the notations (n) and n! should be known); recognise arithmetic and geometric progressions; use the formulae for the <i>n</i>th term and for the sum of the first <i>n</i> terms to solve problems involving arithmetic or geometric progressions; use the condition for the convergence of a geometric progression, and the formula for the sum to infinity of a convergent geometric progression.
8. Differentiation	 understand the idea of the gradient of a curve, and use the notations f'(x), f''(x), dy/dx and d²y/dx² (the technique of differentiation from first principles is not required); use the derivative of xⁿ (for any rational <i>n</i>), together with constant multiples, sums, differences of functions, and of composite functions using the chain rule; apply differentiation to gradients, tangents and normals, increasing and decreasing functions and rates of change (including connected rates of change); locate stationary points, and use information about stationary points in sketching graphs (the ability to distinguish between maximum points and minimum points is required, but identification of points of inflexion is not included).
9. Integration	 understand integration as the reverse process of differentiation, and integrate (ax + b)ⁿ (for any rational n except −1), together with constant multiples, sums and differences; solve problems involving the evaluation of a constant of integration, e.g. to find the equation of the curve through (1, -2) for which

Knowledge of the con	atics 2 (Paper 2) tent of unit P1 is assumed, and candidates may be required to owledge in answering questions.
	Candidates should be able to:
1. Algebra	 understand the meaning of x , and use relations such as a = b ⇔ a² = b² and x - a < b ⇔ a - b < x < a + b in the course of solving equations and inequalities; divide a polynomial, of degree not exceeding 4, by a linear or quadratic polynomial, and identify the quotient and remainder (which may be zero); use the factor theorem and the remainder theorem, e.g. to find factors, solve polynomial equations or evaluate unknown coefficients.
2. Logarithmic and exponential functions	 understand the relationship between logarithms and indices, and use the laws of logarithms (excluding change of base); understand the definition and properties of e^x and ln x, including their relationship as inverse functions and their graphs; use logarithms to solve equations of the form a^x = b, and similar inequalities; use logarithms to transform a given relationship to linear form, and hence determine unknown constants by considering the gradient and/ or intercept.
3. Trigonometry	 understand the relationship of the secant, cosecant and cotangent functions to cosine, sine and tangent, and use properties and graphs of all six trigonometric functions for angles of any magnitude; use trigonometrical identities for the simplification and exact evaluation of expressions and in the course of solving equations, and select an identity or identities appropriate to the context, showing familiarity in particular with the use of: sec²θ = 1 + tan²θ and cosec²θ = 1 + cot²θ, the expansions of sin(A ± B), cos(A ± B) and tan(A ± B), the formulae for sin 2A, cos 2A and tan 2A, the expressions of a sin θ + b cos θ in the forms R sin(θ ± α) and R cos(θ ± α).
4. Differentiation	 use the derivatives of e^x, ln x, sin x, cos x, tan x, together with constant multiples, sums, differences and composites; differentiate products and quotients; find and use the first derivative of a function which is defined parametrically or implicitly.

	• extend the idea of 'reverse differentiation' to include the integration e^{ax+b} , $\frac{1}{ax+b}$, $sin(ax+b)$, $cos(ax+b)$ and $sec^2(ax+b)$ (knowledge of the general method of integration by substitution is not required);
5. Integration	 extend the idea of 'reverse differentiation' to include the integration e^{ax+b}, 1/(ax + b), cos(ax + b) and sec² (ax + b) (knowledge of the general method of integration by substitution is not required); use trigonometrical relationships (such as double-angle formulae) to facilitate the integration of functions such as cos² x; use the trapezium rule to estimate the value of a definite integral, and use sketch graphs in simple cases to determine whether the trapezium rule gives an over-estimate or an under-estimate.
6. Numerical solution of equations	 locate approximately a root of an equation, by means of graphical considerations and/or searching for a sign change; understand the idea of, and use the notation for, a sequence of approximations which converges to a root of an equation; understand how a given simple iterative formula of the form <i>x</i>_{n+1} = F(<i>x</i>_n) relates to the equation being solved, and use a given iteration, or an iteration based on a given rearrangement of an equation, to determine a root to a prescribed degree of accuracy (knowledge of the condition for convergence is not included, but candidates should understand that an iteration may fail to converge).

	·C	
Content Unit P3: Pure Mathematics 3 (Paper 3) Knowledge of the content of unit P1 is assumed, and candidates may be required to demonstrate such knowledge in answering questions.		
	Candidates should be able to:	
1. Algebra	 understand the meaning of x , and use relations such as a = b ⇔ a² = b² and x - a < b ⇔ a - b < x < a + b in the course of solving equations and inequalities; 	
	 divide a polynomial, of degree not exceeding 4, by a linear or quadratic polynomial, and identify the quotient and remainder (which may be zero); 	
	 use the factor theorem and the remainder theorem, e.g. to find factors, solve polynomial equations or evaluate unknown coefficients; 	
	 recall an appropriate form for expressing rational functions in partial fractions, and carry out the decomposition, in cases where the denominator is no more complicated than: 	
	• $(ax+b)(cx+d)(ex+f),$	
	• $(ax + b)(cx + d)^2$,	
	• $(ax + b)(x^2 + c^2)$,	
	and where the degree of the numerator does not exceed that of the denominator;	
	• use the expansion of $(1 + x)^n$, where <i>n</i> is a rational number and $ x < 1$ (finding a general term is not included, but adapting the standard series to expand e.g. $(2 - \frac{1}{2}x)^{-1}$ is included).	
2. Logarithmic and exponential	 understand the relationship between logarithms and indices, and use the laws of logarithms (excluding change of base); 	
and exponential functions	 understand the definition and properties of e^x and ln x, including their relationship as inverse functions and their graphs; 	
	 use logarithms to solve equations of the form a^x = b, and similar inequalities; 	
	 use logarithms to transform a given relationship to linear form, and hence determine unknown constants by considering the gradient and/ or intercept. 	

	Man Curriculum
3. Trigonometry	 understand the relationship of the secant, cosecant and cotangent functions to cosine, sine and tangent, and use properties and graphs of all six trigonometric functions for angles of any magnitude; use trigonometrical identities for the simplification and exact evaluation of expressions and in the course of solving equations, and select an identity or identities appropriate to the context, showing familiarity in particular with the use of: sec² θ ≡ 1 + tan² θ and cosec² θ ≡ 1 + cot² θ, the expansions of sin(A ± B), cos(A ± B) and tan(A ± B), the formulae for sin 2A, cos 2A and tan 2A, the expressions of a sin θ + b cos θ in the forms R sin(θ ± a) and R cos(θ ± a).
4. Differentiation	 use the derivatives of e^x, ln x, sin x, cos x, tan x, together with constant multiples, sums, differences and composites; differentiate products and quotients; find and use the first derivative of a function which is defined parametrically or implicitly.
5. Integration	 extend the idea of 'reverse differentiation' to include the integration of e^{ax+b}, 1/(ax + b), cis(ax + b) and sec²(ax + b); use trigonometrical relationships (such as double-angle formulae) to facilitate the integration of functions such as cos² x; integrate rational functions by means of decomposition into partial fractions (restricted to the types of partial fractions specified in paragraph 1 above); recognise an integrand of the form kt'(x)/f(x), and integrate, for example, x² + 1 or tan x; recognise when an integrand can usefully be regarded as a product, and use integration by parts to integrate, for example, x sin 2x, x² e^x or ln x; use a given substitution to simplify and evaluate either a definite or an indefinite integral; use the trapezium rule to estimate the value of a definite integral, and use sketch graphs in simple cases to determine whether the trapezium rule gives an over-estimate or an under-estimate.

content	MANN, Papac
6. Numerical solution of equations	 locate approximately a root of an equation, by means of graphical considerations and/or searching for a sign change; understand the idea of, and use the notation for, a sequence of approximations which converges to a root of an equation; understand how a given simple iterative formula of the form <i>x</i>_{n+1} = F(<i>x</i>_n) relates to the equation being solved, and use a given iteration, or an iteration based on a given rearrangement of an equation, to determine a root to a prescribed degree of accuracy (knowledge of the condition for convergence is not included, but candidates should understand that an iteration may fail to converge).
7. Vectors	 understand the significance of all the symbols used when the equation of a straight line is expressed in the form r = a + tb; determine whether two lines are parallel, intersect or are skew; find the angle between two lines, and the point of intersection of two lines when it exists; understand the significance of all the symbols used when the equation of a plane is expressed in either of the forms ax + by + cz = d or (r - a).n = 0; use equations of lines and planes to solve problems concerning distances, angles and intersections, and in particular: find the equation of a line or a plane, given sufficient information, determine whether a line lies in a plane, is parallel to a plane, or intersects a plane, and find the point of intersection of a line and a plane when it exists, find the line of intersection of two non-parallel planes, find the angle between two planes, and the angle between a line and a plane.
8. Differential equations	 formulate a simple statement involving a rate of change as a differential equation, including the introduction if necessary of a constant of proportionality; find by integration a general form of solution for a first order differential equation in which the variables are separable; use an initial condition to find a particular solution; interpret the solution of a differential equation in the context of a problem being modelled by the equation.

	 understand the idea of a complex number, recall the meaning of the terms real part, imaginary part, modulus, argument, conjugate, and use the fact that two complex numbers are equal if and only if both real and imaginary parts are equal;
9. Complex numbers	 understand the idea of a complex number, recall the meaning of the terms real part, imaginary part, modulus, argument, conjugate, and use the fact that two complex numbers are equal if and only if both real and imaginary parts are equal;
	 carry out operations of addition, subtraction, multiplication and division of two complex numbers expressed in cartesian form x + iy;
	 use the result that, for a polynomial equation with real coefficients, any non-real roots occur in conjugate pairs;
	 represent complex numbers geometrically by means of an Argand diagram;
	• carry out operations of multiplication and division of two complex numbers expressed in polar form $r(\cos \theta + i \sin \theta) \equiv r e^{i\theta}$;
	• find the two square roots of a complex number;
	 understand in simple terms the geometrical effects of conjugating a complex number and of adding, subtracting, multiplying and dividing two complex numbers;
	 illustrate simple equations and inequalities involving complex numbers by means of loci in an Argand diagram, e.g. z - a < k, z - a = z - b , arg(z - a) = α.

Unit M1: Mechanics 1 (Paper 4)

www.papacambridge.com Questions set will be mainly numerical, and will aim to test mechanical principles without involving difficult algebra or trigonometry. However, candidates should be familiar in particular with the following trigonometrical results: $\sin(90^\circ - \theta) \equiv \cos\theta$, $\cos(90^\circ - \theta) \equiv \sin\theta$, $\tan\theta \equiv \frac{\sin\theta}{\cos\theta}$, $\sin^2\theta + \cos^2\theta \equiv 1$.

Vector notation will not be used in the question papers, but candidates may use vector methods in their solutions if they wish.

In the following content list, reference to the equilibrium or motion of a 'particle' is not intended to exclude questions that involve extended bodies in a 'realistic' context; however, it is to be understood that any such bodies are to be treated as particles for the purposes of the question.

Unit M1: Mechanics 1 (Paper 4)			
	Candidates should be able to:		
1. Forces and equilibrium	 identify the forces acting in a given situation; understand the vector nature of force, and find and use components and resultants; use the principle that, when a particle is in equilibrium, the vector sum of the forces acting is zero, or equivalently, that the sum of the components in any direction is zero; understand that a contact force between two surfaces can be represented by two components, the normal component and the frictional component; use the model of a 'smooth' contact, and understand the limitations of this model; understand the concepts of limiting friction and limiting equilibrium; recall the definition of coefficient of friction, and use the relationship <i>F</i> = μ<i>R</i> or <i>F</i> ≤ μ<i>R</i>, as appropriate; use Newton's third law. 		
2. Kinematics of motion in a straight line	 understand the concepts of distance and speed as scalar quantities, and of displacement, velocity and acceleration as vector quantities (in one dimension only); sketch and interpret displacement-time graphs and velocity-time graphs, and in particular appreciate that: the area under a velocity-time graph represents displacement, the gradient of a displacement-time graph represents velocity, the gradient of a velocity-time graph represents acceleration; use differentiation and integration with respect to time to solve simple problems concerning displacement, velocity and acceleration (restricted to calculus within the scope of unit P1); use appropriate formulae for motion with constant acceleration in a straight line. 		

	 apply Newton's laws of motion to the linear motion of a particle of constant mass moving under the action of constant forces, which may include friction; use the relationship between mass and weight; solve simple problems which may be modelled as the motion of
3. Newton's laws of motion	a particle moving vertically or on an inclined plane with constant acceleration;
	 solve simple problems which may be modelled as the motion of two particles, connected by a light inextensible string which may pass over a fixed smooth peg or light pulley.
4. Energy, work and power	 understand the concept of the work done by a force, and calculate the work done by a constant force when its point of application undergoes a displacement not necessarily parallel to the force (use of the scalar product is not required); understand the concepts of gravitational potential energy and kinetic energy, and use appropriate formulae;
	 understand and use the relationship between the change in energy of a system and the work done by the external forces, and use in appropriate cases the principle of conservation of energy;
	 use the definition of power as the rate at which a force does work, and use the relationship between power, force and velocity for a force acting in the direction of motion;
	 solve problems involving, for example, the instantaneous acceleration of a car moving on a hill with resistance.

	2 (Paper 5) ntent of unit M1 is assumed, and candidates may be required to nowledge in answering questions.
Motion of a projectile	 model the motion of a projectile as a particle moving with constant acceleration and understand any limitations of the model; use horizontal and vertical equations of motion to solve problems on the motion of projectiles, including finding the magnitude and direction of the velocity at a given time of position, the range on a horizontal plane and the greatest height reached;
	 derive and use the cartesian equations of the trajectory of a projectile, including problems in which the initial speed and/or angle of projection may be unknown.
Equilibrium of a rigid body	 calculate the moment of a force about a point, in two dimensional situations only (understanding of the vector nature of moments is not required);
	 use the result that the effect of gravity on a rigid body is equivalent to a single force acting at the centre of mass of the body, and identify the position of the centre of mass of a uniform body using considerations of symmetry;
	 use given information about the position of the centre of mass of a triangular lamina and other simple shapes;
	 determine the position of the centre of mass of a composite body by considering an equivalent system of particles (in simple cases only, e.g. a uniform L-shaped lamina);
	• use the principle that if a rigid body is in equilibrium under the action of coplanar forces then the vector sum of the forces is zero and the sum of the moments of the forces about any point is zero, and the converse of this;
	 solve problems involving the equilibrium of a single rigid body under the action of coplanar forces, including those involving toppling or sliding (problems set will not involve complicated trigonometry).
Uniform motion n a circle	• understand the concept of angular speed for a particle moving in a circle, and use the relation $v = r\omega$;
	 understand that the acceleration of a particle moving in a circle with constant speed is directed towards the centre of the circle, and use
	 the formulae rω² and V²/r; solve problems which can be modelled by the motion of a particle moving in a horizontal circle with constant speed.

	 use Hooke's law as a model relating the force in an elastic string or spring to the extension or compression, and understand the term modulus of elasticity; use the formula for the elastic potential energy stored in a string or
4. Hooke′s law	 use Hooke's law as a model relating the force in an elastic string or spring to the extension or compression, and understand the term modulus of elasticity;
	 use the formula for the elastic potential energy stored in a string or spring;
	 solve problems involving forces due to elastic strings or springs, including those where considerations of work and energy are needed.
5. Linear motion under a variable force	 use dx/dt for velocity, and dv/dt or vdv/dx for acceleration, as appropriate; solve problems which can be modelled as the linear motion of a particle under the action of a variable force, by setting up and solving an appropriate differential equation (restricted to equations in which the variables are separable).

ontent	Statistics 1 (Paper 6) Candidates should be able to: • select a suitable way of presenting raw statistical data, and discuss	
Jnit S1: Probability &	a Statistics 1 (Paper 6)	
	Candidates should be able to:	
1. Representation of data	advantages and/or disadvantages that particular representations may have;	
	 construct and interpret stem-and-leaf diagrams, box-and-whisker plots, histograms and cumulative frequency graphs; 	
	 understand and use different measures of central tendency (mean, median, mode) and variation (range, interquartile range, standard deviation), e.g. in comparing and contrasting sets of data; 	
	 use a cumulative frequency graph to estimate the median value, the quartiles and the interquartile range of a set of data; 	
	• calculate the mean and standard deviation of a set of data (including grouped data) either from the data itself or from given totals such as Σx and Σx^2 , or $\Sigma(x - a)$ and $\Sigma(x - a)^2$.	
2. Permutations and	 understand the terms permutation and combination, and solve simple problems involving selections; 	
combinations	 solve problems about arrangements of objects in a line, including those involving: 	
	 repetition (e.g. the number of ways of arranging the letters of the word 'NEEDLESS'), 	
	 restriction (e.g. the number of ways several people can stand in a line if 2 particular people must — or must not — stand next to each other). 	
3. Probability	 evaluate probabilities in simple cases by means of enumeration of equiprobable elementary events (e.g. for the total score when two fair dice are thrown), or by calculation using permutations or combinations; 	
	 use addition and multiplication of probabilities, as appropriate, in simple cases; 	
	 understand the meaning of exclusive and independent events, and calculate and use conditional probabilities in simple cases, e.g. situations that can be represented by means of a tree diagram. 	
4. Discrete random variables	 construct a probability distribution table relating to a given situation involving a discrete random variable variable X, and calculate E(X) and Var(X); 	
	 use formulae for probabilities for the binomial distribution, and recognise practical situations where the binomial distribution is a suitable model (the notation B(n, p) is included); 	
	 use formulae for the expectation and variance of the binomial distribution. 	

	 understand the use of a normal distribution to model a continuous random variable, and use normal distribution tables; solve problems concerning a variable <i>X</i>, where <i>X</i> ~ N(μ, σ²), including: finding the value of P(<i>X</i> > <i>x</i>₁), or a related probability, given the
5. The normal distribution	 understand the use of a normal distribution to model a continuous random variable, and use normal distribution tables; solve problems concerning a variable <i>X</i>, where <i>X</i> ~ N(μ, σ²), including: finding the value of P(<i>X</i> > <i>x</i>₁), or a related probability, given the values of <i>x</i>₁, μ, σ, finding a relationship between <i>x</i>₁, μ and σ given the value of P(<i>X</i> > <i>x</i>₁) or a related probability; recall conditions under which the normal distribution can be used as an approximation to the binomial distribution (<i>n</i> large enough to ensure that <i>np</i> > 5 and <i>nq</i> > 5), and use this approximation, with a continuity correction, in solving problems.

(nowledge of the co	& Statistics 2 (Paper 7) Intent of unit S1 is assumed, and candidates may be required to howledge in answering questions.
	Candidates should be able to:
1. The Poisson distribution	 calculate probabilities for the distribution Po(μ); use the fact that if X ~ Po(μ) then the mean and variance of X are each equal to μ; understand the relevance of the Poisson distribution to the distribution of random events, and use the Poisson distribution as a model; use the Poisson distribution as an approximation to the binomial distribution where appropriate (n > 50 and np < 5, approximately); use the normal distribution, with continuity correction, as an approximation to the Poisson distribution where appropriate (μ > 15, approximately).
. Linear combinations of random variables	 use, in the course of solving problems, the results that: E(aX + b) = aE(X) + b and Var(aX + b) = a²Var(X) , E(aX + bY) = aE(X) + bE(Y), Var(aX + bY) = a²Var(X) + b²Var(Y) for independent X and Y, if X has a normal distribution then so does aX + b, if X and Y have independent normal distributions then aX + bY has a normal distribution, if X and Y have independent Poisson distributions then X + Y has a Poisson distribution.
3. Continuous random variables	 understand the concept of a continuous random variable, and recall and use properties of a probability density function (restricted to functions defined over a single interval); use a probability density function to solve problems involving probabilities, and to calculate the mean and variance of a distribution (explicit knowledge of the cumulative distribution function is not included, but location of the median, for example, in simple cases by direct consideration of an area may be required).

	 understand the distinction between a sample and a population, and appreciate the necessity for randomness in choosing samples; explain in simple terms why a given sampling method may be unsatisfactory (knowledge of particular sampling methods, such as guota or stratified sampling, is not required, but candidates should
4. Sampling and estimation	 understand the distinction between a sample and a population, and appreciate the necessity for randomness in choosing samples; explain in simple terms why a given sampling method may be unsatisfactory (knowledge of particular sampling methods, such as quota or stratified sampling, is not required, but candidates should have an elementary understanding of the use of random numbers in producing random samples); recognise that a sample mean can be regarded as a random variable, and use the facts that E(X̄) = μ and that Var(X̄) = σ²/n; use the fact that X̄ has a normal distribution if X has a normal distribution; use the Central Limit theorem where appropriate; calculate unbiased estimates of the population mean and variance from a sample, using either raw or summarised data (only a simple understanding of the term 'unbiased' is required); determine a confidence interval for a population mean in cases where the population is normally distributed with known variance or where a large sample is used; determine, from a large sample, an approximate confidence interval for a population proportion.
5. Hypothesis tests	 understand the nature of a hypothesis test, the difference between one-tail and two-tail tests, and the terms null hypothesis, alternative hypothesis, significance level, rejection region (or critical region), acceptance region and test statistic; formulate hypotheses and carry out a hypothesis test in the context of a single observation from a population which has a binomial or Poisson distribution, using either direct evaluation of probabilities or a normal approximation, as appropriate; formulate hypotheses and carry out a hypothesis test concerning the population mean in cases where the population is normally distributed with known variance or where a large sample is used; understand the terms Type I error and Type II errors in relation to hypothesis tests; calculate the probabilities of making Type I and Type II errors in specific situations involving tests based on a normal distribution or direct evaluation of binomial or Poisson probabilities.

Resource list 5.

www.papacambridge.com These titles represent some of the texts available in the UK at the time of printing this booklet. Teachers are encouraged to choose texts for class use which they feel will be of interest to their students. The inclusion of a text does not imply that it is either recommended or approved by Cambridge. Wherever possible, the International Standard Book Number (ISBN) is given.

Endorsed Textbooks

The following textbooks are endorsed by Cambridge for use with the syllabuses in this booklet - please contact Cambridge University Press for further information.

Author	Title	Publisher	ISBN
Neill & Quadling	Pure Mathematics 1	Cambridge University Press	0 521 53011 3
Neill & Quadling	Pure Mathematics 2 & 3	Cambridge University Press	0 521 53012 1
Quadling	Mechanics 1	Cambridge University Press	0 521 53015 6
Quadling	Mechanics 2	Cambridge University Press	0 521 53016 4
Dobbs & Miller	Statistics 1	Cambridge University Press	0 521 53013 X
Dobbs & Miller	Statistics 2	Cambridge University Press	0 521 53014 8

Suggested Books

Pure Mathematics

Author	Title	Publisher	ISBN
Backhouse, Houldsworth & Horrill	Pure Mathematics 1	Longman, 1985	0 582 35386 6
Backhouse, Houldsworth & Horrill	Pure Mathematics 2	Longman, 1985	0 582 35387 4
Backhouse, Houldsworth, Horrill & Wood	Essential Pure Mathematics	Longman, 1991	0582 066581
Bostock & Chandler	Core Maths for Advanced Level	Nelson Thornes, 2000	0 7487 5509 8
Butcher & Megeny	<i>Access to Advanced Level</i> <i>Maths</i> (short introductory course)	Nelson Thornes, 1997	0 7487 2999 2
Emanuel, Wood & Crawshaw	Pure Mathematics 1	Longman, 2001	0 582 40550 5
Emanuel, Wood & Crawshaw	Pure Mathematics 2	Longman, 2001	0 582 40549 1

			MANA Res
Hunt	<i>Graded Exercises in Pure</i> <i>Mathematics</i> (Practice questions)	Cambridge University Press, 2001	Рес мил. рана салир 0 521 637 0 7487 3558 5
Martin, Brown, Rigby & Riley	Complete Advanced Level Mathematics: Pure Mathematics: Core Text	Nelson Thornes, 2000	0 7487 3558 5
Morley	Practice for Advanced Mathematics – Pure Mathematics (Practice questions)	Hodder & Stoughton Educational, 1999	0 340 701676
Sadler & Thorning	Understanding Pure Mathematics	Oxford University Press, 1987	019 914243 2
Smedley & Wiseman	Introducing Pure Mathematics	Oxford University Press, 2001	0 19 914803 1
SMP	Mathematics for AS and A Level – Pure Mathematics	Cambridge University Press, 1997	0 521 56617 7
Solomon	Advanced Level Mathematics: Pure Mathematics	John Murray, 1995	0 7195 5344 X

Integrated Courses

Author	Title	Publisher	ISBN
Berry, Fentern, Francis & Graham	Discovering Advanced Mathematics – AS Mathematics	Collins Educational, 2000	0 00 322502 X
Berry, Fentern, Francis & Graham	Discovering Advanced Mathematics – A2 Mathematics	Collins Educational, 2001	0 00 322503 8

Mechanics

list			trun .
Mechanics			иникали и политика ISBN 0 7487 3559 3
Author	Title	Publisher	ISBN
Adams, Haighton, Trim	Complete Advanced Level Mathematics: Mechanics: Core Text	Nelson Thornes, 2000	0 7487 3559 3
Bostock & Chandler	Mechanics for A Level	Nelson Thornes, 1996	07487 2596 2
Jefferson & Beadsworth	Introducing Mechanics	Oxford University Press, 2000	0 19 914710 8
Kitchen & Wake	<i>Graded Exercises in</i> <i>Mechanics</i> (Practice questions)	Cambridge University Press, 2001	0 521 64686 3
Nunn & Simmons	Practice for Advanced Mathematics (Practice questions)	Hodder & Stoughton Educational, 1998	0 340 70166 8
Sadler & Thorning	Understanding Mechanics	Oxford University Press, 1996	019 914675 6
SMP	Mathematics for AS and A Level – Mechanics	Cambridge University Press, 1997	0 521 56615 0
Solomon	Advanced Level Mathematics: Mechanics	John Murray, 1995	07195 7082 4
Young	Maths in Perspective 2: Mechanics	Hodder & Stoughton Educational, 1989	07131 78221

Statistics

			4
			Mr. Re
			Pla.
			Can
statistics			18
Author	Title	Publisher	ISBN
Clarke & Cooke	A Basic Course in Statistics	Hodder & Stoughton Educational, 1998	Re Мили, рарасали Врасали ISBN 0 340 71995 8
Crawshaw & Chambers	A Concise Course in Advanced Level Statistics	Nelson Thornes, 2001	0 7487 5475X
Crawshaw & Chambers	A-Level Statistics Study Guide	Nelson Thornes, 1997	0 7487 2997 6
McGill, McLennan, Migliorini	Complete Advanced Level Mathematics: Statistics: Core Text	Nelson Thornes, 2000	07487 3560 7
Norris	<i>Graded Exercises in</i> <i>Statistics</i> (Practice questions)	Cambridge University Press, 2000	0 521 65399 1
Rees	Foundations of Statistics	Chapman & Hall, 1987	0 412 28560 6
Smith	<i>Practice for Advanced Mathematics: Statistics</i> (Practice questions)	Hodder & Stoughton Educational, 1998	0 340 70165X
SMP	Mathematics for AS and A Level – Statistics	Cambridge University Press, 1997	0 521 56616 9
Solomon	Advanced Level Mathematics: Statistics	John Murray, 1996	0 7195 7088 3
Upton & Cook	Introducing Statistics	Oxford University Press, 2001	0 19 914801 5
Upton & Cook	Understanding Statistics	Oxford University Press, 1997	0 19 914391 9

Resources are also listed on Cambridge's public website at **www.cie.org.uk**. Please visit this site on a regular basis as the Resource lists are updated through the year.

Access to teachers' email discussion groups, suggested schemes of work and regularly updated resource lists may be found on the Cambridge Teacher Support website at http://teachers.cie.org.uk. This website is available to teachers at registered Cambridge Centres.

www.papacambridge.com List of formulae and tables of the normal distribution 6.

PURE MATHEMATICS

Algebra

For the quadratic equation $ax^2 + bx + c = 0$:

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

For an arithmetic series:

$$u_n = a + (n-1)d$$
, $S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$

For a geometric series:

$$u_n = ar^{n-1},$$
 $S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1),$ $S_\infty = \frac{a}{1-r} \quad (|r| < 1)$

Binomial expansion:

$$(a+b)^{n} = a^{n} + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^{2} + \binom{n}{3}a^{n-3}b^{3} + \dots + b^{n}, \text{ where } n \text{ is a positive integer}$$

and $\binom{n}{r} = \frac{n!}{r!(n-r)!}$
 $(1+x)^{n} = 1 + nx + \frac{n(n-1)}{2!}x^{2} + \frac{n(n-1)(n-2)}{3!}x^{3} \dots, \text{ where } n \text{ is rational and } |x| < 1$

Arc length of circle = $r\theta$ (θ in radians)

Trigonometry

Area of sector of circle
$$= \frac{1}{2}r^2\theta$$
 (θ in radians)
 $\tan \theta = \frac{\sin \theta}{\cos \theta}$
 $\cos^2 \theta + \sin^2 \theta = 1$, $1 + \tan^2 \theta = \sec^2 \theta$, $\cot^2 \theta + 1 = \csc^2 \theta$
 $\sin(A \pm B) \equiv \sin A \cos B \pm \cos A \sin B$
 $\cos(A \pm B) \equiv \cos A \cos B \mp \sin A \sin B$
 $\tan(A \pm B) \equiv \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$
 $\sin 2A \equiv 2 \sin A \cos A$
 $\cos 2A \equiv \cos^2 A - \sin^2 A \equiv 2 \cos^2 A - 1 \equiv 1 - 2 \sin^2 A$
 $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$
Principal values:
 $-\frac{1}{2}\pi \le \sin^{-1} x \le \frac{1}{2}\pi$
 $0 \le \cos^{-1} x \le \pi$
 $-\frac{1}{2}\pi < \tan^{-1} x < \frac{1}{2}\pi$

List of formulae and how normal distribution

Differentiation

manon		
	f(x)	$f'(x) nx^{n-1}$
	x^n	nx^{n-1}
	$\ln x$	$\frac{1}{x}$
	e ^x	e ^x
	$\sin x$	$\cos x$
	$\cos x$	$-\sin x$
	tan x	$\sec^2 x$
	uv	$u\frac{\mathrm{d}v}{\mathrm{d}x} + v\frac{\mathrm{d}u}{\mathrm{d}x}$
	$\frac{u}{v}$	$\frac{v \frac{\mathrm{d}u}{\mathrm{d}x} - u \frac{\mathrm{d}v}{\mathrm{d}x}}{v \frac{\mathrm{d}v}{\mathrm{d}x}}$
	ν	v^2
If $x = f(t)$ and $y = g(t)$	then $\frac{dy}{dx} = \frac{dy}{dt} \div \frac{dx}{dt}$	

Integration

$$f(x) \qquad \int f(x) dx$$

$$x^{n} \qquad \frac{x^{n+1}}{n+1} + c \quad (n \neq -1)$$

$$\frac{1}{x} \qquad \ln|x| + c$$

$$e^{x} \qquad e^{x} + c$$

$$\sin x \qquad -\cos x + c$$

$$\cos x \qquad \sin x + c$$

$$\sec^{2} x \qquad \tan x + c$$

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx$$

$$\int \frac{f'(x)}{f(x)} dx = \ln|f(x)| + c$$

Vectors

If $\mathbf{a} = a_1\mathbf{i} + a_2\mathbf{j} + a_3\mathbf{k}$ and $\mathbf{b} = b_1\mathbf{i} + b_2\mathbf{j} + b_3\mathbf{k}$ then $\mathbf{a}.\mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3 = |\mathbf{a}||\mathbf{b}|\cos\theta$

Numerical integration

Trapezium rule:

$$\int_{a}^{b} f(x) \, dx \approx \frac{1}{2} h\{y_0 + 2(y_1 + y_2 + \dots + y_{n-1}) + y_n\}, \text{ where } h = \frac{b-a}{n}$$

MECHANICS

Uniformly accelerated motion

v

$$= u + at$$
, $s = \frac{1}{2}(u + v)t$, $s = ut + \frac{1}{2}at^2$, $v^2 = u^2 + 2as$

Motion of a projectile

Equation of trajectory is:

$$y = x \tan \theta - \frac{gx^2}{2V^2 \cos^2 \theta}$$

Elastic strings and springs

$$T = \frac{\lambda x}{l}, \qquad \qquad E = \frac{\lambda x^2}{2l}$$

Motion in a circle

For uniform circular motion, the acceleration is directed towards the centre and has magnitude

$$\omega^2 r$$
 or $\frac{v}{r}$

Centres of mass of uniform bodies

Triangular lamina: $\frac{2}{3}$ along median from vertex Solid hemisphere or radius r: $\frac{3}{8}r$ from centre Hemispherical shell of radius r: $\frac{1}{2}r$ from centre Circular arc of radius r and angle 2α : $\frac{r\sin\alpha}{\alpha}$ from centre Circular sector of radius r and angle 2α : $\frac{2r\sin\alpha}{3\alpha}$ from centre Solid cone or pyramid of height h: $\frac{3}{4}h$ from vertex



PROBABILITY AND STATISTICS

Summary statistics

For ungrouped data:

 $\overline{x} = \frac{\Sigma x}{n},$

standard deviation
$$=\sqrt{\frac{\Sigma(x-\overline{x})^2}{n}} = \sqrt{\frac{\Sigma x^2}{n} - \overline{x}^2}$$

For grouped data:

$$\overline{x} = \frac{\Sigma x f}{\Sigma f}$$
, standard deviation $= \sqrt{\frac{\Sigma (x - \overline{x})^2 f}{\Sigma f}} = \sqrt{\frac{\Sigma x^2 f}{\Sigma f}} - \overline{x}^2$

Discrete random variables

$$E(X) = \Sigma xp$$
$$Var(X) = \Sigma x^2 p - \{E(X)\}^2$$

- / - -

For the binomial distribution B(n, p):

$$p_r = \binom{n}{r} p^r (1-p)^{n-r}, \qquad \mu = np, \qquad \sigma^2 = np(1-p)$$

For the Poisson distribution Po(a):

$$p_r = e^{-a} \frac{a^r}{r!}, \qquad \qquad \mu = a , \qquad \qquad \sigma^2 = a$$

Continuous random variables

$$E(X) = \int x f(x) dx$$
$$Var(X) = \int x^2 f(x) dx - \{E(X)\}^2$$

Sampling and testing

Unbiased estimators:

$$=\frac{\Sigma x}{n}, \qquad \qquad s^2 = \frac{1}{n-1} \left(\Sigma x^2 - \frac{(\Sigma x)^2}{n} \right)$$

Central Limit Theorem:

 \overline{x}

$$\overline{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

Approximate distribution of sample proportion:

$$N\left(p, \frac{p(1-p)}{n}\right)$$

List of formulae and tables of the normal distribution

THE NORMAL DISTRIBUTION FUNCTION

www.papacambridge.com

If Z has a normal distribution with mean 0 and variance 1 then, for each value of z, the table gives the value of $\Phi(z)$, where

 $\Phi(z) = \mathbf{P}(Z \le z) \,.$

For negative values of z use $\Phi(-z) = 1 - \Phi(z)$.

0.80.78810.79100.79390.79670.79950.80230.80510.80780.81060.8133358111416190.90.81590.81860.82120.82380.82640.82890.83150.83400.83650.8389358101315181.00.84130.84380.84610.84850.85080.85310.85540.85770.85990.862125791214161.10.86430.86650.86860.87080.87290.87490.87700.87900.88100.883024681012141.20.88490.88690.88880.89070.89250.89440.89620.89800.89970.90152467911131.30.90320.90490.90660.90820.90990.91150.91310.91470.91620.9177235681011	z	0	1	2	3	4	5	6	7	8	9	1	2	3	4	5 DF	6	7	8	9
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$		0.5000	0.5040	0.5090	0.5120	0.51(0	0.5100	0.5220	0.5270	0.5210	0.5250	4	0	10				20	22	20
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$																				
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$																				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	· ·																			
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$,							-
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$,							
0.7 0.7580 0.7611 0.7642 0.7673 0.7734 0.7744 0.7794 0.7823 0.7823 0.7852 3 6 9 12 15 18 21 0.8 0.7881 0.7910 0.7939 0.7967 0.7995 0.8023 0.8051 0.8078 0.8106 0.8133 3 5 8 11 14 16 19 0.9 0.8159 0.8186 0.8212 0.8238 0.8264 0.8289 0.8315 0.8340 0.8365 0.8389 3 5 8 10 13 15 18 1.0 0.8413 0.8438 0.8461 0.8485 0.8508 0.8571 0.8770 0.8790 0.8810 0.8830 2 4 6 8 10 12 14 1.2 0.8849 0.8869 0.8888 0.8907 0.8925 0.8944 0.8962 0.8980 0.8977 0.9015 2 4 6 7 9 11 13 1.3 0.9020 0.9020 0.9207 0.9222 0.9236												-								
0.8 0.7881 0.7910 0.7939 0.7967 0.7995 0.8023 0.8051 0.8078 0.8106 0.8133 3 5 8 11 14 16 19 0.9 0.8159 0.8186 0.8212 0.8238 0.8264 0.8289 0.8315 0.8340 0.8365 0.8389 3 5 8 10 13 15 18 1.0 0.8413 0.8438 0.8461 0.8485 0.8508 0.8531 0.8554 0.8577 0.8599 0.8621 2 5 7 9 12 14 16 1.1 0.8643 0.8665 0.8686 0.8708 0.8729 0.8749 0.8770 0.8790 0.8810 0.8830 2 4 6 8 10 12 14 1.2 0.8849 0.8869 0.8888 0.8907 0.9252 0.8944 0.8962 0.8907 0.9015 2 4 6 7 9 11 13 14 6 7 9 12 14 15 0.9322 0.9043 <td></td> <td>-</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>												-								
0.9 0.8159 0.8186 0.8212 0.8238 0.8264 0.8289 0.8315 0.8340 0.8365 0.8389 3 5 8 10 13 15 18 1.0 0.8413 0.8438 0.8461 0.8485 0.8508 0.8531 0.8577 0.8599 0.8621 2 5 7 9 12 14 16 1.1 0.8643 0.8665 0.8686 0.8708 0.8729 0.8749 0.8770 0.8790 0.8810 0.8830 2 4 6 7 9 12 14 1.2 0.8849 0.8869 0.8888 0.8907 0.8925 0.8944 0.8962 0.8997 0.9015 2 4 6 7 9 11 13 1.3 0.9022 0.9049 0.9066 0.9022 0.9215 0.9255 0.9279 0.9222 0.9306 0.9177 2 3 4 6 7 8 10 1.5 1.5 0.9322 0.9463 0.9474 0.9484 0.9495 0.9505 0.9515																	-		24	
1.00.84130.84380.84610.84850.85080.85310.85540.85770.85990.862125791214161.10.86430.86650.86860.87080.87290.87490.87700.87900.88100.883024681012141.20.88490.88690.88880.89070.89250.89440.89620.89800.89970.90152467911131.30.90320.90490.90660.90820.90990.91150.91310.91470.91620.91772356810111.40.91920.92070.92220.92360.92510.92650.92790.92920.93060.9319134678101.50.93320.93450.93570.93700.93820.93940.94060.94180.94290.944112456781.60.94520.94630.94740.94840.94950.95050.95150.95250.95350.954512344561.80.96410.96560.96640.96710.96780.96860.96930.96990.9706112234451.90.97720.97780.97830.97880.97930.9786 <td></td> <td>-</td> <td></td> <td>-</td> <td></td> <td></td> <td></td> <td></td> <td></td> <td></td>												-		-						
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$												-								
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$																				
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$														-	-				16	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$																~			15	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$																				14
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$																				
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$												1							10	
$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$												1							8	9
1.90.97130.97190.97260.97320.97380.97440.97500.97560.97610.97671122342.00.97720.97780.97830.97880.97930.97980.98030.98080.98120.981701122332.10.98210.98260.98300.98340.98380.98420.98460.98500.98540.985701122232.20.98610.98640.98680.98710.98750.98780.98810.98840.98870.9890011122222.30.98930.98960.98980.99010.99040.99060.99090.99110.99130.9916011112222.40.99180.99200.99220.99250.99270.99290.99310.99320.99340.9936001111112.50.99380.99400.99410.99430.99450.99460.99490.99510.995200111111												1							7	8
2.00.97720.97780.97830.97880.97930.97980.98030.98080.98120.98170112232.10.98210.98260.98300.98340.98380.98420.98460.98500.98540.985701122232.20.98610.98640.98680.98710.98750.98780.98810.98840.98870.989001112222.30.98930.98960.98980.99010.99040.99060.99090.99110.99130.991601111222.40.99180.99200.99220.99250.99270.99290.99310.99320.99340.993600111112.50.99380.99400.99410.99430.99450.99460.99490.99510.9952001111												1							6	6
2.10.98210.98260.98300.98340.98380.98420.98460.98500.98540.985701122232.20.98610.98640.98680.98710.98750.98780.98810.98840.98870.989001112222.30.98930.98960.98980.99010.99040.99060.99090.99110.99130.991601111222.40.99180.99200.99220.99250.99270.99290.99310.99320.99340.993600111112.50.99380.99400.99410.99430.99450.99460.99480.99490.99510.9952001111												1	-						5	5
2.20.98610.98640.98680.98710.98750.98780.98810.98840.98870.98900111222.30.98930.98960.98980.99010.99040.99060.99090.99110.99130.991601111222.40.99180.99200.99220.99250.99270.99290.99310.99320.99340.993600111112.50.99380.99400.99410.99430.99450.99460.99480.99490.99510.9952001111														-					4	4
2.3 0.9893 0.9896 0.9898 0.9901 0.9904 0.9906 0.9909 0.9911 0.9913 0.9916 0 1 1 1 2 2 2.4 0.9918 0.9920 0.9922 0.9925 0.9927 0.9929 0.9931 0.9932 0.9934 0.9936 0 0 1 <td></td> <td>-</td> <td>-</td> <td>2</td> <td></td> <td></td> <td>-</td> <td>3</td> <td>4</td>													-	-	2			-	3	4
2.4 0.9918 0.9920 0.9922 0.9925 0.9927 0.9929 0.9931 0.9932 0.9934 0.9936 0 0 1 1 1 1 2.5 0.9938 0.9940 0.9941 0.9943 0.9945 0.9946 0.9948 0.9949 0.9951 0.9952 0 0 1 1 1 1 1 1 1												-	-	-	1	2			3	3
2.5 0.9938 0.9940 0.9941 0.9943 0.9945 0.9946 0.9948 0.9949 0.9951 0.9952 0 0 0 1 1 1 1													-	-	l	I		2	2	2
												-	-	-	1	1	-	1	2	2
2.6 0.9953 0.9955 0.9956 0.9957 0.9959 0.9960 0.9961 0.9962 0.9963 0.9964 0 0 0 0 1 1 1													-		1	1	-	1	1	1
																1	Ť.	1	1	1
2.7 0.9965 0.9966 0.9967 0.9968 0.9969 0.9970 0.9971 0.9972 0.9973 0.9974 0 0 0 0 1 1												0	-		0	0	1	1	1	1
2.8 0.9974 0.9975 0.9976 0.9977 0.9978 0.9979 0.9979 0.9980 0.9981 0												0	-		0	0	0	0	1	1
2.9 0.9981 0.9982 0.9982 0.9983 0.9984 0.9984 0.9985 0.9985 0.9986 0.9986 0 0 0 0 0 0 0 0	.9 0	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986	0	0	0	0	0	0	0	0	0

Critical values for the normal distribution

If Z has a normal distribution with mean 0 and variance 1 then, for each value of p, the table gives the value of z such that

 $\mathbf{P}(Z \le z) = p \; .$

р	0.75	0.90	0.95	0.975	0.99	0.995	0.9975	0.999	0.9995
Ζ	0.674	1.282	1.645	1.960	2.326	2.576	2.807	3.090	3.291

Mathematical notation 7.

	thematical notation
	N.D.
	Ag Car
7. Math	ematical notation he syllabus in this booklet may use relevant notation from the following list.
Examinations for t	he syllabus in this booklet may use relevant notation from the following list.
1 Set notation	
e	is an element of
¢	is not an element of
$\{x_1, x_2,\}$	the set with elements x_1, x_2, \dots
$\{x :\}$	the set of all x such that
n(A)	the number of elements in set A
Ø	the empty set
E	the universal set
A'	the complement of the set A
\bowtie	the set of natural numbers, $\{1, 2, 3,\}$
\mathbb{Z}	the set of integers, $\{0, \pm 1, \pm 2, \pm 3,\}$
\mathbb{Z}^+	the set of positive integers, {1, 2, 3,}
\mathbb{Z}_n	the set of integers modulo n , $\{0, 1, 2,, n-1\}$
Q	the set of rational numbers, $\left\{ rac{p}{q}: p \in \mathbb{Z}, q \in \mathbb{Z}^{+} ight\}$
\mathbb{Q}^+	the set of positive rational numbers, $\{x \in \mathbb{Q} : x > 0\}$
$\widetilde{\mathbb{Q}}_0^+$	set of positive rational numbers and zero, $\{x \in \mathbb{Q} : x \ge 0\}$
\mathbb{R}	the set of real numbers
\mathbb{R}^+	the set of positive real numbers, $\{x \in \mathbb{R} : x > 0\}$
\mathbb{R}^+_0	the set of positive real numbers and zero, $\{x \in \mathbb{R} : x \ge 0\}$
C	the set of complex numbers
(x, y)	the ordered pair x, y
A imes B	the cartesian product of sets A and B, i.e. $A \times B = \{(a, b) : a \in A, b \in B\}$
\subseteq	is a subset of
\subset	is a proper subset of
\cup	union
\cap	intersection
[<i>a</i> , <i>b</i>]	the closed interval $\{x \in \mathbb{R} : a \leq x \leq b\}$
[<i>a</i> , <i>b</i>)	the interval $\{x \in \mathbb{R} : a \leq x < b\}$
(a, b]	the interval $\{x \in \mathbb{R} : a < x \le b\}$
(a, b)	the open interval $\{x \in \mathbb{R} : a < x < b\}$
y R x	y is related to x by the relation R
$y \sim x$	y is equivalent to x , in the context of some equivalence relation



2 Miscellaneous symbols

=	is equal to
\neq	is not equal to
≡	is identical to or is congruent to
\approx	is approximately equal to
≅	is isomorphic to
oc	is proportional to
<	is less than
≤ >	is less than or equal to, is not greater than
>	is greater than
\geq	is greater than or equal to, is not less than
00	infinity
$p \wedge q$	p and q
$p \lor q$	p or q (or both)
~ <i>p</i>	not p
$p \Rightarrow q$	p implies q (if p then q)
$p \Leftarrow q$	p is implied by q (if q then p)
$p \Leftrightarrow q$	p implies and is implied by q (p is equivalent to q)
Ξ	there exists
\forall	for all

3 Operations

a + b	a plus b
a-b	a minus b
$a \times b$, ab , $a.b$	a multiplied by b
$a \div b, \frac{a}{b}, a / b$	a divided by b
$\sum_{i=1}^{n} a_i$	$a_1 + a_2 + \ldots + a_n$
$\prod_{i=1}^n a_i$	$a_1 \times a_2 \times \ldots \times a_n$
\sqrt{a}	the positive square root of a
a	the modulus of a
n!	n factorial
$\binom{n}{r}$	the binomial coefficient $rac{n!}{r! \ (n-r)!}$ for $n \in \mathbb{Z}^+$
	or $\frac{n(n-1)(n-r+1)}{r!}$ for $n \in \mathbb{Q}$

4 Functions

$\mathbf{f}(x)$	the value of the function f at x
$f: A \to B$	f is a function under which each element of set A has an image in set B
$f: x \to y$	the function f maps the element x to the element y
f^{-1}	the inverse function of the function f
gf	the composite function of f and g which is defined by $gf(x) = g(f(x))$
$\lim_{x \to a} \mathbf{f}(x)$	the limit of $f(x)$ as x tends to a

	 an increment of x the derivative of y with respect to x the <i>n</i>th derivative of y with respect to x
$\Delta x, \delta x$	an increment of x
$\frac{\mathrm{d}y}{\mathrm{d}x}$	the derivative of y with respect to x
$\frac{\mathrm{d}^n y}{\mathrm{d} x^n}$	the <i>n</i> th derivative of <i>y</i> with respect to <i>x</i>
$f'(x), f''(x),, f^{(n)}(x)$	the first, second,, n th derivatives of $f(x)$ with respect to x
$\int y \mathrm{d}x$	the indefinite integral of y with respect to x
$\int_{a}^{b} y \mathrm{d}x$	the definite integral of y with respect to x between the limits $x = a$ and $x = b$
$\frac{\partial V}{\partial x}$	the partial derivative of V with respect to x
$\dot{x}, \ddot{x},$	the first, second, derivatives of x with respect to t

5 Exponential and logarithmic functions

e	base of natural logarithms
e^x , exp x	exponential function of x
$\log_a x$	logarithm to the base a of x
$\ln x$, $\log_e x$	natural logarithm of x
$\lg x, \log_{10} x$	logarithm of x to base 10

6 Circular and hyperbolic functions

sin, cos, tan, cosec, sec, cot	the circular functions
$\sin^{-1}, \cos^{-1}, \tan^{-1}, \ \cos^{-1}, \sec^{-1}, \cot^{-1}$	the inverse circular functions
sinh, cosh, tanh, cosech, sech, coth	the hyperbolic functions

 $\left. \begin{array}{c} \sinh^{-1}, \ \cosh^{-1}, \ \tanh^{-1}, \\ \cosh^{-1}, \ \operatorname{sech}^{-1}, \ \operatorname{coth}^{-1} \end{array} \right\} \qquad \text{the inver}$

the inverse hyperbolic functions

7 Complex numbers

i	square root of -1
Z	a complex number, $z = x + i y = r(\cos \theta + i \sin \theta)$
Re z	the real part of z, $\operatorname{Re} z = x$
Im z	the imaginary part of z , Im $z = y$
z	the modulus of z, $ z = \sqrt{x^2 + y^2}$
arg z	the argument of z, arg $z = \theta, -\pi < \theta \le \pi$
Ζ*	the complex conjugate of z , $x - i y$

8 Matrices

Μ	a matrix M
\mathbf{M}^{-1}	the inverse of the matrix ${f M}$
\mathbf{M}^{T}	the transpose of the matrix ${f M}$
det M or M	the determinant of the square matrix ${\bf M}$

atical notation	MMM. Dahac
9 Vectors	the vector a the vector represented in magnitude and direction by the directed line
a	the vector a
\overrightarrow{AB}	the vector represented in magnitude and direction by the directed line segment <i>AB</i>
â	a unit vector in the direction of a
i, j, k	unit vectors in the directions of the cartesian coordinate axes
$ \mathbf{a} , a$	the magnitude of ${f a}$
$ \overrightarrow{AB} , AB$	the magnitude of \overrightarrow{AB}
a.b	the scalar product of a and b
$\mathbf{a} \times \mathbf{b}$	the vector product of ${f a}$ and ${f b}$

10 Probability and statistics

<i>A</i> , <i>B</i> , <i>C</i> , etc.	events
$A \cup B$	union of the events A and B
$A \cap B$	intersection of the events A and B
P(A)	probability of the event A
A'	complement of the event A
P(A B)	probability of the event A conditional on the event B
<i>X</i> , <i>Y</i> , <i>R</i> , etc.	random variables
<i>x</i> , <i>y</i> , <i>r</i> , etc.	values of the random variables X, Y, R, etc.
<i>x</i> ₁ , <i>x</i> ₂ ,	observations
$f_1, f_2,$	frequencies with which the observations x_1, x_2, \dots occur
p (<i>x</i>)	probability function $P(X = x)$ of the discrete random variable X
$p_1, p_2,$	probabilities of the values x_1, x_2, \dots of the discrete random variable X
f(x), g(x),	the value of the probability density function of a continuous random variable X
F(x), G(x),	the value of the (cumulative) distribution function $P(X \le x)$ of a continuous random variable X
E(X)	expectation of the random variable X
E(g(X))	expectation of $g(X)$
Var(X)	variance of the random variable X
$\mathbf{G}(t)$	probability generating function for a random variable which takes the values $0, 1, 2,$
B(<i>n</i> , <i>p</i>)	binomial distribution with parameters n and p
$Po(\mu)$	Poisson distribution, mean μ
$N(\mu, \sigma^2)$	normal distribution with mean μ and variance σ^2
μ	population mean
σ^2	population variance
σ	population standard deviation
\overline{x} , m	sample mean
$s^2, \hat{\sigma}^2$	unbiased estimate of population variance from a sample,
	$s^2 = \frac{1}{n-1} \sum (x_i - \overline{x})^2$
φ	probability density function of the standardised normal variable with distribution $N(0, 1)$
Φ	corresponding cumulative distribution function
ρ	product moment correlation coefficient for a population
r	product moment correlation coefficient for a sample
$\operatorname{Cov}(X, Y)$	covariance of X and Y

Additional information 8.

8.1 Guided learning hours

www.papaCambridge.com Cambridge International A Level syllabuses are designed on the assumption that candidates have about 360 guided learning hours per subject over the duration of the course. Cambridge International AS Level syllabuses are designed on the assumption that candidates have about 180 guided learning hours per subject over the duration of the course. ('Guided learning hours' include direct teaching and any other supervised or directed study time. They do not include private study by the candidate.)

However, these figures are for guidance only, and the number of hours required may vary according to local curricular practice and the candidates' prior experience of the subject.

8.2 Recommended prior learning

We recommend that candidates who are beginning this course should have previously completed a Cambridge O Level or Cambridge IGCSE course in Mathematics or the equivalent.

8.3 Progression

Cambridge International A Level Mathematics provides a suitable foundation for the study of Mathematics or related courses in higher education.

Cambridge International AS Level Mathematics constitutes the first half of the Cambridge International A Level course in Mathematics and therefore provides a suitable foundation for the study of Mathematics at A Level and thence for related courses in higher education.

Component codes 8.4

Because of local variations, in some cases component codes will be different in instructions about making entries for examinations and timetables from those printed in this syllabus, but the component names will be unchanged to make identification straightforward.

8.5 Grading and reporting

www.papacambridge.com Cambridge International A Level results are shown by one of the grades A*, A, B, C, D or E indicating the standard achieved, Grade A* being the highest and Grade E the lowest. 'Ungraded' indicates that the candidate has failed to reach the standard required for a pass at either Cambridge International AS Level or A Level. 'Ungraded' will be reported on the statement of results but not on the certificate.

If a candidate takes a Cambridge International A Level and fails to achieve grade E or higher, a Cambridge International AS Level grade will be awarded if both of the following apply:

- the components taken for the Cambridge International A Level by the candidate in that series included all the components making up a Cambridge International AS Level
- the candidate's performance on these components was sufficient to merit the award of a Cambridge International AS Level grade.

For languages other than English, Cambridge also reports separate speaking endorsement grades (Distinction, Merit and Pass), for candidates who satisfy the conditions stated in the syllabus.

Percentage uniform marks are also provided on each candidate's statement of results to supplement their grade for a syllabus. They are determined in this way:

- A candidate who obtains...
 - ... the minimum mark necessary for a Grade A* obtains a percentage uniform mark of 90%.
 - ... the minimum mark necessary for a Grade A obtains a percentage uniform mark of 80%.
 - ... the minimum mark necessary for a Grade B obtains a percentage uniform mark of 70%.
 - ... the minimum mark necessary for a Grade C obtains a percentage uniform mark of 60%.
 - ... the minimum mark necessary for a Grade D obtains a percentage uniform mark of 50%.
 - ... the minimum mark necessary for a Grade E obtains a percentage uniform mark of 40%.
 - ... no marks receives a percentage uniform mark of 0%.

Candidates whose mark is none of the above receive a percentage mark in between those stated according to the position of their mark in relation to the grade 'thresholds' (i.e. the minimum mark for obtaining a grade). For example, a candidate whose mark is halfway between the minimum for a Grade C and the minimum for a Grade D (and whose grade is therefore D) receives a percentage uniform mark of 55%.

The percentage uniform mark is stated at syllabus level only. It is not the same as the 'raw' mark obtained by the candidate, since it depends on the position of the grade thresholds (which may vary from one series to another and from one subject to another) and it has been turned into a percentage.

Cambridge International AS Level results are shown by one of the grades a, b, c, d or e indicating the standard achieved, Grade a being the highest and Grade e the lowest. 'Ungraded' indicates that the candidate has failed to reach the standard required for a pass at Cambridge International AS Level. 'Ungraded' will be reported on the statement of results but not on the certificate.

For languages other than English, Cambridge will also report separate speaking endorsement grades (Distinction, Merit and Pass) for candidates who satisfy the conditions stated in the syllabus.

The content and difficulty of a Cambridge International AS Level examination is equivalent to the first half of a corresponding Cambridge International A Level.

www.papaCambridge.com Percentage uniform marks are also provided on each candidate's statement of results to supplement grade for a syllabus. They are determined in this way:

- A candidate who obtains...
 - ... the minimum mark necessary for a Grade a obtains a percentage uniform mark of 80%.
 - ... the minimum mark necessary for a Grade b obtains a percentage uniform mark of 70%.
 - ... the minimum mark necessary for a Grade c obtains a percentage uniform mark of 60%.
 - ... the minimum mark necessary for a Grade d obtains a percentage uniform mark of 50%.
 - ... the minimum mark necessary for a Grade e obtains a percentage uniform mark of 40%.
 - ... no marks receives a percentage uniform mark of 0%.

Candidates whose mark is none of the above receive a percentage mark in between those stated according to the position of their mark in relation to the grade 'thresholds' (i.e. the minimum mark for obtaining a grade). For example, a candidate whose mark is halfway between the minimum for a Grade c and the minimum for a Grade d (and whose grade is therefore d) receives a percentage uniform mark of 55%.

The percentage uniform mark is stated at syllabus level only. It is not the same as the 'raw' mark obtained by the candidate, since it depends on the position of the grade thresholds (which may vary from one series to another and from one subject to another) and it has been turned into a percentage.

8.6 Access

Reasonable adjustments are made for disabled candidates in order to enable them to access the assessments and to demonstrate what they know and what they can do. For this reason, very few candidates will have a complete barrier to the assessment. Information on reasonable adjustments is found in the Cambridge Handbook which can be downloaded from the website www.cie.org.uk

Candidates who are unable to access part of the assessment, even after exploring all possibilities through reasonable adjustments, may still be able to receive an award based on the parts of the assessment they have taken

8.7 Resources

Copies of syllabuses, the most recent question papers and Principal Examiners' reports for teachers are on the Syllabus and Support Materials CD-ROM, which we send to all Cambridge International Schools. They are also on our public website – go to **www.cie.org.uk/alevel**. Click the Subjects tab and choose your subject. For resources, click 'Resource List'.

You can use the 'Filter by' list to show all resources or only resources categorised as 'Endorsed by Cambridge'. Endorsed resources are written to align closely with the syllabus they support. They have been through a detailed quality-assurance process. As new resources are published, we review them against the syllabus and publish their details on the relevant resource list section of the website.

Additional syllabus-specific support is available from our secure Teacher Support website http://teachers.cie.org.uk which is available to teachers at registered Cambridge schools. It provides past question papers and examiner reports on previous examinations, as well as any extra resources such as schemes of work or examples of candidate responses. You can also find a range of subject communities on the Teacher Support website, where Cambridge teachers can share their own materials and join discussion groups.

University of Cambridge International Examinations 1 Hills Road, Cambridge, CB1 2EU, United Kingdom Tel: +44 (0)1223 553554 Fax: +44 (0)1223 553558 Email: international@cie.org.uk www.cie.org.uk

© University of Cambridge International Examinations 2011

www.papacambridge.com