

**MARK SCHEME for the May/June 2010 question paper  
for the guidance of teachers**

**9231 FURTHER MATHEMATICS**

**9231/13**

Paper 13, maximum raw mark 100

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

### Mark Scheme Notes

Marks are of the following three types:

**M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

**A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

**B** Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol  $\surd$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking  $g$  equal to 9.8 or 9.81 instead of 10.

Page 3	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

The following abbreviations may be used in a mark scheme or used on the scripts:

- AEF Any Equivalent Form (of answer is equally acceptable)
- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- CWO Correct Working Only – often written by a 'fortuitous' answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

### **Penalties**

- MR –1 A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through  $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA –1 This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

www.PapaCambridge.com

1 Relevant working from  

$$\mathbf{A} - 5\mathbf{I} = \begin{pmatrix} 0 & -3 & 0 \\ 1 & -3 & 1 \\ -1 & 3 & -1 \end{pmatrix}$$
 or the equivalent in equations  
 to obtain an eigenvector of the form  $\begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}$  A1

[2]

An eigenvalue of  $\mathbf{A} + \mathbf{A}^2$  is  $5 + 25 = 30$  B1  
 Corresponding eigenvector, as above B1  
[2]

No penalty for not hence methods.

Accept  $\lambda = 6$  with  $\begin{pmatrix} -1 \\ -1 \\ 1 \end{pmatrix}$

and  $\lambda = 20$  with  $\begin{pmatrix} 3 \\ 1 \\ -1 \end{pmatrix}$

Accept linear scaling of eigenvectors.

2  $2 \sin \alpha \sum_{n=1}^N \sin(2n\alpha) = \cos \alpha - \cos[(2N + 1)\alpha]$  M1A1

$\Rightarrow$  displayed result (AG) M1A1  
[4]

$\cos(2N + 1)\pi/3$  oscillates finitely as  $n \rightarrow \infty \Rightarrow \sum_{n=1}^{\infty} \sin(2n\pi/3)$  does not converge (CWO) B1

Require  $\alpha = \frac{\pi}{3}$ , 'oscillate' or values of  $\cos(2N + 1)\frac{\pi}{3}$  given as  $\frac{1}{2}$  or  $-1$  [1]

3  $H_k : x_k > 2$  for some  $k$  B1

$x_{k+1} - 2 = (2x_k^2 - 8)/(2x_k + 3)$  M1A1

$H_k \Rightarrow 2x_k^2 - 8 > 0 \Rightarrow x_{k+1} > 2 \Rightarrow H_{k+1}$  A1

$x_1 = 3 > 2 \Rightarrow H_1$  is true B1 CWO

Completion of the induction argument A1  
[6]

<b>Page 5</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>
	<b>GCE A LEVEL – May/June 2010</b>	<b>9231</b>

Alternatively for lines 2 and 3:

$$x_{k+1} = x_k + \frac{1}{2} - \frac{3\frac{1}{2}}{(2x_k + 3)}$$

$$H_k \Rightarrow 2x_k + 3 > 7 \Rightarrow H_{k+1}$$

**OR**  $x_{k+1} = x_k + \frac{x_{k-2}}{(2x_k + 3)}$

$$x_k > 2 \Rightarrow x_{k+1} > 2$$

**OR**  $x_{k+1} - x_k = \frac{x_{k-2}}{(2x_k + 3)}$

$$x_k > 2 \Rightarrow x_{k+1} > x_k > 2$$

Minimum conclusion is 'Hence true for  $n \geq 1$ '.

**4**  $dx/dt = t \cos t, dy/dt = t \sin t$  (both)

$$\sqrt{(dx/dt)^2 + (dy/dt)^2} = t$$

$$S = 2\pi \int_0^{\pi/2} (t \sin t - t^2 \cos t) dt \text{ (AEF)}$$

$$\int t \sin t dt = -t \cos t + \sin t$$

$$\int t^2 \cos t dt = t^2 \sin t + 2t \cos t - 2 \sin t$$

$$S = (\pi/2)[12 - \pi^2]$$

Accept forms such as  $6\pi - \pi^3/2$ , etc.

**OR** for lines 4 and 5

$$2\pi \left[ (-t \cos t) + \int \cos t dt - \left( t^2 \sin t - \int 2t \sin t dt \right) \right]_0^{\pi/2} \text{ (LNR)}$$

$$= 2\pi \left[ -t \cos t + \sin t - t^2 \sin t - 2t \cos t + 2 \sin t \right]_0^{\pi/2} \text{ (LNR)}$$

Page 6	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

- 5  $(c + is)^5 = c^5 + 5c^4(is) + 10c^3(-s^2) + 10c^2(is)^3 + 5c(is)^4 + (is)^5$
- $\theta = 5c^4s - 10c^2s^3 + s^5$
- $= 5s(1-s^2)^2 - 10s^3(1-s^2) + s^5$  M1
- $= 16s^5 - 20s^3 + 5s$  (AG) A1 OEW [4]
- $x = \sin \theta \Rightarrow \sin 5\theta = -1/2$  M1
- Roots are  $\sin q\pi$  where  $q = 7/30, 11/30, 31/30, 35/30, 43/30$  A3
- A1 for 1 root: + A1 for 2 further roots: + A1 for completion CWO CWO [4]

Alternative answers

$q = \frac{11}{30}, \frac{23}{30}, \frac{35}{30}, \frac{47}{30}, \frac{59}{30}$

or

$q = \frac{7}{30}, \frac{19}{30}, \frac{31}{30}, \frac{43}{30}, \frac{55}{30}$

- 6 (i) One asymptote is  $x = -1$  B1
- $y = x - 4 - 3/(x + 1)$  M1
- Require  $y = x +$  non-zero constant.
- Other asymptote is  $y = x - 4$  A1 [3]
- Alternatively for last two marks:
- OR  $x + k \approx (x^2 - 3x - 7)/(x + 1)$  for large  $x \Rightarrow x^2 + (k + 1)x + k \approx x^2 - 3x - 7$  for large  $x$  M1
- $\Rightarrow k + 1 = -3 \Rightarrow k = -4 \Rightarrow$  other asymptote is  $y = x - 4$  A1
- OR  $(x^2 - 3x - 7)/(x + 1) = mx + c \Rightarrow (m - 1)x^2 + (m + c + 3)x + c + 7 = 0$  M1
- Put  $m - 1 = 0, m + c + 3 = 0$  to obtain other asymptote is  $y = x - 4$  A1
- (ii) Obtains any correct result for  $dy/dx$  M1
- $\Rightarrow \dots \Rightarrow dy/dx = [(x + 1)^2 + 3]/(x + 1)^2 = 1 + 3/(x + 1)^2$  A1
- No comment required. [2]
- (iii) Axes and asymptotes correctly placed B1
- Right-hand branch B1
- Left-hand branch B1
- Deduct 1 overall for bad form(s) at infinity [3]

Page 7	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

- 7 (i)  $dx/dt = (2t - 2t^3)e^{-t^2}$ ,  $dy/dt = (1 - 2t^2)e^{-t^2}$  (both, AEF)
- Obtains displayed result (AG) M1 [3]
- (ii) Any correct result for  $d(dy/dx)/dt$  in terms of  $t$  (\*) M1A1A1  
M1 – Quotient Rule A1A1 for terms in numerator  
 $d^2y/dx^2 = (*) \times dt/dx$  expressed in terms of  $t$  M1
- Simplify to, e.g.,  $\frac{(-1 + t^2 - 2t^4)e^{t^2}}{4t^3(1 - t^2)^3}$  A1
- Two terms in numerator must be combined. [5]
- 8 Complementary function is  $Ae^{-x} + Be^{-4x}$  M1A1
- Particular integral of form  $P \sin 3x + Q \cos 3x$ , so that
- $-5P + 15Q = 10$  and  $15P - 5Q = -20$  M1
- $\Rightarrow P = -7/5, Q = -1/5$  A1
- General solution is  $y = Ae^{-x} + Be^{-4x} - 1.4 \sin 3x - 0.2 \cos 3x$  A1  
[5]
- $Ae^{-x} + Be^{-4x} \rightarrow 0$  as  $x \rightarrow +\infty$ , M1
- $\tan \phi = \frac{1}{7}$  M1
- $R = \sqrt{2} = 1.41, \phi = \pi + \arctan(1/7) = 3.28$  B1A1
- Accept  $R = \sqrt{2}$ , but must be positive. [4]

<b>Page 8</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>
	<b>GCE A LEVEL – May/June 2010</b>	<b>9231</b>

www.PapaCambridge.com

9  $D[s^{n+1}c] = -s^{n+2} + (n+1)s^n c^2$ , where  $D \equiv d/dx$ ,  $s = \sin x$ ,  $c = \cos x$

$$= \dots = -(n+2)s^{n+2} + (n+1)s^n$$

Integrates w.r.t.  $x$  over the range  $[0, \pi/2]$  to obtain  $0 = -(n+2)I_{n+2} + (n+1)I_n$  M1

Result (AG) A1  
[4]

**OR**  $I_n = -[cs^{n-1}]_0^{\pi/2} + (n-1) \int_0^{\pi/2} c^2 s^{n-2} dx$  M1

$$\Rightarrow (n-1)I_{n-2} - (n-1)I_n \quad (n \geq 2) \quad \text{M1A1}$$

$$\Rightarrow I_n = [(n-1)/n]I_{n-2} \Rightarrow I_{n+2} = [(n+1)/(n+2)]I_n \text{ for } n \geq 0 \quad \text{A1}$$

**OR** Starts with  $I_{n+2}$  and relates to  $I_n$  directly: mark as above

**OR**  $I_n = \int_0^{\pi/2} \sin^{n+2} \theta \operatorname{cosec}^2 \theta d\theta$

$$= [-\sin^{n+2} \theta \cot \theta]_0^{\pi/2} + \int_0^{\pi/2} (n+2) \sin^{n+1} \theta \cos \theta \cot \theta d\theta \quad (\text{LNR}) \quad \text{M1}$$

$$= 0 + (n+2) \int_0^{\pi/2} \sin^n \theta (1 - \sin^2 \theta) d\theta \quad (\text{LR}) \quad \text{M1}$$

$$= (n+2)I_n - (n+2)I_{n+2} \quad \text{A1}$$

$$\Rightarrow I_{n+2} = \frac{(n+1)}{(n+2)} I_n \quad \text{A1}$$



Page 9	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

$$\bar{y} = \frac{\frac{1}{2} \int_0^{\pi/2m} \sin^8 mx \, dx}{\int_0^{\pi/2m} \sin^4 mx \, dx}$$

let  $u = mx$

$$\bar{y} = \frac{\frac{1}{2} \int_0^{\pi/2} \sin^8 u \, du}{\int_0^{\pi/2} \sin^4 u \, du}$$

$$I_0 = \frac{\pi}{2} \quad I_2 = \frac{1}{2} \times \frac{\pi}{4} \quad I_4 = \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2} \left( = \frac{3\pi}{16} \right)$$

$$I_6 = \frac{5}{6} \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2} \quad I_8 = \frac{7}{8} \times \frac{5}{6} \times \frac{3}{4} \times \frac{1}{2} \times \frac{\pi}{2} \left( = \frac{105\pi}{768} \right)$$

$$\bar{y} = \frac{1}{2} \frac{I_8}{I_4} = \frac{35}{96} \quad (\text{or } 0.365)$$

M1

A1

B1

B1

B1

[6]

**OR** for last 3 marks

$$\frac{I_8}{I_4} = \frac{35}{48} \quad (\text{oe})$$

$$\therefore \bar{y} = \frac{35}{96}$$

M1A1

A1

**10 (i)**  $x = 1/y \Rightarrow 2y^4 - 4y^3 - cy^2 - y - 1 = 0$

M1A1

[2]

**(ii)**  $\sum \alpha^2 = 1 - 2c$

M1A1

$$\sum \alpha^{-2} = 4 + c$$

A1

(M1 is for use of  $\sum \alpha^2 = (\sum \alpha)^2 - 2 \sum \alpha\beta$  in either part.)

[3]

**(iii)**  $S = \sum (\alpha - \alpha^{-1})^2 = \sum \alpha^2 + \sum \alpha^{-2} - 8 = -c - 3$

M1A1✓

A1ft is for adding answers to **(ii)** correctly and subtracting 8.

[2]

**(iv)**  $c = -3 \Rightarrow S = 0$  so that if all roots are real then  $\alpha = \pm 1$

and similarly for  $\beta, \gamma, \delta$

M1A1 CWO

This is impossible since e.g.,  $\alpha\beta\gamma\delta = -2$ , or any other contradiction

A1 CWO

[3]

Page 10	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

www.PapaCambridge.com

11 (i)  $dr/d\theta = -a/(1+\theta)^2$

Since  $a > 0$ , then  $dr/d\theta < 0$ ,  $\forall$  points of  $C$

[4]

OR  $a(> 0)$  constant and as  $\theta(> 0)$  increases,  $1 + \theta$  increases

M1

$\therefore \frac{a}{1+\theta}$  decreases

A1

(ii)  $y = a \sin \theta / (1 + \theta)$

M1

$dy/d\theta = 0 \Rightarrow \dots \Rightarrow (1 + \theta) \cos \theta - \sin \theta = 0$

A1

$\Rightarrow \tan \theta = 1 + \theta$  (AG)

A1

$\tan \theta - 1 - \theta = -0.135$  when  $\theta = 1.1$ ;  $\tan \theta - 1 - \theta = +0.372$  when  $\theta = 1.2$

B1

[4]

OR equivalent argument for B1 such as:

$\tan(1.1) \approx 1.96 < 2.1$ ,  $\tan(1.2) \approx 2.37 > 2.2$

B1

or  $f(\theta) = \tan \theta - (1 + \theta) \Rightarrow f(1.1) = -0.14$   $f(1.2) = 0.37$

(iii) Sketch:

Approximately correct shape and placement for  $0 \leq \theta \leq \pi/2$  and passing through  $(a, 0)$  and  $(0.4a, \pi/2)$ , approximately, indicated in some way

B1B1

Maximum in interval  $(\pi/4, \pi/2)$

B1

[3]

(iv)  $A = (a^2/2) \int_0^{\pi/2} (1+\theta)^{-2} d\theta$

M1

$= (a^2/2) \left[ -(1+\theta)^{-1} \right]_0^{\pi/2}$

A1

$= \dots = \pi a^2 / 2 (\pi + 2)$

A1

Do not accept double minus signs or fractions in the denominator for the final mark.

[3]

Page 11	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

## 12 EITHER

(i)  $(\mathbf{i} + \mathbf{j}) \times (\mathbf{j} + 2\mathbf{k}) = 2\mathbf{i} - 2\mathbf{j} + \mathbf{k}$

$$PQ = (4\mathbf{i} + \mathbf{j} + 3\mathbf{k}) \cdot (2\mathbf{i} - 2\mathbf{j} + \mathbf{k}) / 3 = 9/3 = 3$$

M1A1  
[4]

(ii)  $(4 + \lambda)/2 = (1 + \lambda - \mu)/(-2) = (3 - 2\mu)/1$  (AEF)

**OR**  $(4 + \lambda) + (1 + \lambda - \mu) = 0$ ,  $(1 + \lambda - \mu + 2(3 - 2\mu)) = 0$ , both

M1A1

$$\Rightarrow \dots \Rightarrow \mu = 1$$

M1A1

Position vector of Q is  $-\mathbf{i} + \mathbf{j} + \mathbf{k}$

A1  
[5]

Parts (i) and (ii) together:

$$\vec{PQ} = \begin{pmatrix} 4 + \lambda \\ 1 + \lambda - \mu \\ 3 - 2\mu \end{pmatrix}$$

B1

$$\vec{PQ} \cdot \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix} = 0 \Rightarrow 5 + 2\lambda - \mu = 0$$

M1A1

$$\vec{PQ} \cdot \begin{pmatrix} 0 \\ 1 \\ 2 \end{pmatrix} = 0 \Rightarrow 7 + \lambda - 5\mu = 0$$

A1

$$\lambda = -2, \mu = 1$$

M1A1

$$\vec{PQ} \cdot \begin{pmatrix} 2 \\ -2 \\ 1 \end{pmatrix} \Rightarrow |\vec{PQ}| = 3$$

M1A1

$$\vec{OQ} = \begin{pmatrix} -1 \\ 1 \\ 1 \end{pmatrix}$$

A1

[9]

Page 12	Mark Scheme: Teachers' version	Syllabus
	GCE A LEVEL – May/June 2010	9231

(iii)  $(\mathbf{i} + \mathbf{j}) \times (4\mathbf{i} + \mathbf{j} + 3\mathbf{k}) = 3\mathbf{i} - 3\mathbf{j} - 3\mathbf{k}$

$$p_2 = \left| \frac{[(\mathbf{i} - \mathbf{j} - \mathbf{k})(4\mathbf{i} + \mathbf{k})] \cdot \sqrt{3}}{\sqrt{3}} \right|$$

$$= \dots = \sqrt{3}$$

A1

[4]

for final 2 marks

$$\pi : x - y - z = 0$$

$$\text{perpendicular distance} = \left| \frac{-1-1-1}{\sqrt{1^2+1^2+1^2}} \right| = \sqrt{3}$$

or any other method.

M1 is for a complete strategy and A1 for  $\sqrt{3}$ .

12 OR

(i) (a) 
$$\begin{pmatrix} 1 & 1 & 5 & 7 \\ 3 & 9 & 17 & 25 \\ 1 & 7 & 7 & 11 \\ 3 & 6 & 16 & 23 \end{pmatrix} \rightarrow \dots \rightarrow \begin{pmatrix} 1 & 1 & 5 & 7 \\ 0 & 6 & 2 & 4 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

OR Establishes equivalent result for  $\mathbf{M}^T$

OR Establishes 2 linearly independent relations for rows or columns

M1A1

$$\Rightarrow r(\mathbf{M}) = 2 \Rightarrow \dim(R) = 2$$

A1

(b) Basis for  $R_T$  is  $\left\{ \begin{pmatrix} 1 \\ 3 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 9 \\ 7 \\ 6 \end{pmatrix} \right\}$

M1A1

[5]

(ii) Let 
$$\begin{pmatrix} 1 \\ -15 \\ -17 \\ -6 \end{pmatrix} = p \begin{pmatrix} 1 \\ 3 \\ 1 \\ 3 \end{pmatrix} + q \begin{pmatrix} 1 \\ 9 \\ 7 \\ 6 \end{pmatrix}$$

Solves any 2 of:

$$p + q = 1, 3p + 9q = -15, p + 7q = -17, 3p + 6q = -6 \text{ to obtain } p = 4, q = -3$$

M1A1

Checks consistency with other 2 equations

A1

[3]

(iii)  $\mathbf{M} \begin{pmatrix} 4 \\ -3 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ -15 \\ -17 \\ -6 \end{pmatrix}$  OR Works with equations ( $\mathbf{M}$  written out fully.) (AG) BI

$\mathbf{e}_1$  and  $\mathbf{e}_2$  basis vectors for  $\mathbf{T} \Rightarrow \mathbf{M}\mathbf{e}_1 = 0, \mathbf{M}\mathbf{e}_2 = 0$  M1

$$\mathbf{M} \begin{pmatrix} 4 \\ -3 \\ 0 \\ 0 \end{pmatrix} + \lambda \mathbf{M}\mathbf{e}_1 + \mu \mathbf{M}\mathbf{e}_2 = \begin{pmatrix} 1 \\ -15 \\ -17 \\ -6 \end{pmatrix} + \lambda 0 + \mu 0 = \begin{pmatrix} 1 \\ -15 \\ -17 \\ -6 \end{pmatrix}, \forall \lambda, \mu$$
A1

[3]

(iv) Need  $-3 + \lambda + 2\mu = 0$  M1

$$\lambda = 1 \Rightarrow \mu = 1 \Rightarrow \mathbf{x} = \begin{pmatrix} 37 \\ 0 \\ -3 \\ -3 \end{pmatrix}$$
A1

[2]

Accept a parametric answer

e.g.  $\begin{pmatrix} 46 \\ 0 \\ -9 \\ 0 \end{pmatrix} + t \begin{pmatrix} -3 \\ 0 \\ 2 \\ -1 \end{pmatrix}$

Augmented Matrix Method

(i) (a)  $\left( \begin{array}{cccc|c} 1 & 1 & 5 & 7 & 1 \\ 3 & 9 & 17 & 25 & -15 \\ 1 & 7 & 7 & 11 & -17 \\ 3 & 6 & 16 & 23 & -6 \end{array} \right) \sim \left( \begin{array}{cccc|c} 1 & 1 & 5 & 7 & 1 \\ 0 & 6 & 2 & 4 & -18 \\ 0 & 6 & 2 & 4 & -18 \\ 0 & 3 & 1 & 2 & -9 \end{array} \right) \sim \left( \begin{array}{cccc|c} 1 & 1 & 5 & 7 & 1 \\ 0 & 3 & 1 & 2 & -9 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{array} \right)$  M1A1

$\Rightarrow r(\mathbf{M}) = \dim(R) = 2$  A1

(b) Basis for  $R$  is  $\left\{ \begin{pmatrix} 1 \\ 3 \\ 1 \\ 3 \end{pmatrix}, \begin{pmatrix} 1 \\ 9 \\ 7 \\ 6 \end{pmatrix} \right\}$  M1A1

[5]

(ii) and (iii)

$$x + y + 5z + 7t = 1$$

$$3y + z + 2t = -9$$

$$\Rightarrow \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 46 \\ 0 \\ -9 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} 14 \\ 1 \\ -3 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ 0 \\ -2 \\ 1 \end{pmatrix}$$

A1

$$\lambda = -3, \mu = 0 \Rightarrow \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 4 \\ -3 \\ 0 \\ 0 \end{pmatrix}$$

B1

$$\therefore \mathbf{M} \begin{pmatrix} 4 \\ -3 \\ 0 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ -15 \\ -17 \\ -6 \end{pmatrix}$$

$$\mathbf{M}\mathbf{e}_1 = \mathbf{M}\mathbf{e}_2 = 0$$

M1

$$\therefore \mathbf{M} \left( \begin{pmatrix} 4 \\ -3 \\ 0 \\ 0 \end{pmatrix} + \lambda \mathbf{e}_1 + \mu \mathbf{e}_2 \right) = \begin{pmatrix} 1 \\ -15 \\ -17 \\ -6 \end{pmatrix}$$

A1

$$\text{(iv) } \lambda = 0 \quad \begin{pmatrix} x \\ y \\ z \\ t \end{pmatrix} = \begin{pmatrix} 46 \\ 0 \\ -9 \\ 0 \end{pmatrix} + \mu \begin{pmatrix} 3 \\ 0 \\ -2 \\ 1 \end{pmatrix}$$

M1A1

[8]