



Cambridge International AS & A Level

CANDIDATE NAME



CENTRE NUMBER

--	--	--	--	--

CANDIDATE NUMBER

--	--	--	--



FURTHER MATHEMATICS

9231/33

Paper 3 Further Mechanics

May/June 2024

1 hour 30 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.
- Where a numerical value for the acceleration due to gravity (g) is needed, use 10 ms^{-2} .

INFORMATION


- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.





1



Two smooth uniform spheres A and B of equal radii have masses m and $2m$ respectively. The two spheres are moving on a smooth horizontal surface when they collide with speeds u and $\frac{1}{2}u$ respectively. Immediately before the collision, A 's direction of motion is along the line of centres, and B 's direction of motion makes an angle θ with the line of centres (see diagram).

As a result of the collision, the direction of motion of A is reversed and its speed is reduced to $\frac{1}{4}u$. The direction of motion of B again makes an angle θ with the line of centres, but on the opposite side of the line of centres. The speed of B is unchanged.

Find the value of the coefficient of restitution between the spheres. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN



- 2 A particle P of mass m is attached to one end of a light elastic string of natural length a and modulus of elasticity $2mg$. A particle Q of mass km is attached to the other end of the string. Particle P lies on a smooth horizontal table. The string has part of its length in contact with the table and then passes through a small smooth hole H in the table.

Particle P moves in a horizontal circle on the surface of the table with constant speed $\sqrt{\frac{1}{2}ga}$. Particle Q hangs in equilibrium vertically below the hole with $HQ = \frac{1}{4}a$.

- (a) Find, in terms of a , the extension in the string. [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

- (b) Find the value of k . [2]

.....

.....

.....

.....

.....





3 A particle P of mass m is attached to one end of a light inextensible string of length a . The other end of the string is attached to a fixed point O . When the particle is hanging vertically below O , it is projected horizontally with speed u so that it begins to move along a circular path. When P is at the lowest point of its motion, the tension in the string is T . When OP makes an angle θ with the upward vertical, the tension in the string is S .

(a) Show that $S = T - 3mg(1 + \cos \theta)$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN




(b) Given that $u = \sqrt{4ag}$, find the value of $\cos \theta$ when the string goes slack. [2]

Handwriting practice lines consisting of multiple horizontal dotted lines for writing the answer to question (b).

DO NOT WRITE IN THIS MARGIN





A light spring of natural length a and modulus of elasticity kmg is attached to a fixed point O on a smooth plane inclined to the horizontal at an angle θ , where $\sin \theta = \frac{3}{4}$. A particle of mass m is attached to the lower end of the spring and is held at the point A on the plane, where $OA = 2a$ and OA is along a line of greatest slope of the plane (see diagram).

The particle is released from rest and is moving with speed V when it passes through the point B on the plane, where $OB = \frac{3}{2}a$. The speed of the particle is $\frac{1}{2}V$ when it passes through the point C on the plane, where $OC = \frac{3}{4}a$.

Find the value of k .

[7]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





Handwriting practice area with horizontal dotted lines.

DO NOT WRITE IN THIS MARGIN





The object ABC is suspended from C . In its equilibrium position, the side AB makes an angle θ with the vertical, where $\tan \theta = \frac{6}{5}$.

(b) Find x in terms of a . [4]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN





(b) It is given that the velocity of P after 5 seconds is perpendicular to the initial velocity.

Find, in either order, the value of u and the value of $\sin \theta$. [5]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN





7 A parachutist of mass m kg opens his parachute when he is moving vertically downwards with a speed of 50 ms^{-1} . At time t s after opening his parachute, he has fallen a distance x m from the point where he opened his parachute, and his speed is $v \text{ ms}^{-1}$. The forces acting on him are his weight and a resistive force of magnitude mv N.

(a) Find an expression for v in terms of t . [6]

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN

DO NOT WRITE IN THIS MARGIN





(b) Find an expression for x in terms of t .

[3]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

(c) Find the distance that the parachutist has fallen, since opening his parachute, when his speed is 15 m s^{-1} . [2]

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

DO NOT WRITE IN THIS MARGIN





Additional page

If you use the following page to complete the answer to any question, the question number must be clearly shown.

A large area of the page consisting of 22 horizontal dotted lines, intended for writing an answer to a question.



* 0019655479115 *



DO NOT WRITE IN THIS MARGIN





BLANK PAGE

DO NOT WRITE IN THIS MARGIN

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in the Cambridge Assessment International Education Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge Assessment International Education is part of Cambridge Assessment. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which is a department of the University of Cambridge.



DO NOT WRITE IN THIS MARGIN