

**Q1.**

- 10 (i) Express  $2x^2 - 4x + 1$  in the form  $a(x + b)^2 + c$  and hence state the coordinates of the minimum point,  $A$ , on the curve  $y = 2x^2 - 4x + 1$ . [4]

The line  $x - y + 4 = 0$  intersects the curve  $y = 2x^2 - 4x + 1$  at points  $P$  and  $Q$ . It is given that the coordinates of  $P$  are  $(3, 7)$ .


- (ii) Find the coordinates of  $Q$ . [3]  
(iii) Find the equation of the line joining  $Q$  to the mid-point of  $AP$ . [3]

**Q2.**

- 2 Find the set of values of  $m$  for which the line  $y = mx + 4$  intersects the curve  $y = 3x^2 - 4x + 7$  at two distinct points. [5]

**Q3.**

5



The diagram shows the curve  $y = 7\sqrt{x}$  and the line  $y = 6x + k$ , where  $k$  is a constant. The curve and the line intersect at the points  $A$  and  $B$ .

- (i) For the case where  $k = 2$ , find the  $x$ -coordinates of  $A$  and  $B$ . [4]  
(ii) Find the value of  $k$  for which  $y = 6x + k$  is a tangent to the curve  $y = 7\sqrt{x}$ . [2]

**Q4.**

- 10 The equation of a line is  $2y + x = k$ , where  $k$  is a constant, and the equation of a curve is  $xy = 6$ .
- (i) In the case where  $k = 8$ , the line intersects the curve at the points  $A$  and  $B$ . Find the equation of the perpendicular bisector of the line  $AB$ . [6]  
(ii) Find the set of values of  $k$  for which the line  $2y + x = k$  intersects the curve  $xy = 6$  at two distinct points. [3]

**Q5.**

- 7 A curve has equation  $y = x^2 - 4x + 4$  and a line has equation  $y = mx$ , where  $m$  is a constant.
- (i) For the case where  $m = 1$ , the curve and the line intersect at the points  $A$  and  $B$ . Find the coordinates of the mid-point of  $AB$ . [4]
- (ii) Find the non-zero value of  $m$  for which the line is a tangent to the curve, and find the coordinates of the point where the tangent touches the curve. [5]

**Q6.**


- 2 A curve has equation  $y = 3x^3 - 6x^2 + 4x + 2$ . Show that the gradient of the curve is never negative. [3]

**Q7.**

- 9 A line has equation  $y = kx + 6$  and a curve has equation  $y = x^2 + 3x + 2k$ , where  $k$  is a constant.
- (i) For the case where  $k = 2$ , the line and the curve intersect at points  $A$  and  $B$ . Find the distance  $AB$  and the coordinates of the mid-point of  $AB$ . [5]
- (ii) Find the two values of  $k$  for which the line is a tangent to the curve. [4]

**Q8.**

3



The diagram shows the curve  $y = 2x^5 + 3x^3$  and the line  $y = 2x$  intersecting at points  $A$ ,  $O$  and  $B$ .

- (i) Show that the  $x$ -coordinates of  $A$  and  $B$  satisfy the equation  $2x^4 + 3x^2 - 2 = 0$ . [2]
- (ii) Solve the equation  $2x^4 + 3x^2 - 2 = 0$  and hence find the coordinates of  $A$  and  $B$ , giving your answers in an exact form. [3]

**Q9.**

- 7 (i) A straight line passes through the point  $(2, 0)$  and has gradient  $m$ . Write down the equation of the line. [1]
- (ii) Find the two values of  $m$  for which the line is a tangent to the curve  $y = x^2 - 4x + 5$ . For each value of  $m$ , find the coordinates of the point where the line touches the curve. [6]
- (iii) Express  $x^2 - 4x + 5$  in the form  $(x + a)^2 + b$  and hence, or otherwise, write down the coordinates of the minimum point on the curve. [2]

**Q10.**

10 A straight line has equation  $y = -2x + k$ , where  $k$  is a constant, and a curve has equation  $y = \frac{2}{x-3}$ .

(i) Show that the  $x$ -coordinates of any points of intersection of the line and curve are given by the equation  $2x^2 - (6 + k)x + (2 + 3k) = 0$ . [1]

(ii) Find the two values of  $k$  for which the line is a tangent to the curve. [3]

The two tangents, given by the values of  $k$  found in part (ii), touch the curve at points  $A$  and  $B$ .

(iii) Find the coordinates of  $A$  and  $B$  and the equation of the line  $AB$ . [6]

**Q11.**

1 Solve the inequality  $x^2 - x - 2 > 0$ . [3]