


Chapter 4 Differentiation

May/June 2002

- 5 The equation of a curve is $y = 2 \cos x + \sin 2x$. Find the x -coordinates of the stationary points on the curve for which $0 < x < \pi$, and determine the nature of each of these stationary points. [7]

10



The function f is defined by $f(x) = (\ln x)^2$ for $x > 0$. The diagram shows a sketch of the graph of $y = f(x)$. The minimum point of the graph is A . The point B has x -coordinate e .

- (i) State the x -coordinate of A . [1]
- (ii) Show that $f''(x) = 0$ at B . [4]
- (iii) Use the substitution $x = e^u$ to show that the area of the region bounded by the x -axis, the line $x = e$, and the part of the curve between A and B is given by

$$\int_0^1 u^2 e^u du. \quad [3]$$

- (iv) Hence, or otherwise, find the exact value of this area. [3]

Oct/Nov 2002

- 4 The curve $y = e^x + 4e^{-2x}$ has one stationary point.
- (i) Find the x -coordinate of this point. [4]
- (ii) Determine whether the stationary point is a maximum or a minimum point. [2]

May/June 2003

8 The equation of a curve is $y = \ln x + \frac{2}{x}$, where $x > 0$.

(i) Find the coordinates of the stationary point of the curve and determine whether it is a maximum or a minimum point. [5]

(ii) The sequence of values given by the iterative formula

$$x_{n+1} = \frac{2}{3 - \ln x_n},$$

with initial value $x_1 = 1$, converges to α . State an equation satisfied by α , and hence show that α is the x -coordinate of a point on the curve where $y = 3$. [2]

(iii) Use this iterative formula to find α correct to 2 decimal places, showing the result of each iteration. [3]

Oct/Nov 2003

4 The equation of a curve is

$$\sqrt{x} + \sqrt{y} = \sqrt{a},$$

where a is a positive constant.

(i) Express $\frac{dy}{dx}$ in terms of x and y . [3]

(ii) The straight line with equation $y = x$ intersects the curve at the point P . Find the equation of the tangent to the curve at P . [3]

May/June 2004


3 Find the gradient of the curve with equation

$$2x^2 - 4xy + 3y^2 = 3,$$

at the point $(2, 1)$. [4]

Oct/Nov 2004

7



The diagram shows the curve $y = x^2 e^{-\frac{1}{2}x}$.

- (i) Find the x -coordinate of M , the maximum point of the curve. [4]
- (ii) Find the area of the shaded region enclosed by the curve, the x -axis and the line $x = 1$, giving your answer in terms of e . [5]

Oct/Nov 2005

- 3 The equation of a curve is $y = x + \cos 2x$. Find the x -coordinates of the stationary points of the curve for which $0 \leq x \leq \pi$, and determine the nature of each of these stationary points. [7]

May/June 2006

- 3 The parametric equations of a curve are

$$x = 2\theta + \sin 2\theta, \quad y = 1 - \cos 2\theta.$$

Show that $\frac{dy}{dx} = \tan \theta$. [5]

Oct/Nov 2006

- 3 The curve with equation $y = 6e^x - e^{3x}$ has one stationary point.
- (i) Find the x -coordinate of this point. [4]
- (ii) Determine whether this point is a maximum or a minimum point. [2]
- 6 The equation of a curve is $x^3 + 2y^3 = 3xy$.
- (i) Show that $\frac{dy}{dx} = \frac{y - x^2}{2y^2 - x}$. [4]
- (ii) Find the coordinates of the point, other than the origin, where the curve has a tangent which is parallel to the x -axis. [5]

May/June 2007

- 3 The equation of a curve is $y = x \sin 2x$, where x is in radians. Find the equation of the tangent to the curve at the point where $x = \frac{1}{4}\pi$. [4]

Oct/Nov 2007

- 4 The curve with equation $y = e^{-x} \sin x$ has one stationary point for which $0 \leq x \leq \pi$.
- (i) Find the x -coordinate of this point. [4]
- (ii) Determine whether this point is a maximum or a minimum point. [2]

May/June 2008

- 6 The equation of a curve is $xy(x+y) = 2a^3$, where a is a non-zero constant. Show that there is only one point on the curve at which the tangent is parallel to the x -axis, and find the coordinates of this point. [8]

Oct/Nov 2008

- 3 The curve $y = \frac{e^x}{\cos x}$, for $-\frac{1}{2}\pi < x < \frac{1}{2}\pi$, has one stationary point. Find the x -coordinate of this point. [5]
- 4 The parametric equations of a curve are
- $$x = a(2\theta - \sin 2\theta), \quad y = a(1 - \cos 2\theta).$$
- Show that $\frac{dy}{dx} = \cot \theta$. [5]

May/June 2009

- 6 The parametric equations of a curve are
- $$x = a \cos^3 t, \quad y = a \sin^3 t,$$
- where a is a positive constant and $0 < t < \frac{1}{2}\pi$.
- (i) Express $\frac{dy}{dx}$ in terms of t . [3]
- (ii) Show that the equation of the tangent to the curve at the point with parameter t is
- $$x \sin t + y \cos t = a \sin t \cos t. [3]$$
- (iii) Hence show that, if this tangent meets the x -axis at X and the y -axis at Y , then the length of XY is always equal to a . [2]

Oct/Nov 2009/31

- 4 A curve has equation $y = e^{-3x} \tan x$. Find the x -coordinates of the stationary points on the curve in the interval $-\frac{1}{2}\pi < x < \frac{1}{2}\pi$. Give your answers correct to 3 decimal places. [6]

Oct/Nov 2009/32


3 The equation of a curve is $x^3 - x^2y - y^3 = 3$.

(i) Find $\frac{dy}{dx}$ in terms of x and y . [4]

(ii) Find the equation of the tangent to the curve at the point $(2, 1)$, giving your answer in the form $ax + by + c = 0$. [2]

May/June 2010/31

9



The diagram shows the curve $y = \sqrt{\left(\frac{1-x}{1+x}\right)}$.

(i) By first differentiating $\frac{1-x}{1+x}$, obtain an expression for $\frac{dy}{dx}$ in terms of x . Hence show that the gradient of the normal to the curve at the point (x, y) is $(1+x)\sqrt{1-x^2}$. [5]

(ii) The gradient of the normal to the curve has its maximum value at the point P shown in the diagram. Find, by differentiation, the x -coordinate of P . [4]