

CANDIDATE
NAME


--

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--



PHYSICAL SCIENCE

8780/03

Paper 3 Structured Questions

October/November 2014

1 hour 30 minutes

Candidates answer on the Question Paper.

Additional Materials: Data Booklet

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

A Data Booklet is provided.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **18** printed pages and **2** blank pages.

Answer **all** the questions in the spaces provided.

Relevant Data, Formulae and the Periodic Table are provided in the Data Booklet.

- 1 At room temperature, both sodium metal and sodium chloride are crystalline solids which contain ions.

- (a) On Fig. 1.1, mark the charge on each ion in sodium metal and in sodium chloride.




Fig. 1.1

[2]

- (b) (i) Describe how the ions are held together in solid sodium metal.

.....
 [1]

- (ii) Describe how the ions are held together in solid sodium chloride.

.....
 [1]

- (c) (i) Explain why both molten and solid sodium metal conduct electricity.

.....
 [1]

- (ii) Explain why molten sodium chloride conducts electricity, but solid sodium chloride does not.

.....

 [2]

[Total: 7]

- 2 In this question you are to estimate the average rate of energy loss as an airliner descends from its cruising height, lands, and comes to rest on the runway.

The airliner starts to lose height 30 minutes before it comes to rest on the runway.

Table 2.1 shows the quantities and their estimated values needed in order to calculate the energy lost by the airliner.

Table 2.1

quantity	estimated value
cruising height above the runway	8 000 m
mass of the loaded airliner	$3 \times 10^5 \text{ kg}$
cruising speed	200 m s^{-1}

- (a) Use the values in Table 2.1 to estimate

- (i) the total energy loss of the airliner,

total energy loss = J [4]

- (ii) the average rate of energy loss.

average rate of energy loss = W [2]

(b) (i) The speed of the airliner as its wheels touch the runway is 250 km h^{-1} .

The average braking force of the airliner during landing is 400 kN .

Calculate the distance the airliner travels along the runway before it comes to rest.


distance = m [3]

(ii) Suggest why the runway needs to be significantly longer than the distance calculated in **(b)(i)**.

.....
.....[1]

[Total: 10]

- 3 Nitroglycerine, $C_3H_5N_3O_9$, is an explosive. On detonation, it decomposes rapidly to form a large number of gaseous molecules. The equation for this decomposition is given below.



- (a) State what is meant by the term *one mole* of molecules.

.....
[1]

- (b) A sample of nitroglycerine was detonated and produced 0.350 g of oxygen gas.

- (i) Show that the amount of oxygen gas produced is 1.09×10^{-2} mol and hence deduce the total number of moles of gas formed in this reaction.

total number of moles of gas = [2]

- (ii) Calculate the number of moles, and the mass, of nitroglycerine detonated.

number of moles of nitroglycerine =

mass of nitroglycerine = g
 [2]

- (c) A second sample of nitroglycerine is placed in a strong, sealed container and detonated. The container does not break or change volume.

The volume of the container is $1.00 \times 10^{-3} \text{ m}^3$.

The resulting decomposition produces a total of 0.873 mol of gaseous products at a temperature of 1100 K.

- (i) State the ideal gas equation.

.....[1]

- (ii) Use the ideal gas equation to calculate the pressure in the container after detonation. Give the unit for your answer.

(the gas constant, $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$)

pressure = unit [3]

[Total: 9]

- 4 Blood pressure is defined as the excess pressure in a person's arteries compared to atmospheric pressure.

Fig. 4.1 and Fig. 4.2 show apparatus that can be used to measure blood pressure. The cuff is placed around the patient's arm and air is pumped into the tube in the cuff. This causes the mercury to be pushed around the U-tube. The valve, when closed, prevents air from coming back out of the cuff.




Fig. 4.1




Fig. 4.2

- (a) Explain, in terms of the kinetic molecular model,
- (i) why the atmosphere exerts a pressure on the mercury at point **A**,

.....

.....

.....

.....[3]

(ii) why the pressure is greater at **C** than at **A**.

.....
.....
.....[2]

(b) The air in the cuff is released slowly by opening the valve. The valve is closed again when the pressure at **C** is equal to the maximum pressure of the patient's blood.

For a particular patient, the distance **BD** is 280 mm and the distance **AD** is 395 mm.

The density of mercury is 13600 kg m^{-3} .

Calculate the blood pressure of the patient in pascal.

pressure = Pa [2]

[Total: 7]

- 5 Fig. 5.1 shows an experiment designed to demonstrate interference of visible light. In card **A** there is a single slit. In card **B** there are two slits, separated by distance x .

A series of bright and dark fringes is observed on the screen.




Fig. 5.1

- (a) Explain why the single slit in card **A** must be narrow.

.....

[2]

- (b) State the effect of using a different card **B** which has

- (i) a smaller distance x between the slits,

.....[1]

- (ii) narrower slits.

.....[1]

(c) (i) State what is meant by *monochromatic light*.

.....
.....[1]


(ii) In the experiment shown in Fig. 5.1, white light is used instead of monochromatic light.
Predict what would be observed on the screen and give a reason for your answer.

.....
.....
.....
.....[2]

[Total: 7]

- 6 The dimerisation of nitrogen dioxide, NO_2 , to form dinitrogen tetroxide, N_2O_4 , is exothermic.

This reaction is reversible as shown in the equation below.



In a sealed container, a dynamic equilibrium mixture of the two gases will eventually form.

- (a) Explain what is meant by the term *dynamic equilibrium* as applied to a chemical reaction.

.....

 [2]

- (b) A teacher prepared three sealed glass containers, **A**, **B** and **C**, each filled with nitrogen dioxide. The containers were left at a temperature of 35°C until the contents reached equilibrium. After some time, the contents of each container were light brown in colour.

- (i) The teacher then heated container **A** to 45°C and cooled container **B** to 25°C . Container **C** was kept at 35°C .

Suggest how, if at all, the appearance of the contents of **A** and **B** would change relative to that of the contents of **C**.

Explain your answers.

.....

 [3]

- (ii) Explain why the contents of **A** take less time to reach equilibrium than the contents of **B**.

.....

 [3]

- (c) (i) Table 6.1 gives some values for enthalpy change of formation ΔH_f° .

Table 6.1

compound	$\Delta H_f^\circ / \text{kJ mol}^{-1}$
$\text{NO}_2(\text{g})$	33.18
$\text{N}_2\text{O}_4(\text{g})$	9.16

Use the data in Table 6.1 to calculate the enthalpy change for the dimerisation of NO_2 .

enthalpy change = kJ mol^{-1} [1]

- (ii) Write an equation for the reaction whose enthalpy change is the enthalpy change of formation of nitrogen dioxide.
Include state symbols.

.....[1]

- (d) In the laboratory, nitrogen dioxide can be produced by strongly heating solid magnesium nitrate, $\text{Mg}(\text{NO}_3)_2$. Write an equation for this reaction.

.....[2]

[Total: 12]

- 7 In the apparatus shown in Fig. 7.1, the cathode and anode are connected to a power supply fitted with a safety resistor. Electrons are emitted from the hot cathode and are accelerated towards the anode.

The ammeter reads $24 \mu\text{A}$.




Fig. 7.1

- (a) Calculate the total resistance in the circuit.

total resistance = Ω [2]

- (b) (i) Calculate the power dissipated in the safety resistor.

power = W [1]

- (ii) Calculate the output power from the power supply and hence the power dissipated inside the glass container.

power dissipated inside the glass container = W [2]

- (c) (i) Deduce the charge hitting the anode each second and give the unit.

charge = unit [2]

- (ii) Calculate the number of electrons hitting the anode each second.
(electronic charge = $1.6 \times 10^{-19} \text{ C}$)


number of electrons = [1]

- (iii) Use your answers to (b)(ii) and (c)(ii) to calculate the energy received by each electron as it moves from the cathode to the anode.

energy = J [1]

[Total: 9]

- 8 The structural and displayed formulae of compound **P** are shown below.



Some reactions of compound **P** are shown in Fig. 8.1.




Fig. 8.1

- (a) Name the type of reaction involved in the following conversions.

P into **Q**

P into **T**

[2]

- (b) Compound **T** has the following composition by mass: C, 55.81 %; H, 6.98 %; O, 37.21 %.

Calculate the empirical formula of **T**.

empirical formula of **T** = [2]

- (c) (i) In the appropriate boxes in Fig. 8.1, draw the structural formulae for the organic compounds **R**, **S** and **T**. [3]

- (ii) In the space below, draw the **displayed** formula of compound **Q**.

[1]

- (d) Consider the functional groups present in **P** and **Q**.

- (i) State a test that would allow you to distinguish between pure samples of these two compounds. Describe the observations you would make if your chosen test is performed on **P** and on **Q**.

test

observation with **P**

.....

observation with **Q**

.....

[3]

- (ii) State the functional group identified by your test.

.....[1]

[Total: 12]

9 (a) (i) State Kirchhoff's second law.

.....
[1]

(ii) Explain the relationship between Kirchhoff's second law and the principle of conservation of energy.

.....
[1]

(b) Two cells are connected to a network of resistors as shown in Fig. 9.1. The cells and ammeter have negligible resistance.




Fig. 9.1

(i) Use Kirchhoff's first law to deduce the relationship between I_1 , I_2 and I_3 .

[1]

(ii) Use Kirchhoff's second law for loop **XWVY** to deduce the relationship between E_2 , I_3 and R .

[1]

(iii) Use Kirchhoff's second law for loop **SWUT** to express E_1 in terms of R , I_1 and I_3 .

[1]

(iv) For a particular pair of cells, E_1 and E_2 , the ammeter shows a zero reading.

Show that the ratio $E_2 : E_1 = 4 : 9$.

[2]

[Total: 7]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.