



**1. Nov/2023/Paper\_9702/11/No.38**

Two alpha-particles with the same kinetic energy are moving towards, and are then deflected by, a gold nucleus.



Which diagram could show the paths of the two alpha-particles?



**2. Nov/2023/Paper\_9702/11/No.39**

Which nuclide is formed when  $^{10}_6\text{C}$  undergoes  $\beta^+$  decay?

- A  $^{11}_6\text{C}$       B  $^9_6\text{C}$       C  $^{10}_5\text{B}$       D  $^{10}_7\text{N}$

**3. Nov/2023/Paper\_9702/11/No.40**

A particular hadron is composed of three quarks and has zero charge.

What is a possible quark composition of the hadron?

- A down, down, strange
- B up, down, strange
- C up, up, down
- D up, up, strange

4. Nov/2023/Paper\_ 9702/12/No.30

A fine mist of oil droplets is sprayed into air. As the oil droplets leave the nozzle of the spraying device they can become electrically charged.

What is **not** a possible value for the charge on an oil droplet?

- A zero
- B  $1.0 \times 10^{-19} \text{ C}$
- C  $4.8 \times 10^{-19} \text{ C}$
- D  $8.0 \times 10^{-19} \text{ C}$

5. Nov/2023/Paper\_ 9702/12/No.38

Two nuclides are different isotopes of the same element.

Which statement about the nuclides is correct?

- A Neutral atoms of the nuclides have the same number of electrons.
- B Nuclei of the nuclides have different numbers of protons.
- C Nuclei of the nuclides have the same number of nucleons.
- D Nuclei of the nuclides have the same number of neutrons.

6. Nov/2023/Paper\_ 9702/12/No.39

The charge-to-mass ratio  $r$  of a particle is given by the equation shown.

$$r = \frac{\text{charge on particle}}{\text{mass of particle}}$$

The value of  $r$  is determined for an  $\alpha$ -particle, a  $\beta^+$  particle and a proton p.

Which list shows the particles in order of increasing magnitude of  $r$  from left to right?

- A  $\alpha \rightarrow \beta^+ \rightarrow \text{p}$
- B  $\alpha \rightarrow \text{p} \rightarrow \beta^+$
- C  $\text{p} \rightarrow \alpha \rightarrow \beta^+$
- D  $\text{p} \rightarrow \beta^+ \rightarrow \alpha$

7. Nov/2023/Paper\_ 9702/12/No.40

Which combination of up (u) and down (d) quarks forms a neutron?

- A u u u
- B u u d
- C u d d
- D d d d

8. Nov/2023/Paper\_9702/13/No.38

The table shows the number of nucleons and the total number of particles (protons, neutrons and electrons) in neutral atoms of four nuclides W, X, Y and Z.

|   | number of nucleons | total number of particles |
|---|--------------------|---------------------------|
| W | 19                 | 30                        |
| X | 19                 | 31                        |
| Y | 21                 | 32                        |
| Z | 22                 | 31                        |

Which two nuclides are isotopes of each other?

- A W and X      B W and Y      C X and Z      D Y and Z

9. Nov/2023/Paper\_9702/13/No.39

When a sample of a radioactive isotope decays by  $\alpha$ -particle emission, the  $\alpha$ -particles emitted have a single discrete energy.

When a sample of a radioactive isotope decays by  $\beta^-$  particle emission, the  $\beta^-$  particles emitted have a continuous range of energies.

What is the explanation for this?

- A An antineutrino is emitted with a  $\beta^-$  particle but not with an  $\alpha$ -particle.  
B An antineutrino is emitted with an  $\alpha$ -particle but not with a  $\beta^-$  particle.  
C The  $\alpha$ -particles have much more energy than the  $\beta^-$  particles.  
D The  $\beta^-$  particles have much more energy than the  $\alpha$ -particles.

10. Nov/2023/Paper\_9702/13/No.40

Some particles are a combination of three quarks.

Which combination of quarks does **not** result in a particle with a charge of either  $+1.6 \times 10^{-19} \text{ C}$  or zero?

- A up, down, down  
B up, strange, strange  
C up, up, down  
D up, up, up

- (a) The results of the  $\alpha$ -particle scattering experiment led to the development of the nuclear model for the atom.

State the results that suggested that most of the mass of the atom is concentrated in a very small region and most of the atom is empty space.

.....

.....

.....

..... [2]

- (b) State the composition of  $\gamma$ -radiation.

..... [1]

- (c) Table 7.1 lists the names of three particles and possible classifications for them.

**Table 7.1**

| particle name | classification |        |        |
|---------------|----------------|--------|--------|
|               | baryon         | hadron | lepton |
| neutrino      |                |        |        |
| neutron       |                |        |        |
| positron      |                |        |        |

Complete Table 7.1 by placing ticks ( $\checkmark$ ) in the boxes to indicate the classifications that apply to each particle. [2]

(d) The discovery of a particle with an unusual charge was an important step in the development of the theory of quarks. The particle is a hadron with a mass of  $2.19 \times 10^{-27}$  kg and a charge of  $+2e$ , where  $e$  is the elementary charge.


(i) Calculate the mass, in  $u$ , of the particle. Give your answer to three significant figures.

mass = .....  $u$  [1]

(ii) Determine a possible quark composition of a hadron with a charge of  $+2e$ . Explain your reasoning.

[2]

[Total: 8]



(a) In the following list, underline **all** the particles that are **not** fundamental.

antineutrino                  baryon                  nucleon                  positron                  [1]

(b) A nucleus of thorium-230 ( ${}^{230}_{90}\text{Th}$ ) decays in stages, by emitting  $\alpha$ -particles and  $\beta^-$  particles, to form a nucleus of lead-206 ( ${}^{206}_{82}\text{Pb}$ ).

Determine the total number of  $\alpha$ -particles and the total number of  $\beta^-$  particles that are emitted during the sequence of decays that form the nucleus of lead-206 from the nucleus of thorium-230.

number of  $\alpha$ -particles = .....

number of  $\beta^-$  particles = ..... [2]

(c) A meson has a charge of  $-1e$ , where  $e$  is the elementary charge. The quark composition of the meson includes a charm antiquark.

State and explain a possible flavour (type) of the other quark in the meson.




.....  
..... [2]

[Total: 5]

13. Nov/2023/Paper\_9702/21/No.8

- (a) The nuclide  ${}_{12}^{23}\text{Mg}$  is an isotope of magnesium that undergoes  $\beta^+$  decay to form a new nuclide X according to the equation



Four numbers are missing from the equation.

- (i) For the nuclide  ${}_{12}^{23}\text{Mg}$ , state what is represented by the numbers 23 and 12.

23 represents: .....


12 represents: .....

[2]

- (ii) Complete the equation by inserting the missing numbers. [2]

- (iii) State the name of the group (class) of fundamental particles to which the positron and neutrino belong.

..... [1]



- (b) A radioactive source emits particles from its nuclei when it decays. Fig. 8.1 shows, for the source, the variation with kinetic energy of the number of particles emitted.




Fig. 8.1

State how Fig. 8.1 shows that these nuclei do **not** undergo beta-decay.

.....  
..... [1]

[Total: 6]

