
Centre Number	Candidate Number	Name	SN. Por	as l
		GE INTERNATIONAL EX ertificate of Secondary E	XAMINATIONS Education 0610/06	abaCambride
BIOLOGY				
Paper 6 Alte	rnative to Practical		May/June 2004	1
			1 hou	
	wer on the Question Pap dditional Materials.	ber.	Thou	
	er, candidate number an	nd name on all the work you		
o not use staples, pap inswer <b>all</b> questions. It the end of the examir	cil for any diagrams, grap er clips, highlighters, glu nation, fasten all your wo	phs or rough working. e or correction fluid.		
You may use a soft pen No not use staples, pap Answer <b>all</b> questions. At the end of the examir	cil for any diagrams, grap er clips, highlighters, glue nation, fasten all your wo given in brackets [ ] at t a label, look at the e incorrect or ur correct details e top of this page.	phs or rough working. e or correction fluid. ork securely together.		er's Use

**1** Fig. 1.1 shows the appearance of a cell from the epidermis of a leaf.





Three similar pieces of epidermis have been placed in different solutions and left submerged for 30 minutes. One solution was pure water, another contained 1.5% sugar solution and the third 5% sugar solution.

Figs. 1.2, 1.3 and 1.4 show a cell from each of these three pieces of epidermis.




Fig. 1.2

Fig. 1.3

Fig. 1.4

(These cells are all drawn to the same magnification)

(a) Measure the width of the cell contents along the lines drawn across each cell.

Fig. 1.1	•
Fig. 1.2	
Fig. 1.3	
Fig. 1.4	

[3]

www.papaCambridge.com

	42	
	3	For Examiner's
(b)	3 Suggest in which solution each of the cells, in Figs. 1.2, 1.3 and 1.4, was placed and the cells in Figs. 1.2 and 1.4 and 1.	Use
	Fig. 1.2	11dg
	explanation	Com
	Fig. 1.3	
	explanation	•
	Fig. 1.4	
	explanation	
	[8]	
	[Total : 11]	

- www.papaCambridge.com 2 Starch is broken down during digestion by an enzyme, amylase. The test for starch uses iodine solution. When all the starch has been completely digested, the orange brown iodine solution do not change colour.
  - (a) Three test tubes **A**, **B** and **C** each contained  $5 \text{ cm}^3$  of 1% starch solution.

To tube **A**,  $1 \text{ cm}^3$  of water and  $2 \text{ cm}^3$  of 1% amylase solution were added. The stop clock was started immediately and the mixture was stirred. Every minute a drop was removed from the mixture and added to iodine solution on a white tile.

To tube **B**,  $1 \text{ cm}^3$  of water an  $2 \text{ cm}^3$  of boiled and cooled 1% amylase solution were added.

The same method of testing was used.

To tube **C**,  $1 \text{ cm}^3$  of 0.5% sodium chloride solution and  $2 \text{ cm}^3$  of 1% amylase solution were added.

The same method of testing was used.

The colours observed are shown in Table 2.1.

These investigations were carried out at room temperature.

## Table 2.1

time/mins	tube A	tube <b>B</b>	tube C
1	black	black	black
2	black	black	dark brown
3	black	black	dark brown
4	black	black	lighter brown
5	dark brown	black	lighter brown
6	dark brown	black	orange brown
7	dark brown	black	orange brown
8	dark brown	black	orange brown
9	lighter brown	black	orange brown
10	lighter brown	black	orange brown
11	lighter brown	black	orange brown
12	lighter brown	black	orange brown
13	orange brown	black	orange brown
14	orange brown	black	orange brown
15	orange brown	black	orange brown


(i) State how long it took for the starch to be completely broken down in tubes A and С.

*tube* **A** ......[2]

4

(ii)	5 Describe the effect on the reaction of using sodium chloride solution. [2]	For Examiner's Use
(iii)	[2] State the purpose of tube <b>B</b> in this investigation.	Age com
	the human alimentary canal, starch is broken down by amylase. The pH of the	
	nentary canal varies between pH 2 and pH 8.5. ggest how you could investigate the effect of pH on the activity of amylase.	
	[5]	

[Total : 10]



Label the position of the seed.

	the second second
	7
(ii)	7 Calculate the magnification of your drawing using the maximum length of the Fig. 3.1 that you have drawn. maximum length of your drawing of one fruit
	maximum length of your drawing of one fruit cm
	maximum length of the same fruit in Fig. 3.1 cm
	working:
	magnification[2]
The	e background in Fig. 3.1 is a grid, with squares of 1 mm x 1 mm.
(iii)	Determine the surface area of one of the fruits including the wing-like extension. Explain how you worked out your answer.
	[3]


www.papacambridge.com (b) Using paper and plasticine to represent a wind dispersed fruit, such as A possible to adjust the size of the wing. The effect of different wing size was investig by dropping a model fruit with different surface areas, from the same height in the same wind conditions. The horizontal distance travelled by the model was measured. Each model fruit was dropped five times.

The results are shown below.

## Table 3.1

surface area of	distance travelled/cm				mean	
wing-like extension / cm <sup>2</sup>	drop 1	drop 2	drop 3	drop 4	drop 5	distance travelled/cm
32	30	40	20	15	20	
64	20	30	30	25	40	
96	30	40	26	50	35	
128	45	20	40	45	65	
160	72	40	54	50	34	

- (i) Complete Table 3.1 by calculating the mean (average) distance travelled by the model fruits. [2]
- (ii) Plot the mean distance the model fruit travelled horizontally against the surface area of the model as a line graph. [4]



	9 Describe the relationship between the surface area and the mean of travelled.	For Examiner's
(iii)	Describe the relationship between the surface area and the mean a travelled.	Use
	[2]	ase com
(iv)	Outline the importance of seed dispersal away from the parent plant.	
	[2]	
	[Total : 19]	



**BLANK PAGE** 

10



BLANK PAGE

11



**BLANK PAGE** 

12

Every reasonable effort has been made to trace all copyright holders where the publishers (i.e. UCLES) are aware that third-party material has been reproduced. The publishers would be pleased to hear from anyone whose rights they have unwittingly infringed.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.