

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

Middle con

•	
•	
•	
•	
1	
`	
J	
J	
`	
1	
•	

CANDIDATE NAME							
CENTRE NUMBER				CANDIDATI NUMBER	E		

BIOLOGY 0610/31

Paper 3 Extended

October/November 2012

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Exam	iner's Use
1	
2	
3	
4	
5	
6	
Total	

This document consists of 22 printed pages and 2 blank pages.



www.PapaCambridge.com

Question 1 begins on page 3.

ropods.

rids.

rids.

1 Arachnids, crustaceans, insects and myriapods are all classified as arthropods.

Scorpions, such as *Heterometrus swammerdami* shown in Fig. 1.1, are arachnids.




Fig. 1.1

(a)	State three features, shown by <i>H. swammerdami</i> and visible in Fig. 1.1, that arachnids
	share with other arthropods.

1	
2	
2	
3	[3

(b) Fig. 1.2 shows seven species of arachnid.




Fig. 1.2

Key

	5 key to identify each species. Write the letter of de the key. One has been done for you. Key	each species (A to G) in the Abaliella dicranotarsalis	bridge
1 (a)	Abdomen with a tail	Abaliella dicranotarsalis E	
(b)	Abdomen without a tail	go to 2	
2 (a)	Legs much longer than abdomen and cephalothorax	go to 3	
(b)	Legs not much longer than abdomen and cephalothorax	go to 4	
3 (a)	Hairs on the legs	Tegenaria domestica	
(b)	No hairs on the legs	Odielus spinosus	-
4 (a)	Cephalothorax or abdomen segmented	Chelifer tuberculatus	
(b)	Cephalothorax and abdomen not segmented	go to 5	
5 (a)	Abdomen and cephalothorax about the same size	Poecilotheria regalis	
(b)	Abdomen larger than cephalothorax	go to 6	
6 (a)	Body covered in long hairs	Tyroglyphus longior	
(b)	Body not covered in hairs	Ixodes hexagonus	

[4]

[Total: 7]

2 Blood flows through the hepatic portal vein from some organs to the liver.

Fig. 2.1 shows the hepatic portal vein and these organs.




Fig. 2.1

(a) Blood in the hepatic portal vein is deoxygenated.

oxygena	-	tne	DIOOG	ın	tne	nepatic	portai	vein	IS	deoxygenated	ratner	tnan
												[2]

www.PapaCambridge.com (b) Name four organs, other than the spleen, that are shown in Fig. 2.1 and from blood flows into the hepatic portal vein. 1 2 3 4 [4] (c) Describe the role of the hepatic portal vein in the transport of absorbed nutrients. (d) Explain how the liver is involved in regulating the composition of the blood and in protecting the body against toxic substances.

(e)	The spleen contains lymphatic tissue which is full of phagocytes and lymphocyte.
	Describe how phagocytes and lymphocytes protect the body against the spread disease-causing organisms.
	phagocytes
	lymphocytes
	[4]

[Total: 18]

www.PapaCambridge.com

Question 3 begins on page 10.

3 The ribcage and diaphragm are involved in the breathing mechanism to ventilate the

Fig. 3.1 is a flow chart that shows the changes that take place when breathing in.




Fig. 3.1

(a) Complete Fig. 3.1 by writing appropriate words in the spaces provided.

[6]

(b) Fig. 3.2 shows part of the epithelium that lines the trachea.




Fig. 3.2

Explain how the cells labelled ${\bf A}$ and ${\bf B}$ in Fig. 3.2 protect the gas exchange system.


Α	
В	
	[4

[Total: 10]

www.PanaCambridge.com

Question 4 begins on page 13.

(a) Complete the balanced chemical equation for photosynthesis.



A student investigated the effect of increasing the concentration of carbon dioxide on the rate of photosynthesis of Cabomba, an aquatic plant.

Fig. 4.1 shows the apparatus that the student used.




Fig. 4.1

The concentration of carbon dioxide in the water surrounding the plant was changed by adding different concentrations of sodium hydrogencarbonate solution to the water.

The student recorded the time taken for the meniscus to travel 50 mm down the tubing.

The rate of photosynthesis was calculated as:

rate of photosynthesis =
$$\frac{1000}{f}$$

where \mathbf{t} = time taken in seconds for the meniscus to travel 50 mm.

Table 4.1

The student's results are sho	14 nown in Table 4.1.	www.Por	Dal Cambridge
	Table 4.1		Orida
concentration of sodium hydrogencarbonate solution / mol per dm ³	t, time taken for meniscus to travel 50 mm / s	rate of photosynthesis (1000/t)	100
0.00	4998	0.20	
0.01	2500	0.40	
0.02	1175	0.85	
0.05	350	2.86	
0.07	201		
0.10	199	5.03	

(b)	Calculate	the	rate	of	photosynthesis	for	the	concentration	of	sodium
	hydrogenc	arbona	ate soli	ution	of 0.07 mol per dr	n^3 .				

Write your answer in Table 4.1. [1]

(c)	(i)	Explain why the lamp must be kept at a fixed distance from the syringe.	
			[2]
	(ii)	Explain what caused the meniscus to move down the capillary tubing.	
			[2]

(d) Fig. 4.2 is a partially completed graph of the student's results.

Complete the graph by labelling the axes, adding the missing point and drawing suitable line.




Fig. 4.2

	4	
	16 WW. D	
	700	
(e)	Explain, using the term limiting factors , the effect of carbon dioxide concentrathe rate of photosynthesis as shown by the student's results.	For vine
	Explain, using the term limiting factors , the effect of carbon dioxide concentrathe rate of photosynthesis as shown by the student's results. You will gain credit for using the data in the table and the graph to answer the question.	de.c
	[5]	
	[Total: 16]	


Table 5.1

Table 5.1 shows some information about air pollution. Table 5.1 pollutant source of air pollutant effect of pollutant on the environment			
pollutant	source of air pollutant	effect of pollutant on the environment	
	combustion of fossil fuels	increased greenhouse effect and global warming	
methane		increased greenhouse effect and global warming	
sulfur dioxide	combustion of high sulfur fuels	acid rain	
nitrogen oxides	fertilisers	acid rain	

(a)	Complete Table 5.1 by writing answers in the spaces indicated.	[2]
(b)	Explain how the increased greenhouse effect is thought to lead to global warming.	
		[3]

(c) Fig. 5.1 shows changes in the emissions of sulfur dioxide in Europe between 18 2004.

60






Fig. 5.1

dioxide in Europe between 1880 and 2004.
[4]

1880 1890 1900 1910 1920 1930 1940 1950 1960 1970 1980 1990 2000 2010

year

(i) Use the information in Fig. 5.1 to describe the changes in the emissions of sulfur

		Why.
		19 A. D.
	(ii)	Describe the effects of acid rain on the environment.
		[3]
	(iii)	Outline the methods that have been used to reduce the emissions of sulfur dioxide.
		[3]
		[Total: 15]
		[Total: To]
6		wers of pea plants, <i>Pisum sativum</i> , are produced for sexual reproduction. The are naturally self-pollinating, but they can be cross-pollinated by insects.
	(a) Exp	lain the difference between self-pollination and cross-pollination.
		[2]

	20
(b)	Explain the disadvantages for plants, such as <i>P. sativum</i> , of reproducing sexual.
	[4]
	a seeds develop inside pea pods after fertilisation. They contain starch. A gene controls production of an enzyme involved in the synthesis of starch grains.
	e allele, R , codes for an enzyme that produces normal starch grains. s results in seeds that are round.
	e allele, ${f r}$, does not code for the enzyme. The starch grains are not formed normally. This ults in seeds that are wrinkled.
Fig.	6.1 shows round and wrinkled pea seeds.
	round pea seed wrinkled pea seed
	Fig. 6.1
bre	e bred plants are homozygous for the gene concerned. A plant breeder had some pure d pea plants that had grown from round seeds and some pure bred plants that had wn from wrinkled seeds.
(c)	State the genotypes of the pure bred plants that had grown from round and from wrinkled seeds.
	round

wrinkled [1]

www.PapaCambridge.com These pure bred plants were cross-pollinated (cross 1) and the seeds collected. All the seeds were round. These round seeds were germinated, grown into adult pa (offspring 1) and self-pollinated (cross 2).

The pods on the offspring 1 plants contained both round and wrinkled seeds.

Further crosses (3 and 4) were carried out as shown in Table 6.1.

Table 6.1

cross		phenotype of seeds in the seed pods		ratio of round to
		round seeds	wrinkled seeds	wrinkled seeds
1	pure bred for round seeds x pure bred for wrinkled seeds	✓	×	1:0
2	offspring 1 self-pollinated	✓	✓	
3	offspring 1 x pure bred for round seeds			
4	offspring 1 x pure bred for wrinkled seeds			

(d) Complete Table 6.1 by indicating

- the type of seeds present in the pods with a tick (✓) or a cross (✗)
- the ratio of round to wrinkled seeds.

You may use the space below and on page 22 for any rough working.

[3]

		or	
3	,	iner	's
	1		
	00		•
7	1	0	
		.C	
	7	1	ク

(e)	Seed shape in peas is an example of discontinuous variation. Suggest one reason why seed shape is an example of discontinuous variation.
	[1]
Plaı	nts have methods to disperse their seeds over a wide area.
(f)	Explain the advantages of having seeds that are dispersed over a wide area,
	rol
	[3]

[Total: 14]

BLANK PAGE

www.PapaCambridge.com

24

BLANK PAGE

MMM. Papa Cambridge.com

 $Copyright\ Acknowledgements:$

Question 3 Figure 3.2

© Reference A18K2A – Electronmicrograph Peter Arnold, Inc. / Alamy

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.