

ZNOTES.ORG

UPDATED TO 2020-21 SYLLABUS

CAIE IGCSE


BIOLOGY (0610)

SUMMARIZED NOTES ON THE THEORY SYLLABUS

1. Organization of the Organism

1.1. Cell structure and Organization

- All living things are made of cells.
- All typical cells have:
 - Cell membrane: differentially or partially permeable to allow certain substances to enter and leave the cell.
 - Cytoplasm: where chemical reactions take place
 - Nucleus: contains DNA and controls the cell
 - Mitochondria: organelle where aerobic respiration happens
 - Ribosome: makes protein and can be found floating within the cytoplasm
- A typical animal cell (e.g. the liver cell) has all above
- Only plant cells have:
 - Vacuole: stores food & water & helps to maintain shape of cell
 - Cell wall: rigid to keep shape of cell
 - Chloroplasts: contain chlorophyll, which absorbs light energy for photosynthesis
- A typical plant cell (e.g. the palisade cell) has all the above things.



1.2. Levels of Organization

| Cell | Function | Adaptation(s) | Diagram |
|----------------|---|--|---------|
| Red blood cell | Transport of oxygen | Biconcave shape | |
| | | No nucleus | |
| | | Flexible | |
| | | Has haemoglobin | |
| Muscle cell | Contracts to get structures closer together | Long Many protein fibres in cytoplasm to shorten cell when energy available | |

| Cell | Function | Adaptation(s) | Diagram |
|----------------|-----------------------------------|--|---------|
| Ciliated cell | Move and push mucus | Tiny hairs called cilia | |
| Root hair cell | Absorb minerals and water | Elongated shape for more surface area | |
| Xylem vessel | Transport water and support plant | No cytoplasm so water passes freely | |
| | | No cross walls so cells connect to form tube | |
| | | Lignin makes it strong and waterproof | |
| Palisade cell | Photo-synthesizes | Regular shape so many can fit in a small space | |
| | | Many chloroplasts | |

respiratory system

1.3. Size of Specimens

$$\text{Magnification} = \frac{\text{size of drawing}}{\text{size of specimen}} = \frac{\text{image}}{\text{actual}} = \frac{I}{A}$$

2. Characteristics and Classification of Living Organisms

2.1. Characteristics of Living Organisms

- Movement:** action by an organism or part of an organism causing a change of position or place
- Respiration:** the chemical reactions that break down nutrient molecules in living cells to release energy
- Sensitivity:** ability to detect or sense changes in the environment (stimuli) and to make responses
- Growth:** permanent increase in size and dry mass by an increase in cell number or cell size or both
- Reproduction:** processes that make more of the same kind of organism
- Excretion:** removal from organisms of toxic materials, the waste products of metabolism (chemical reactions in cells)

including respiration) and substances in excess of requirements

- **Nutrition:** taking in of nutrients which are organic substances and mineral ions, containing raw materials or energy for growth and tissue repair, absorbing and assimilating them

2.2. Concept & Use of a Classification System

- Organisms can be classified into groups by the features that they share
- Classification systems aim to reflect evolutionary relationships (change of adaptive features of a population over time, as a result of natural selection)
- **Species:** organisms which can reproduce successfully
- Classification is traditionally based on studies of morphology and anatomy
- **Morphology:** the overall form and shape of their bodies e.g. wings or legs
- **Anatomy:** the detailed body structure determined by dissection
- **Binomial system:** a system of naming species in which the scientific name of an organism is made up of two parts showing the genus (starting with a capital letter) and species (starting with a lower-case letter), written in italics when printed (therefore underlined when written) e.g. *Homo sapiens*

KING PHILIP CAME OVER FOR GOOD SPAGHETTI

- Kingdom, Phylum, Class, Order, Family, Genus, Species
- Kingdom --> Species = Similarity increases
- DNA is the chemical from which chromosomes are made
- Each DNA molecule is made up of strings of smaller molecules containing four bases
- Biologists compare the sequences of the bases in the DNA of organisms from two different species
- The more similar the base sequence, the more closely related the species are to one another
- Organisms which share a more recent ancestor have base sequences in DNA that are more similar than those that share only a distant ancestor
- The sequences of bases in DNA and of amino acids in proteins are used as a more accurate means of classification (cladistics)

2.3. Kingdoms

- **Animal:** Multi-cellular ingestive heterotrophs (eat living organisms)
- **Plant:** Multi-cellular photosynthetic autotrophic (make their own food) organism with a cellulose cell wall.
- **Fungi:** Single celled or multi cellular heterotrophic organism with cell wall not made of cellulose, spread by

spreading of spores in moist/dark/warm environment, saprotrophs (feed off dead organisms) or parasites

- **Prokaryotes:** Single celled organism with no true nucleus
- **Protocist:** Single celled organism with a nucleus

2.4. Vertebrates

MR FAB

- **Mammals**
 - Fur/hair on skin
 - Can live on land and in water
 - 4 limbs
 - Lungs to breathe
 - Give birth to live young
- **Reptiles:**
 - Scales on skin
 - Usually 4 legs
 - Lungs to breathe
 - Hard eggs
- **Fish:**
 - Wet scales
 - External fertilization and soft eggs
 - Gills to breathe
- **Amphibians:**
 - Smooth, moist skin
 - External fertilization and soft eggs
 - Gills/lungs to breathe so can live on land and in water
 - 4 legs
- **Birds**
 - Feathers on body and scales on legs
 - Have 2 legs and 2 wings
 - Lungs to breathe
 - Hard eggs

2.5. Arthropods (Invertebrates with Legs)

CAMI

- **Crustaceans:** (e.g. crabs)
 - Have an exoskeleton
 - 1 pair of compound eyes
 - 2 body segment – cephalothorax and abdomen
 - More than four pairs of legs
 - 2 pairs of antennae sensitive to touch and chemicals
- **Arachnids:** (e.g. spiders)
 - 2 body segment – cephalothorax and abdomen
 - Four pairs of legs
 - Pair of chelicerae to hold prey
 - Two pedipalps for reproduction
 - Simple eyes
- **Myriapods:** (e.g. centipede)
 - Segmented body
 - Additional segments formed
 - One pair of antennae
 - 70+ pairs of legs – 1 or 2 pairs on each segment

- Fused head and thorax and segmented abdomen
- Simple eyes
- Insects: (e.g. bees)
 - 3 body segments – head, thorax and abdomen
 - 3 pairs of legs
 - 1 pair of antennae
 - 1 or 2 pairs of wings
 - Compound and simple eyes

2.6. Classifying Plants


- **Ferns:**
 - Do not produce flowers
 - They are plants with roots, stems and leaves
 - Have leaves called fronds
 - Reproduce by spores
- **Flowering plants:**
 - They are plants with roots, stems and leaves
 - Reproduce sexually by means of flowers and seeds
 - Seeds are produced inside the ovary in the flower

| Monocotyledons | Dicotyledons |
|--------------------|--------------------------|
| One cotyledon | Two cotyledons |
| Parallel veins | Veins netlike |
| Fibrous root | Taproot present |
| Floral parts in 3s | Floral parts in 4s or 5s |

2.7. Viruses and Bacteria

| | Virus | Bacteria |
|--------------------------|-------------------------------|------------------------------------|
| Covered by: | Protein coat | Cell wall |
| Cell membrane: | No | Yes |
| Cytoplasm: | No | Yes |
| Genetic material: | DNA or RNA – only a few genes | DNA – enough for several 100 genes |
| Living or not? | Non-living unless in host | Living |

- Two examples of virus structure



2.8. Dichotomous Keys

- **Dichotomous key:** uses visible features to classify organisms. It is which gives you a choice of two features and you follow the one that applies: each choice leads to another choice until the organism is narrowed down to its genus and finally species.

3. Movement in and Out of Cells

3.1. Diffusion

- This is the movement of molecules from a region of high concentration to a region of low concentration down the concentration gradient.
- This results in random movement of molecules until equilibrium is reached
- The diffusion of gases and solutes is important as without it, molecules which are needed for life, for example glucose and oxygen for respiration, would not be able to get to the places they are needed. Water is needed as a solvent
- Factors influencing faster diffusion:
 - Larger concentration gradient
 - Higher temperature
 - Larger surface area

3.2. Osmosis

- Movement of water molecules from a region of high-water potential to a region of low water potential, through a partially permeable membrane
- Conc. of solute outside cell = conc. inside cell → no change in size
- Conc. of solute outside cell > conc. inside cell → cell shrinks (Plasmolysis)
- Conc. of solute outside cell < conc. inside cell → cell swells (Turgid)

- In animals:
 - Increasing solute concentration inside of cell can cause cell to explode as a result of it having too much water, crenation.
- In plants:
 - Increasing solute concentration inside of cell causes cell to become turgid, vacuole fills up.
 - Decreasing solute concentration inside of cell causes cell to become flaccid, losing water and vacuole gets smaller. Cell body shrinks, pulling away from cell wall

3.3. Active Transport

- Movement of particles through a cell membrane, from a region of lower concentration to a region of higher concentration against a concentration gradient, using energy released during respiration
- Active transport is needed when an organism wants to optimize the amount of nutrients it can take up - ion uptake by root hairs and uptake of glucose by epithelial cells of villi.

4. Biological Molecules

- **Carbohydrates:** made from Carbon, Hydrogen and Oxygen (CHO)
- **Fats and oils:** made from Carbon, Hydrogen and Oxygen (CHO)
- **Proteins:** made from Carbon, Hydrogen, Oxygen, Nitrogen and sometimes Sulfur (CHON{S})

| Basic units (monomers) | Larger molecules (macromolecules) |
|--------------------------|-----------------------------------|
| Simple sugars | Starch and glycogen |
| Fatty acids and glycerol | Fats and oils |
| Amino acids | Proteins |

4.2. Chemical Tests

- **Starch:** Add few drops of iodine, +ve result = blue-black colour
- **Reducing sugars:** Add Benedict's reagent, then mixture is heated in water bath for 2 to 3 minutes. +ve result = brick-red precipitate, -ve result = remains blue
- **Proteins:** Add few drops of Biuret reagent, +ve result = mauve colour
- **Fats:** Emulsion test; ethanol is added to mixture, and this is poured into a test tube with an equal amount of distilled water, +ve result = milky-white emulsion
- **Vitamin C:** Decolourisation of DCPIP shows that a vitamin C is probably present.


4.3. Structure of a DNA

- Chromosomes are made of a molecule called DNA

- Each chromosome is a very long molecule of tightly coiled DNA
- Two strands coiled together to form a double helix
- Each strand contains chemicals called bases
- Cross-links between strands are formed by pairs of bases
- The bases always pair up in the same way:
 - A and T
 - C and G

5. Enzymes

- **Catalyst:** a substance that speeds up a chemical reaction and is not changed by the reaction
- **Enzymes:** proteins that function as biological catalysts
- Enzymes lowers amount of energy needed for reaction to take place
- Enzyme lowers the activation energy needed for reaction to take place
- Lock and key theory:



- **Substrate:** the molecule(s) before they are made to react
- **Product:** the molecule(s) that are made in a reaction
- Catabolic reaction: molecules are broken down
- Anabolic reaction: molecules are combined


5.2. Effect of Temperature on Enzymes

- Enzymes have an optimum temperature: the temperature at which they work best giving the fastest reaction $\approx 37^{\circ}\text{C}$ in animals
- When temperature increases, molecules move faster so collide with an enzyme in less time
- Having more energy makes them more likely to bind to active site.
- If temperature is too high, enzyme molecules vibrate too vigorously and enzyme is denatured; it loses its shape and will no longer bind with a substrate.
- When the temperature is too low there is not enough kinetic energy for the reaction so it reacts too slowly.

5.3. Effect of pH on Enzymes

- Enzymes are sensitive to pH
- Some enzymes work best in an acid and others in an alkaline
- Enzymes work best at their optimum pH
- If the pH is changed then the enzyme will denature and will no longer fit with substrate- no reaction takes place

5.4. Graphs for Changes in Enzyme Activity





5.5. Enzymes and their Uses

- **Seeds to germinate:** the enzymes turn insoluble food stores to soluble.
- **Biological washing powders:** enzymes are added to washing powders to help remove stains for example:
 - Lipase for lipids from fatty foods and greasy fingerprints
 - Protease for proteins from blood stains
- **Food industry:**
 - Isomerase converts glucose to fructose which is sweeter, so less is needed to give a sweet taste
 - Pectinase helps break down cell walls in fruit juice production so it increases yield, lowers viscosity and reduces cloudiness

6. Plant Nutrition

- **Photosynthesis:** process by which plants manufacture carbohydrates from raw materials using energy from light.




- The carbon dioxide diffuses through the open stomata of the leaf of a plant and water is taken up through roots.
- Chlorophyll is a dye, which traps light energy and converts it into chemical energy for the formation of carbohydrates and their subsequent storage.

6.2. Chlorophyll Is Necessary for Photosynthesis


- Take a potted plant with variegated (green and white) leaves.
- Destarch the plant by keeping it in complete darkness for about 48 hours.
- Expose the plant to the sunlight for a few days.
- Leaf boiled in water for 2 minutes to break down cell walls, denature enzymes and allow for easier penetration by ethanol.

- Warmed in ethanol until leaf is colourless to extract chlorophyll, which would mask observation
- Dipped in water briefly: to soften leaf
- Leaf is placed on a white tile and iodine is added. If starch is present, colour will be blue-black and if absent, it will remain orange




6.3. Light Is Necessary for Photosynthesis

- Destarch the plant by keeping it in darkness for 48hrs
- Place a stencil over part of a leaf
- Place the leaf in sunlight for 4-6 hours
- Remove the stencil and test for starch
- +ve result = parts which received light turn black
- -ve result = parts which didn't receive light remain yellow/brown



6.4. Carbon Dioxide is Necessary for Photosynthesis


- Take two destarched potted plants.
- Cover both the plants with bell jars and label them as A and B.
- Inside A, keep NaHCO_3 (sodium bicarbonate). It produces CO_2 .
- Inside B, keep NaOH (Sodium hydroxide). It absorbs CO_2 .
- Keep both the set-ups in the sunlight for at least 6 hours.
- Perform the starch test on both plants.



- The leaves of Plant A will turn black after the starch test
- The leaves of Plant B will remain orange/brown after starch test

6.5. Limiting Factors

- Limiting factor: something present in the environment in such short supply that it restricts life processes.




6.6. Glasshouse Systems

To increase the crop yield, farmers control the limiting factors:


- CO₂ enrichment: paraffin is burnt to increase CO₂ concentration by three times the original amount and doubling the yield
- Optimum temperature: thermostatically controlled heaters make the temperature right for enzymes to work
- Optimum light: light has a high intensity for more photosynthesis, the correct wavelengths (red and blue not green) and duration controls production of fruit

6.7. Leaf Structure




- Cuticle:** waxy layer that prevents water loss from top of the leaf
- Epidermis:** transparent cell that allows sunlight to pass through to the palisade cell
- Palisade:** found at the top of the cell and contains many chloroplasts which absorb sunlight.
- Spongy mesophyll layer:** irregularly shaped cells which create air spaces to allow gaseous exchange to take place; do not contain many chloroplasts
- Vascular Bundle:** made up of xylem and phloem
- Xylem:** vessel which transports water and dissolved minerals and has lignified walls made of cellulose
- Phloem:** vessel which transports nutrients

- Stomata:** little holes that open and close to allow gaseous exchange to take place. The stomata close to prevent water loss and open to let gases come in and out. When guard cells lose water, the stomata close (at night), while the stomata open when guard cells gain water & swell (during the day).




6.8. Xylem



- Unidirectional vessel which transports water and dissolved minerals.
- Walls are made out of waterproof lignin.
- Water moves up due to transpiration and osmosis

6.9. Phloem



- Bidirectional vessel
- Contains sieve elements which allow sugars to pass from one cell to next downwards
- Contains companion cells which provide energy for active transport of sugars all over plant.
- Translocation moves organic molecules (sugars, amino acids) from source to sink.
- Phloem vessels still have cross walls called sieve plates that contain pores.
- Companion cells actively load sucrose into the phloem.
- Water follows high solute in phloem by osmosis. A positive pressure potential develops moving mass of phloem sap forward.
- Phloem still contains small amount of cytoplasm along the walls but the organelle content is greatly reduced.

- Companion cells actively unload (ATP used) the organic molecules

6.10. Mineral Requirements

| Nitrogen | Magnesium |
|---|--|
| Needed for protein synthesis | Needed for chlorophyll synthesis |
| Deficiency: small plant, slow growth, top leaves pale, bottom leaves dead and roots slightly affected | Deficiency: plant lack chlorophyll, leaves yellow but normal roots |

- Nitrogen fertilizers:** provide nitrogen in the form of nitrate ions, nitrite ions or ammonium ions. But using fertilisers can lead to eutrophication, which is when the fertiliser is transported by rain and leaches into stagnant water e.g. pond or river

7. Human Nutrition

- Balanced Diet:** getting all the right nutrients in correct proportions
- Diet related to age/sex/activity:
 - Children Below 12:** Require more calcium
 - Teenagers:** Highest calorie Intake
 - Adults:** Balanced meal with less calories
 - Pregnant Women:** more iron, calcium and folic acid
 - Males:** Generally, require more energy

7.2. Malnutrition

- A condition caused by eating an unbalanced diet. Several forms:
 - Overnutrition:** balanced diet but eating too much of everything
 - Undernutrition:** having too little food
 - Eating foods in incorrect proportions

7.3. Effects of Malnutrition

- Starvation:** losing strength & finally dying because of lack of food
- Coronary heart disease:** eating too much fats which are rich in saturated fatty acids and cholesterol, may lead to heart attack
- Constipation:** lack of roughages in food causes constipation because roughages are indigestible and form bulks. Friction between bulks and walls of intestine stimulate the peristalsis
- Obesity:** Eating too much fats and carbohydrates leads to their storage in the body mainly in the forms of fats and causing an increase in body weight. This can cause; heart attack, stroke, joint pain, mobility impairment, high blood pressure


7.4. Uses

| Nutrient | Uses |
|---------------|--|
| Carbohydrates | Energy |
| Fats | Source of energy, building materials, energy store, insulation, buoyancy, making hormones |
| Proteins | Energy, building materials, enzymes, haemoglobin, structural material (muscle), hormones, antibodies |
| Vitamin C | Protect cells from ageing, production of fibres |
| Vitamin D | Absorption of calcium |
| Calcium | Development and maintenance of strong bones and teeth |
| Iron | Making haemoglobin |
| Fiber | Provides bulk for faeces, helps peristalsis |
| Water | Chemical reactions, solvent for transport |

7.5. Deficiencies

- Vitamin C:** *Scurvy*; loss of teeth, pale skin & sunken eyes
- Vitamin D:** *Rickets*; weak bones and teeth
- Calcium:** *Rickets*; weak bones and teeth, also poor clotting of blood, spasms
- Iron:** *Anaemia*: Fatigue (less iron → less haemoglobin → less oxygen transported → less respiration → less energy)

7.6. Human Alimentary Canal



- Ingestion:** taking substances (e.g. food, drink) into the body through the mouth.
- Egestion:** passing out of food that has not been digested, as faeces, through the anus.
- Digestion:** the break-down of large, insoluble food molecules into small, water soluble molecules using mechanical and chemical processes
- Mouth:** contains teeth used for mechanical digestion, area where food is mixed with salivary amylase & where ingestion takes place
- Salivary glands:** produce saliva which contains amylase and helps food slide down oesophagus
- Oesophagus:** tube-shaped organ which uses peristalsis to transport food from mouth to stomach
- Stomach:** has sphincters to control movement into and also has pepsin (a protease) to break down proteins into





peptides, it also kills bacteria with hydrochloric acid. They also have elastic walls.

- **Small intestine:** tube shaped organ composed of two parts the:
 - **Duodenum:** fats are emulsified by bile, and digested by pancreatic lipase to form fatty acids and glycerol. Pancreatic amylase and trypsin (a protease) break down starch and peptides into maltose and amino acids
 - **Ileum:** Maltase breaks down maltose to glucose. This is where absorption takes place; adapted by having villi and microvilli.
- **Pancreas:** produces pancreatic juice which contains amylase, trypsin and lipase and hydrogencarbonate.
- **Liver:** produces bile, stores glucose as glycogen, interconverting them to keep glucose concentration constant. Also carries out interconversion of amino acids (transamination), deamination and removal of old red blood cells and storage of their iron. Also, site of breakdown of alcohol and other toxins.
- **Gall bladder:** stores bile from liver
- **Bile:** produced by liver and stored in gall bladder, its role is to emulsify fats, to increase surface area for the action of enzymes.
- **Large intestine:** tube shaped organ composed of two parts:
 - **Colon:** organ for absorption of minerals and vitamins, and reabsorbing water from waste to maintain body's water levels
 - **Rectum:** where faeces are temporarily stored
- **Anus:** ring of muscle which controls when faeces is released.

7.7. Diarrhoea


- **Diarrhoea:** when not enough water is absorbed from the faeces
- To cure this is to give oral rehydration therapy
- One of these infectious by a bacterium causing the diseases cholera (spreads rapidly)
- The cholera bacterium produces a toxin that causes secretion of chloride ions into the small intestine, causing osmotic movement of water into the gut, causing diarrhoea, dehydration and loss of salts from the blood

7.8. Teeth

| Incisor | Canine | Premolar | Molar |
|---|---|---|---|
|  |  |  |  |
| Incisors | Canines | Premolars | Molars |

| Incisor | Canine | Premolar | Molar |
|---|---------------------------------------|--------------------------------|---------------------------------------|
| Rectangular shape, sharp for cutting and biting | Sharp-pointed for holding and cutting | Blunt for chewing and crushing | Blunt chewing and crushing. Two roots |


7.9. Structure of a Tooth



- **Enamel:** the strongest tissue in the body made from calcium salts
- **Cement:** helps to anchor tooth
- **Pulp cavity:** contains tooth-producing cells, blood vessels, and nerve endings which detect pain.
- **Dentine:** calcium salts deposited on a framework of collagen fibres

Neck: in between crown and root, it is the gums

7.10. Tooth Decay



7.11. Prevention


- Eating food with low sugar content
- Regular and effective teeth brushing to remove plaque
- Finishing a meal with a crisp vegetable and a glass of water

7.12. Chemical Digestion

- Where enzymes are used to break down large insoluble substances such as proteins into smaller soluble substances like amino acids so that they can be absorbed.
- **Amylase:** breaks down starch into maltose, it is produced in the pancreas (but also in the salivary gland)
- **Protease:** breaks down proteins to peptides (done by pepsin) then into amino acids (done by trypsin). Pepsin comes from the stomach and trypsin comes from the pancreas.
- **Lipase:** breaks down lipids into fatty acids and glycerol, produced by the pancreas.

- **Hydrochloric acid in gastric juice:**
 - Denaturing enzymes in harmful microorganisms in food
 - Giving the optimum pH for pepsin activity

7.13. Absorption






- Movement of digested food molecules through wall of the intestine into the blood or lymph.
- The small intestine is the region for absorption of digested food.
- The small intestine is folded into many villi which increase the surface area for absorption. One villus will have tiny folds on the cells on its outside called microvilli.
- More surface area means more absorption can happen
- **Capillary:** transports glucose and amino acids
- **Vein:** delivers absorbed products to liver via hepatic portal vein.
- **Gland:** produces enzymes
- **Lacteal:** absorbs fatty acid and glycerol
- **Epithelium:** only one cell thick for faster transport. The cells of the epithelium are folded to form microvilli.
- Small intestine and colon absorb water
 - The small intestine absorbs 5–10 dm³ per day
 - The colon absorbs 0.3–0.5 dm³ per day


8. Transport in Plants

8.1. Xylem and Phloem

- Functions of xylem and phloem
 - To transport substances from source, where they are taken in or made, to the sinks, where they are used
 - To support the stem




8.2. Root Hair Cell



- **Function:** to absorb water and minerals from the soil
- They have an elongated shape for more surface area which increases the rate of absorption of water by osmosis and ions by active transport

8.3. Pathway Taken by Water



- Water enters root hair cell from moist soil via osmosis because water potential is higher in soil than in cytoplasm.
- Water passes through the cortex cells by osmosis but mostly by "suction".
- Water and minerals are forced to cross the endodermis.
- Water enters xylem then leaves when it gets to mesophyll cells

8.4. Transpiration

- **Transpiration:** evaporation of water at surfaces of the mesophyll cells followed by loss of water vapour from plant leaves, through stomata.
- Water leaves mesophyll cells, into air spaces created by irregular shape of spongy mesophyll cells, then diffuses out of the stomata.
- **Wilting:** occurs if water loss is greater than water uptake – cells become flaccid, tissues become limp and plant no longer supported

8.5. Uptake of Water

- Caused by water loss in leaves which lowers its water potential
- Water moves from xylem to leaf tissues via osmosis
- Water moves up the stem in the xylem due to tension (because of cohesion of water molecules to each other) caused by water loss from the leaves
- Ends with the gain of water through roots
- This upward flow of water is called the transpiration stream

8.6. Factors Affecting Rate of Transpiration

- **Temperature:** higher temperatures increase water-holding capacity of air and increases transpiration rate
- **Humidity:** low humidity increases water potential gradient between leaf and atmosphere ∴ increasing transpiration rate

- **Light intensity:** high light intensity causes stomata to open (to allow more photosynthesis) which causes increase in transpiration

8.7. Translocation

- Movement of sucrose and amino acids in phloem; from regions of production (sources) to regions of storage or to regions of utilization in respiration or growth (sinks).
- Translocation in different seasons:
 - Spring: sucrose transported from stores in roots to leaves
 - Summer & early autumn: sucrose goes from photosynthesizing leaves to root stores,


9. Transport in Animals

- **Circulatory system:** system of tubes (veins, capillaries, arteries) with a pump (heart) and valves (in heart and veins) to ensure one-way flow of blood.

9.2. Transport Systems

- **Single circulation system (fish):**
 - Two heart chambers
 - Blood absorbs oxygen in gills
 - Released in body cells then back to heart
- **Double circulation system:**
 - Four heart chambers
 - Blood passes through heart twice
 - Oxygenated in lungs, to heart, to body, back to heart
 - **Advantage:** delivers greater blood flow rate to tissues around the body as the heart pumps the oxygenated blood to it from the lungs


9.3. The Heart



- **Right atrium:** collect deoxygenated blood & pump it to right ventricle
- **Right ventricle:** pumps deoxygenated blood to lungs
- **Pulmonary artery:** carries deoxygenated blood from right ventricle to lungs
- **Septum:** separates left and right sides of the heart
- **Pulmonary vein:** carry oxygenated blood from lungs to left atrium

- **Left atrium:** collect oxygenated blood and pump it to left ventricle
- **Left ventricle:** pumps oxygenated blood to the body via the aorta
- **Aorta:** carries oxygenated blood from left ventricle to rest of body
- **Tricuspid and bicuspid valves:** prevent backflow of blood into the atria when ventricles contract (atria ventricular valves)
- **Pulmonary and aortic valves:** prevent backflow of blood from the arteries into the ventricles (semi-lunar valves)

9.4. Cardiac Cycle



- Physical activity makes the heart beat more quickly and more deeply, for an increased circulation of blood so that more oxygen and glucose can get to the muscles.

9.5. ECG Trace

- The electrical activity of the heart can be monitored by the electrocardiogram, pulse rate and listening to the sounds of the valves closing
- Physical activity makes the heart beat more quickly and more deeply, for an increased circulation of blood so that more oxygen and glucose can get to the muscle

9.6. Coronary Heart Disease

- Coronary artery becomes blocked, interrupting the supply of blood to the heart muscle.
- The heart muscle cells are deprived of oxygen & glucose, and poisonous wastes such as lactic acid build up.
- Part of the heart muscle stops contracting, causing a heart attack
- Caused by stress, smoking, poor diet, poor lifestyle & genetically
- Can be prevented by not smoking, avoiding fatty food and exercising regularly

- Treated by aspirin and surgery (stents, angioplasty and by-pass)

9.7. Arteries, Veins and Capillaries


| Vessel | Function | Structure |
|-----------|---|---|
| Artery | Transport high pressure blood away from heart | Elastic walls expand and relax as blood is forced out; causes pulse |
| | | Thick walls withstand high pressure |
| | | Rings of muscle narrow or widen artery to control blood flow. |
| Vein | Transport low pressure blood to the heart | Valves prevent backflow of blood. |
| | | Blood is at low pressure, but nearby muscles squeeze veins and help push blood to the heart |
| | | Large diameter and thin walls reduce resistance to flow of blood |
| Capillary | Allow substances to diffuse into cells | One cell thick wall for easy diffusion |
| | | Highly branched; large surface area |
| | | Capillary beds constantly supplied with fresh blood, so diffusion occurs |

9.8. Lymphatic System

- Circulation of body fluids, and the production of lymphocytes.
- Lymph node contains many lymphocytes which filter lymph.
- Tissue fluid: made when plasma is squeezed out of capillaries.
- Substances diffuse between cells and tissue fluid.
- Lymph vessels collect lymph and return it to the blood.
- Tissue fluid returns to the capillaries by osmosis.

9.9. Blood

- Red blood cells: haemoglobin and oxygen transport
- White blood cells: phagocytosis and antibody formation
- Platelets: causing clotting
- Plasma: transport of blood cells, ions, soluble nutrients, hormones, carbon dioxide, urea and plasma proteins



9.10. Immune system

| Phagocyte | Lymphocyte |
|---|---|
| Phagocyte has lobed nucleus and vesicles containing digestive enzymes. | Lymphocytes are found in blood and in lymph nodes |
| Phagocytosis: engulf pathogen, vesicles fuse with vacuole, enzymes digest bacteria. | Large nucleus and they produce antibodies, |
| Antigen: protein/ carbohydrate on surface of pathogen which provokes immune system | Antibodies: Y-shaped protein, bind to label pathogens. |
| | Then either destroyed by being ingested by phagocytes, or the antibodies may do it. |

9.11. Blood Clotting

- Reduces blood loss and keeps pathogens out
- Fibrinogen (inactive) turns to fibrin (activated), and forms a mesh to trap red blood cells, which eventually dries to form a scab.

10. Diseases and Immunity

10.1. Pathogens

- **Pathogen:** a disease-causing organism.
- **Transmissible disease:** a disease in which the pathogen can be passed from one host to another.
- The pathogen for a transmissible disease may be transmitted either:
 - Direct contact e.g. through blood, body fluids
 - Indirectly e.g. from contaminated surfaces/food, from animals, from air

10.2. Body Defences

- The human body has many natural defences against pathogens.
- **Mechanical barriers:**
 - Nostrils contain hairs that help trap dust
 - Skin has a thick outer layer of dead cells
- **Chemical barriers:**
 - Sticky mucus which can trap pathogens

- In the stomach, hydrochloric acid is secreted which kills many of the bacteria in food
- **Cells:** Pathogens that manage to get through all these defences are usually destroyed by white blood cells:
 - Some of these cells take in and digest the pathogens by phagocytosis
 - Others produce antibodies that incapacitate or kill the pathogen
 - Vaccination against disease helps antibodies to produce very quickly

10.3. The Immune System


- An antibody is a protein molecule which fits into another molecule
- Pathogen molecules are called **antigens**.
- To destroy a pathogen, antibody molecules must be made which are exactly the right shape to fit into molecules (antigens) on the outside of the pathogen.
- Antibodies lock onto antigens leading to destruction of pathogen / marking of pathogens for destruction by phagocytes
- If a pathogen enters the body, it meets a large number of lymphocytes. One of these will recognize the pathogen and divide rapidly by mitosis
- These lymphocytes then secrete antibody, destroying the pathogens
- **Active immunity:** defence against a pathogen by antibody production in the body.
- Active immunity is gained after an infection by a pathogen or by vaccination.
- **Vaccines** immunise children against diseases caused by pathogens
- **Process of vaccination:**
 - Harmless pathogen given which has antigens
 - Antigens trigger an immune response by lymphocytes which produce antibodies
 - Memory cells are produced that give long-term immunity
- In controlling the spread of disease, it is important to consider hygienic food preparation, good personal hygiene, waste disposal and sewage treatment
- **Passive immunity:** short-term defence against a pathogen by antibodies acquired from another individual, e.g. mother to infant
- Memory cells are not produced in passive immunity
- Babies get passive immunity by breast feeding.
 - Breast milk contains antibodies from the mother which are passed on to her baby.
 - Useful because a young baby's immune system is not well developed and so the mother's antibodies can protect it against any diseases to which she is immune for the first few months of its life
- Some diseases are caused by the immune system targeting and destroying body cells (Type 1 diabetes)

11. Gas Exchange in Humans

11.1. Gas Exchange

| Property of surface | Reason |
|-----------------------|--|
| Thin (one cell thick) | Short distance to diffuse |
| Large surface area | Many molecules can diffuse at once |
| Moist | Cells die if not kept moist |
| Well ventilated | Concentration gradients for oxygen and carbon dioxide are kept up by regular fresh supplies of air |
| Close to blood supply | Gases can be carried to/from the cells that need/produce them |

11.2. Structure



- **Cartilage (in trachea):** prevents the trachea from collapsing during absence of air and also to protect it
- **Ribs:** to protect vital organs and blood vessels and expands and contracts (and efficient breathing)
- **Intercostal (internal & external) muscles:** situated between the ribs that create and move the chest wall
- **Diaphragm:** produces volume and pressure changes in the thorax leading to the ventilation of the lungs

| | Inspired Air | Expired Air |
|----------------|-------------------|------------------------|
| Oxygen | 21% | 16% |
| Carbon Dioxide | 0.04% | 4% |
| Nitrogen | 78% | 78% |
| Water | Varies to climate | Saturated water vapour |
| Temperature | Varies to climate | Always warm |

- **Test for CO₂:** Blow CO₂ through limewater. +ve result = turn cloudy


11.3. Effect of Physical Activity on Breathing

- Physical activity increases the breathing rate – more breaths per minute, and the tidal volume – more air per breath

- This is measured with a spirometer to produce a spirogram.
- During exercise, tissues respire at a higher rate, the change in breathing volume and rate helps to keep CO₂ concentration and pH at safe levels.

11.4. Breathing

| Breathing In | Breathing Out |
|---|---|
| External intercostal muscles contract – pulls rib cage upwards and outwards | External intercostal muscles relax – rib cage falls downwards and inwards |
| Diaphragm muscles contract – diaphragm moves downwards | Diaphragm muscles relax – returns to dome shape |
| Lung volume increases – and pressure falls | Lung volume decreases – and pressure increases |
| Air rushes in to equalise pressure | Air is forced out |





- **Internal intercostal muscles:** are used in coughing and sneezing.
- **Mucus & cilia:** goblet cells produce sticky mucus to trap and eliminate particulate matter and microorganisms.
- **Ciliated cells** have cilia: little hairs which sweep back and forward in a coordinated way to brush mucus up the lungs into the mouth

12. Respiration

- Chemical reactions that break down nutrient molecules in living cells to release energy.
- Uses of energy in the body of humans: muscle contraction, protein synthesis, cell division, active transport, growth, the passage of nerve impulses and the maintenance of a constant body temperature.
- Respiration involves the action of enzymes in cells

12.2. Aerobic Respiration

- Release of a relatively large amount of energy in cells by the breakdown of food substances in the presence of oxygen.



12.3. Anaerobic Respiration



- Release of a relatively small amount of energy by the breakdown of food substances in the absence of oxygen.
- **In muscles:**
 - $\text{Glucose} \rightarrow \text{Lactic Acid}$
 - $C_6H_{12}O_6 \rightarrow 2C_3H_6O_3$
- **In yeast** (single-cell fungi):
 - $\text{Glucose} \rightarrow \text{Ethanol} + \text{Carbon Dioxide}$
 - $C_6H_{12}O_6 \rightarrow 2C_2H_5OH + 2CO_2$
- **Disadvantages of anaerobic respiration:**
 - Only produces 1/20 of the energy per glucose molecule that aerobic respiration would
 - Produces poisonous lactic acid
- **Lactic acid:**
 - Transported in blood to heart, liver and kidneys, which oxidize it
 - The heart, liver and kidneys need extra oxygen to do this which causes you to continue breathing heavily after exercise.
 - The extra oxygen is called the oxygen debt.

13. Excretion in Humans

- **Excretion:** the removal from organisms of toxic materials, the waste products of metabolism (chemical reactions in cells including respiration) and substances in excess of requirements.
- Substances should include carbon dioxide, urea and salts.


13.2. Function of Kidney

- Removal of urea and excess water and the re-absorption of glucose and some salts
- Urea is formed in the liver from excess amino acids
- Alcohol, drugs & hormones are broken down in the liver



- **Cortex:** contains Bowman's capsules and coiled tubules
- **Ureter:** carries urine from kidney to bladder
- **Medulla:** contains loops of Henlé and collecting ducts
- **Loop of Henlé:** selectively absorbs water/solutes
- **Collecting ducts:** reabsorbs water into blood and store wastes until they are passed into ureter
- **Urethra:** carried urine from bladder to the outside.
- **Bladder:** stores urine
- **Renal capsule:** filters from blood: water, glucose, urea and salts.
- **Tubule:** (yellow) reabsorbs 100% of glucose, most of the water and some salts back into the blood (red), leading to concentration of urea in the urine as well as loss of excess water and salts into the tubule.
- **Renal artery:** brings wastes and water from blood
- **Renal vein:** reabsorbs water and useful molecules and leaves wastes behind

13.3. Structure of the Kidney



1. **Ultrafiltration:** blood from renal artery enters the glomerulus. Water, urea, salts and glucose are forced into the Bowman's capsule. Blood cells and large proteins cannot pass through.
2. **Selective reabsorption:** in the proximal tubule two thirds of the salt and water and all the glucose moves out of the nephron, by active transport. These substances are reabsorbed back into the blood capillary.
3. **Loop of Henlé:** this part of the loop of Henlé is permeable to water but not salt. Water is drawn out of the filtrate in the nephron by osmosis because of the low water potential of the medulla tissue fluid.
4. **Loop of Henlé:** this part is permeable to salt but not water. The loss of water from the filtrate in the

previous stage increases the salt concentration. Some salt passively diffuses out of the loop here.

- 5. **Collecting duct:** the remaining substances move through the second coiled tubule (distal tubule), into the collecting duct. The permeability of this part of the nephron to water is controlled

13.4. Dialysis

- When a kidney machine takes a patient's blood and cleans it, then returns the blood to circulation.
- This is how it works:
 - Blood enters machine from patient
 - The pump passes the blood through the dialysis tubing which is semi-permeable therefore acting as a filter
 - The surrounding liquid contains some salts, glucose but no urea so waste materials pass from blood by diffusion;
 - The 'cleaned' blood returns to patient's circulation/body

| Dialysis | Transplant |
|---|--|
| More expensive in the long run | Less expensive in the long run |
| Very disruptive (three 6-8 hrs sessions per week) | Not very disruptive (only have to take medication) |
| Do not need to find kidney | Need a kidney |
| Need a machine & must live near one | Can go anywhere, anytime |
| | Risk of rejection |

14. Coordination and Response


14.1. Nervous Control in Humans

- The nervous system consists of two parts:
 - **Central nervous system (CNS)** consisting of the brain and spinal cord, which are the areas of coordination
 - **Peripheral nervous system (PNS)** made up of nerves and neurones, which coordinate and regulate bodily functions.
- Involuntary actions: not under conscious control e.g. reflex action
- Voluntary actions: are done if we decide to carry them out


14.2. Types of Neurons

- **Nerve impulse:** an electrical signal that passes along nerve cells called neurones


Motor Neurone:



Sensory Neurone:




Relay (connector) neurone:



14.3. Reflex arc

- A reflex action is an involuntary, quick action to respond to a stimulus, in order to protect the body from danger
- E.g. quickly removing your hand from hot metal surface
- They involve three neurones: a sensory neurone, relay neurone and motor neurone.
- The gap between neurones is called a synapse.
- **How the reflex arc works:**
 - A stimulus affects a receptor (cell or organ that converts a stimulus into an electrical impulse)
 - A sensory neurone carries impulse from the receptor to the CNS
 - Connector/relay neurone carries impulse slowly (because it has no myelin sheath) across the spinal chord
 - Motor neurone carries impulse from the CNS to the effector
 - Effector (either a muscle or a gland) carries out the response



- **Reflex action:** means of automatically and rapidly integrating and coordinating stimuli with the responses of effectors (muscles and glands)

14.4. Synapses


- **Synapse:** a junction between two neurones, consisting of a gap across which impulses pass by diffusion of a neurotransmitter
- **Synaptic cleft:** small gap between each pair of neurones
- Inside the neurones axon, there are 100s of tiny vacuoles (vessicles each contain a chemical called neurotransmitter)
- When an impulse arrives, the vessicles move to the cell membrane and empty their content into the synaptic cleft
- The neurotransmitter quickly diffuses across the tiny gap and attaches to receptor molecules in the cell membrane of the relay neurone
- This can happen because the shape of the neurotransmitter molecules is complimentary to the shape of the receptor molecule
- Many drugs e.g. heroin act upon synapses

14.5. Antagonistic Muscle

- A muscle that opposes the action of another; e.g. biceps and triceps are antagonistic muscles or circular and radial muscles in the eye
- **Agonist:** a muscle that contracts while another relaxes; e.g. when bending the elbow, the biceps are the agonist
- **Antagonist:** a muscle that relaxes while another contracts; e.g. when bending the elbow, the triceps are the antagonist
- **Sense organ:** groups of receptor cells responding to specific stimuli: light, sound, touch, temperature and chemicals.

14.6. The Eye


- The sense organ responsible for sight



- **Cornea:** refracts light
- **Iris:** controls how much light enters pupil
- **Lens:** focuses light onto retina
- **Retina:** contains light receptors, some sensitive to light of different colours
- **Optic nerve:** carries impulses to the brain

14.7. Accomodation

- Adjusting for near and distant objects.



| Near Object | Distant Object |
|----------------------------|----------------------------|
| Ciliary muscles contract | Ciliary muscles relax |
| Ligaments relax | Ligaments are tight |
| Lens becomes short and fat | Lens becomes long and thin |

14.8. Pupil Reflex



- Adjusting for high and low light intensity

| Low Light Intensity | High Light Intensity |
|---|---|
| Radial muscles (straight lines) contract and become shorter to pull the pupil (black dot) making it wider, to let more light enter, to form a clear image on retina | Circular muscles (circular lines) contract and become shorter to reduce the size of the pupil to protect retina from bleaching. |

14.9. Rods and Cones

| Rods | Cones |
|---|--|
| Provide low detail, black & white images, good for seeing in low intensity light (at night). | Provide detailed, coloured images; they work in high light intensity. |
| Packed most tightly around edge of retina so you can see things most clearly when not looking directly at them. | Most tightly packed at centre of retina, so objects are seen most clearly when being directly looked at. |

- **Fovea:**
 - Part of the retina where the receptor cells are pushed most closely together
 - Where light is focused when you look straight at an object

14.10. Hormones

- A chemical substance, produced by a gland, carried by the blood, which alters the activity of one or more specific target organs and is then destroyed by the liver.

14.11. Adrenaline

- A hormone secreted by the adrenal gland.
- It increases pulse rate, makes the glycogen in muscles is converted to glucose and released into blood, makes you breathe deeper and more rapidly, airways become wider, and makes skin become pale as blood is diverted away.
- Increases blood glucose concentration for respiration.
- Adrenaline is secreted for example: while bungee jumping or riding a rollercoaster

| Gland | Hormone | Function |
|---------------|--------------|---|
| Adrenal gland | Adrenaline | Prepares body for vigorous action |
| Pancreas | Insulin | Reduces conc. of glucose in blood |
| Testis | Testosterone | Causes development of male sexual characteristics |
| Ovary | Oestrogen | Causes development of female sexual characteristics |

14.12. Nervous and Hormonal Systems

| Comparison | Nervous system | Endocrine system |
|----------------------|--|---|
| Speed of action | Very rapid | Can be slow |
| Nature of message | Electrical impulses, travelling along nerves | Chemical messenger (hormones) travelling in bloodstream |
| Duration of response | Usually within seconds | May take years (puberty) |

| Comparison | Nervous system | Endocrine system |
|-------------------------------|--|--------------------------------------|
| Area of response | Localized response (only one area usually) | Widespread response (in many organs) |
| Example of process controlled | Reflexes such as blinking | Development of reproductive system |

- Hormones are used in food production, for example oestrogen is used to boost growth rate of chickens.
- Advantage: chickens grow quickly meaning more profit.
- Disadvantages: this may cause human males to develop feminine characteristics, and it is unnatural.

14.13. Homeostasis


- The maintenance of a constant internal environment.
- Homeostasis is the control of internal conditions within set limits

14.14. Negative Feedback

- Feedback controls the production of hormones – the hormones regulate their own production.
- A negative feedback control is when the change in hormone level acts as a signal to cancel out that change, so when blood hormone level is low, hormone production is stimulated, when it is high, it is inhibited.

14.15. Glucoregulation


- Blood glucose levels are monitored and controlled by the pancreas
- The pancreas produces and releases different hormones depending on the blood glucose level
- Insulin is released when blood glucose levels are high – the liver stores excess glucose as glycogen
- Glucagon is released when blood glucose levels are low – the liver converts stored glycogen into glucose and releases it into the blood



- When the control of blood glucose does not work, a person is said to have diabetes
- **Type 1 diabetes** is caused by the death of the cells that secrete insulin
 - **Symptom:** hyperglycaemia (feel unwell, dry mouth, blurred vision and feel thirsty) or hypoglycaemia (tired, show confusion and irrational behaviour)


- **Treatment:** eating little and often and avoiding large amount of carbohydrates, injecting insulin to reduce blood glucose concentration

14.16. Thermoregulation



Constant body temperature is maintained by:

- **Insulation:** provided by fatty tissue retains heat. Hairs become erect to trap warm air by contracting erector muscles and vice versa.
- **Vasodilatation:** when it is hot, arterioles, which supply blood to the skin-surface capillaries, dilate (become wider) to allow more blood near to skin surface to increase heat loss (face redder)
- **Vasoconstriction:** when it is cold, arterioles, which supply blood to the skin-surface capillaries, constrict (become smaller) to allow less blood near to skin surface to decrease heat loss
- **Sweating:** the water evaporates giving a cooling effect
- **Skin receptors:** sense heat and sensory neurons send impulses to the hypothalamus
- **Shivering:** muscular activity generates heat
- **Thermoregulatory centre:** in the hypothalamus, it controls the use of corrective mechanisms (e.g. sweating and shivering).



14.17. Homeostatic Organs

- Cells: change composition of blood as they remove nutrients and O₂ and add wastes and CO₂
- Heart: keeps blood pressure constant to deliver oxygen and nutrients around body
- Skin: to maintain heat exchange with external environment
- Kidneys: regulate water and salt levels (osmoregulation) and the removal of wastes like urea (excretion)

- Lungs: regulate gas exchange
- Intestines: supply soluble nutrients and water to blood
- Liver: regulates blood solutes and removes toxins

14.18. Tropic Responses

- **Auxin:**
 - Plant hormones or growth substances
 - Controls tropisms
 - It is produced by cells at the tip of roots and shoots of plants
- **Gravitropism:** a response in which a plant grows towards (positive) or away (negative) from gravity.
- **Auxins' role in gravitropism:**
 - Tend to settle at the bottom end of the root.
 - However, this does not make the cells of the tip of the root grow longer; auxins prevent cells at bottom tip of root from growing, making cells at top of root grow faster.
 - When cells of top of the root grow faster, they push root deeper into soil and root gets longer.
 - The root grows in direction of the gravitational pull.
- **Phototropism:** a response in which a plant grows towards (positive) or away (negative) from the direction from which light is coming.
- **Auxins' role in phototropism:**
 - If sun shines on right side of a plant's shoot, auxins will accumulate on dark opposite left side.
 - Auxins accumulating makes cells on left side grow faster than cells on right side.
 - When left side of shoot starts growing faster than right side, shoot will start to bend to right side towards sunlight.
- **Hormones can be used as weed killers:** spraying with high concentrations of hormone (2,4-D) upsets normal growth patterns. It affects different species differently so might only kill one species not the other (this is good).

15. Drugs

- **Drugs:** Any substance taken into the body that modifies or affects chemical reactions in the body.

15.2. Antibiotics

- Antibiotics work by stopping a metabolic practice performed by the bacteria you are trying to get rid of, but not performed by human cells.
- Some bacteria are resistant to antibiotics which reduces the effectiveness of antibiotics
- Development of resistant bacteria such as MRSA can be minimised by limiting use of antibiotics only when essential and ensuring treatment is completed
- Antibiotics don't work on viruses because they are not really living and they make the host cell perform the tasks for them.

15.3. Heroin

Effects of the abuse of heroin: a powerful depressant

- Problems of addiction
- Severe withdrawal symptoms (vomiting, restlessness)
- Malnourishment as drug depresses appetite
- Financial problems – stealing, loss of job
- Infection from sharing needles e.g. HIV/AIDS
- Heroin affects the nervous system by its effect on the function of synapses

15.4. Alcohol

Effects of excessive consumption of alcohol –a depressant:

- Causes coronary heart diseases
- Reduced self-control
- Depression
- Effect on reaction times
- Damage to liver – cirrhosis

15.5. Smoking

Some effects of tobacco smoke:

- Drying effect and heat irritate lungs – destroys cilia
- Nicotine is addictive, it is also a stimulant, it increases pulse rate and narrows blood vessels which can cause damage
- Tobacco smoking can cause chronic obstructive pulmonary disease (COPD), lung cancer and coronary heart disease
- Tar causes cancer, and is an irritant so causes coughing.
- There are other irritants in tobacco smoke including: smoke particles, ammonia, and sulphur dioxide
- Emphysema: walls between alveoli break making large sacs, reducing surface area massively and making you breathless after a couple of steps
- Loss of limbs due to poor circulation, CHD and lower sperm-count
- Carbon monoxide irreversibly bonds with haemoglobin which can lead to oxygen starvation
- Cancer of the stomach, pancreas and bladder etc.
- Liver is the site of breakdown of alcohol and other toxins

15.6. Hormones and Sports

- **Hormones:** can be used to improve sporting performance
- Testosterone
 - Improved hand-eye coordination
 - Improved body-fat composition
 - Increased muscle mass
- Anabolic Androgenic Steroids
 - Affects limbic system
 - Mood swings
 - Impaired judgement

- High blood pressure
- Kidney failure
- Increased risks of prostate cancer (male)
- Inconsistencies of menstrual cycle (female)
- Changes in blood cholesterol

16. Reproduction

16.1. Asexual Reproduction

- The process resulting in the production of genetically identical offspring from one parent.
- **Bacteria:**
 - Reproduce by binary fission, each bacterium divides into two.
 - The generation time is the time taken for a cell to divide into 2.
- **Fungi:**
 - Single-celled yeast reproduces by binary fission.
 - All other fungi produce via spores.
 - When the sporangium bursts it spreads the spores.
 - Spores land and grow mycelium (roots) for example mushrooms
- **Potatoes:**
 - The shoot from a potato goes back underground and the stem swells to form a new genetically identical potato.
 - The swollen stem acts as a storage organ.

| Advantages | Disadvantages |
|--|---------------------------------|
| Fast: no need to find mate, fertilise etc. | No variation |
| Good characteristics are kept | Harmful genes transferred |
| | Overcrowding- fighting for food |


16.2. Sexual Reproduction

- **Sexual reproduction:** process involving the fusion of the nuclei of two gametes (sex cells) to form a zygote and the production of offspring that are genetically different from each other
- **Fertilisation:** the fusion of gamete nuclei
- Nuclei of gametes are haploid and that the nucleus of a zygote is diploid


| Advantages | Disadvantages |
|--|---|
| Produces genetically different offspring | Takes lots of time and energy |
| don't all die from change in the environment | Good characteristics can be lost |
| | Energy on improving appearances or pollen volume for pollination (plants) |

16.3. Sexual Reproduction in Plants

- Insect pollinated, dicotyledonous flowering plant: foxglove



Wind pollinated flower structure: grass



16.4. Functions

- **Sepal:** protect the flower bud.
- **Petal:** brightly coloured and scented and may have nectarines which are all used to attract insects, petals in wind pollinated flowers are tiny, and used for pushing the bracts (leaf-like structures) apart to expose stamens and stigma
- **Anther:** has pollen sacs with pollen grains which contain the male nucleus (male gamete).
- **Stigma:** platform on which pollen grains land
- **Ovary:** hollow chamber, ovules grow from the walls.


16.5. Pollination

- **Pollination:** transfer of pollen grains from the male part of the plant (anther of stamen) to the female part of the plant (stigma).
- **Agents of pollination:** insects, birds, mammals, water and wind

| Insect Pollinated | Wind Pollinated |
|---|---------------------------|
| Large colourful petals – attract | Dull petals |
| Sweetly scented | No scent |
| Nectaries | No nectaries |
| Moderate amount of pollen | Huge amount of pollen |
| Pollen is spiky/sticky | Pollen round and smooth |
| Anther & stigma inside flower | Anther & stigma hangs out |
| Stick stigma | Stigma hairy |
| Flowers have stripes which act as guide-lines for insects | |





- **Pollen tube:** pollen grain lands on stigma and creates a tunnel down the style, through the micropyle, to the ovules.

- Structure of non-endospermic seed:



Plumule + Radicle = embryo

- **Formation of a seed:** the zygote divides many times by mitosis to form an embryo. The cotyledon is the food store. The testa stops drying out of embryo.
- Wind and animal dispersal are used by plants to colonise new areas; done because new areas have less competition for light, space and nutrients, so seeds are more likely to develop.


| Wind Dispersed Seed | Animal Dispersed Seed |
|---|--|
| Dandelion  | Apple (internal)  |
| Sycamore  | Bur (external)  |

16.6. Germination

- A process controlled by enzymes
- **Water:** activates enzymes to turn insoluble food stores into soluble substances, and makes tissues swell so that the testa splits
- **Oxygen:** enters through the gaps in the testa (along with water), and is used in aerobic respiration.
- **Temperature:** must be suitable for enzymes to work (at optimum temperature).


16.7. Sexual Reproduction In Humans

- Male reproductive system:




- **Testes:** have many coiled tubes which produce sperm, and the cells between tubes produce testosterone.
- **Scrotum:** holds testicles
- **Sperm duct:** carries sperm from testicles to urethra.
- **Prostate gland:** makes seminal fluid
- **Urethra:** carries semen from sperm duct to tip of penis
- **Penis:** male sex organ, used to transfer semen to the female.

• **Female reproductive system:**



- **Ovary:** contains follicles which develop into the ova and produces progesterone and oestrogen
- **Oviduct (fallopian tube):** carries the ovum to uterus
- **Uterus (womb):** where the fetus develops.
- **Cervix:** neck of uterus: a strong rigid muscle, moist by mucus with a small opening
- **Vagina:** receives penis during intercourse, and way out for baby at birth. Moist tube of muscle, flexible and secretes mucus

16.8. Menstrual Cycle



- **Day 1 to 5:**
 - In the ovary, *FSH* secreted by the *Pituitary Gland* to stimulate the maturation of ONE *follicle* in the ovary.
 - In the uterus: the *endometrium* breaks down; menstruation
- **Day 5 to 12:**
 - In the ovary the follicle keeps maturing
 - In the uterus, *oestrogen* is secreted by *follicle* and the ovarian tissues to prepare the *endometrium*
- **Day 13/14/15:**
 - In the ovary, *LH* is also secreted by the *Pituitary Gland* to trigger the release of the egg from follicle into the fallopian tube
- **Day 15 to 28:**
 - In the ovary, *LH* triggers formation of *Corpus Luteum*
 - In the uterus: *progesterone* is secreted by *Corpus Luteum* to keep *endometrium* thick, waiting for possible embryo implant.
- **Day 28 – Scenario 1:** Egg not fertilized
 - No implantation takes place, the *Corpus Luteum* degenerates, causing a lack of progesterone.
 - This means that *endometrium* is no longer thick, back to Day 1
- **Day 28 – Scenario 2:** Egg is fertilized
 - Implantation occurs.
 - This makes the hormones to keep the *Corpus Luteum* maintained which means that *progesterone* is high.
 - This keeps the *Endometrium* thick for pregnancy

16.9. Hormones in Menstrual Cycle

- **Oestrogen** is secreted by the ovaries. It stops FSH being produced - so that only one egg matures in a cycle and it stimulates the pituitary gland to release hormone LH.
- **Progesterone** is a hormone secreted by ovaries. It maintains the lining of the uterus during the middle part of the menstrual cycle and during pregnancy.
- **Follicle stimulating hormone (FSH)** is secreted by the pituitary gland. It causes an egg to mature in an ovary and it stimulates ovaries to release hormone oestrogen
- **Luteinizing hormone (LH):** is also secreted by pituitary gland and causes mature egg to be released from ovary.

16.10. Sexual Intercourse

- Penis fills with blood and becomes erect
- Vagina walls secrete a lubricant.
- Rubbing of the glans (end of penis) against the vagina wall sets of a reflex action, causes sperm to be released from the testes, and is transported by peristalsis along sperm ducts and urethra, where seminal fluid is added to make semen.
- The exit of semen from the penis is called ejaculation.
- Sperm then swim through the cervix and oviducts to the first third of the oviduct (from the ovary) where one combines with the egg.

16.11. Fertilization

- The fusion of an ovum and a sperm to form a zygote.
- **Development of zygote:**
 - One sperm penetrates
 - Ovum membrane alters to form barrier against sperm
 - Head of sperm (male nucleus) approaches and then fuses with the nucleus of the ovum.
 - Zygote divides over and over, to make a ball of cells called an embryo.
 - It implants itself in the wall of the nucleus (implantation) which is followed by conception
- **Development of foetus:** zygote is changed through growth (mitosis) and development (organization of cells into tissues and organs)
- **Umbilical cord:** contains umbilical artery which carries deoxygenated blood and waste products from foetus to placenta and umbilical vein which carries oxygenated blood and soluble food from placenta to foetus. (Contains foetus' blood)
- **Placenta:** organ for exchange of soluble materials such as foods, wastes and oxygen between mother and foetus; physical attachment between uterus and foetus. (Contains mother's blood)
- **Amniotic sac:** membrane which encloses amniotic fluid, broken at birth.
- **Amniotic fluid:** protects foetus against mechanical shock, drying out and temperature fluctuations

16.12. Antenatal Care:

- **Change in diet:**
 - More proteins → growth of foetus
 - Slightly more fat → the new cells' cell membrane
 - More vitamin C and D → blood vessel walls and bones
 - Iron → haemoglobin
 - Calcium → growth of bones and teeth
- Guidance on motherhood
- Checks on foetus and mother including: weight check, blood tests, urine tests, blood pressure checks, ultrasound scanning etc.

16.13. Labour and Birth

- **Labour:** The uterine muscular wall contract and cervix tries to relax, then contractions get more frequent. Contractions cause amniotic membrane to break and release amniotic fluid.
- **Expulsion:** Powerful Contraction pushes baby out.
- **Afterbirth:** Placenta is expelled out. All contraction & pain gone

| Gamete | Size | Mobility | Number |
|--------|---------|---------------------------------|-------------------------|
| Sperm | Smaller | Very mobile - use its tail | Many more (300,000,000) |
| Egg | Larger | Immobile - moved by peristalsis | Fewer and limited |

| Damage beauty | | No additives /preservatives | | Builds mother-child bond | | No cost/preparation | | Causes decline in uterus size | |

Bottle feeding:

| Advantages | Disadvantages |
|--|--------------------------------|
| Less painful | More likely to develop illness |
| Other people can feed baby | Risk of wrong mixture |
| May contain supplement vitamins and minerals | Expensive |

16.14. Sex Hormones

- At puberty, the pituitary gland starts to stimulate the primary sex organs; the testes in males and the ovaries in females.
- Sex hormones - testosterone in males and oestrogen in females are released into the bloodstream.
- They only affect the target organs which have receptors which can recognize them.
- Causes secondary sexual characteristics such as the growth of pubic hair and maturation of sexual organs.

16.15. Methods of Birth Control

- **Natural:**
 - Abstinence: don't have sex
 - Rhythm method: don't have sex during the fertile period, only during the safe period
- **Chemical:**

- Progesterone-only pill: pill which affects the uterus and makes implantation difficult
- Spermicide: a chemical applied as a gel, cream or foam which kills sperm. It is very unreliable on its own but makes barrier methods of contraception more effective.
- **Mechanical:**
 - Condom: thin rubber covering over penis, it protects from impregnation and STDs, used by man
 - Diaphragm: used by woman, prevent sperm entering uterus, reliable, must stay in place 6 hours after sex, needs a correct size
 - Femidom: closed end, has a ring which gets pushed through cervix and open end's ring lies against the labia
 - IUD: plastic-coated copper coil, can be left inside for months or even years, has a string which is used to remove it out of the vagina, reliable, it irritates uterus wall preventing implantation
- **Surgical:**
 - Vasectomy: sperm ducts are cut and tied
 - Female sterilization: oviducts are cut and tied

16.16. Artificial Insemination

- **By donor:** man's sperm has a problem, making impregnation impossible, so a donor gives his sperm.
- **In vitro fertilization:** an ovum is fertilized outside a woman's body. The fertilized ovum is implanted into the uterus.
- **Fertility drugs:** drugs which enhance reproductive fertility. For women, fertility medication is used to stimulate follicle development of the ovary. The side effect is multiple pregnancies. They contain varying amount of FSH and LH.

16.17. Human Immunodeficiency Virus (HIV)

- **Transmission:** Intercourse, blood transfusion, organ transplant or sharing needle with infected person
- **Prevention:**
 - Avoid intercourse with many partners
 - Use a condom
 - Don't come in contact with other people's blood
- **How it affects the immune system:**
 - Infects and destroys lymphocytes
 - Decreases efficiency of immune system
 - Body becomes liable to infection by other pathogens

17. Inheritance

- **Inheritance:** The transmission of genetic information from generation to generation

17.2. Chromosome

- **Chromosome:** a thread-like structure of DNA, made up of a string of genes
- **Gene:** a length of DNA that is the unit of heredity and codes for a specific protein. A gene may be copied and passed on to the next generation
- **Allele:** any of two or more alternative forms of a gene
- **Haploid nucleus:** a nucleus containing a single set of unpaired chromosomes (e.g. sperm and egg)
- **Diploid nucleus:** a nucleus containing two sets of chromosomes (e.g. in body cells)
- **Inheritance of gender in humans:** woman's gamete can only carry an "X" chromosome, and a male gamete can carry either an "X" or "Y" chromosome; females are "XX" while males are "XY". There is always a 50% chance of getting a boy and vice versa.

17.3. DNA & Protein Synthesis

- **DNA:** control cell functions by controlling production of proteins, antibodies and receptors for neurotransmitters
- DNA has 2 long strands and 4 nucleotides, AT and CG
- How proteins are made:
 - Made from long chains of amino acids (20)
 - DNA bases are read as triplets
 - DNA is found in the nucleus
 - Protein synthesis happens on ribosome in cytoplasm
 - mRNA carries information from DNA to ribosome
 - When a protein is made, mRNA is made in the nucleus, copying the base sequence
 - mRNA moves out of the nucleus into the cytoplasm and attaches to the ribosome
 - ribosome assembles amino acids into protein molecules
 - the specific order of amino acids is determined by the sequence of bases in the mRNA
- All body cells in an organism contain the same genes, but many genes in a particular cell are not expressed because the cell only makes the specific proteins it needs

17.4. Mitosis

- The nuclear division giving rise to genetically identical cells in which the chromosome number is maintained by the exact duplication of chromosomes.
- Mitosis is needed for:
 - **Growth:** in animals each tissue provides its own new cells when they are needed.
 - **Repair** of damaged tissues: for example, when you cut your skin, mitosis provides new cells to cover up cut.
 - **Replacement** of worn out cells
 - **Asexual reproduction:** in plants
- **Stem cells:** unspecialized cells that divide by mitosis to produce daughter cells that can become specialized for specific functions

17.5. Meiosis


- Reduction division in which the chromosome number is halved from diploid to haploid
- Gametes are the result of meiosis
- Meiosis results in genetic variation so the cells produced are not all genetically identical.

17.6. Monohybrid Inheritance

- **Genotype:** genetic makeup of an organism in terms of the alleles present (e.g. Tt or GG)
- **Phenotype:** physical or other features of an organism due to both its genotype and its environment (e.g. tall plant or green seed)


genotype + environment + random variation → phenotype

- **Homozygous:** having two identical alleles of a particular gene (e.g. TT or gg). Two identical homozygous individuals that breed together will be pure-breeding
- **Heterozygous:** having two different alleles of a particular gene (e.g. Tt or Gg), not pure-breeding
- **Dominant:** an allele that is expressed if it is present (e.g. T or G)
- **Recessive:** an allele that is only expressed when there is no dominant allele of the gene present (e.g. t or g)
- **Pedigree diagrams:**



Genetic diagrams:
1:1 Monohybrid Crosses

A Genetic Diagram




F1 Phenotype: 2 : 2 = 1 : 1 Boy : Girl

Note the gender of the baby is determined by the sperm!

3:1 Monohybrid Crosses

A Genetic Diagram



F1 Phenotype: 3 : 1 Brown eyes : blue eyes

Note the gametes are always put in circles

- **Co-dominance:** when neither of two alleles is dominant to each other.
- There are three alleles for blood group given by the symbols I^A , I^B and I^O .
- I^A and I^B are co-dominant giving blood group AB or $I^A I^B$, and both dominant to I^O .
- **Sex-linked characteristic:** a characteristic in which the gene responsible is located on a sex chromosome and that this makes it more common in one sex than in the other
 - Colour blindness as an example of sex linkage

18. Variation & Selection

18.1. Variation

- **Variation:** differences between individuals of the same species
- Phenotypic variation is caused by both genetic and environmental factors

- **Continuous variation** is influenced by genes and environment, resulting in a range of phenotypes between two extremes, e.g. height in humans
- **Discontinuous variation** is caused by genes alone and results in a limited number of distinct phenotypes (e.g. you are either blood group O, A, B or AB, nothing else)
- **Mutation:** genetic change
- **Gene mutation:** a change in the base sequence of DNA
- Mutation is the way in which new alleles are formed
- Ionising radiation and some chemicals increase the rate of mutation
- Mutation is a source of variation e.g. in Down's syndrome, where a parent's chromosomes are unevenly distributed in meiosis. In fertilisation, a zygote with a number of chromosomes that is not 46 is created (e.g. 23 + 24). Characteristics: broad forehead, short neck, downward-sloping eyes, short nose and mental retardation.

18.2. Sickle Cell Anaemia

- Disease in which the red blood cell has a sickle shape instead of a round biconcave shape, controlled by a recessive allele, which causes weakness, aching joints and poor circulation.
- The fact that it is recessive means that a heterozygous person can be a carrier: they have the allele but it is not expressed.
- Being a carrier of sickle cell anaemia makes you resistant to malaria
- In equatorial Africa, having sickle cell anaemia causes death. Malaria also causes death. But the carriers have immunity to malaria and have some symptoms of anaemia, in severe cases, they are very weak

18.3. Adaptive Features

- **Adaptive feature:**
 - inherited feature that helps an organism to survive and reproduce in its environment
 - the inherited functional features of an organism that increase its fitness
- **Fitness:** the probability of an organism surviving and reproducing in the environment in which it is found
- **Xerophytes:** live in deserts where water is scarce and evaporation is rapid, or in windy habitats. Their features are:
 - Deep roots to reach water far underground
 - Leaves reduced spines with minimum surface area for transpiration
 - Shallow spreading roots to collect occasional rain
 - Rolled leaves, leaf hairs and stomata sunk in pits to trap moist air
 - Waxy leaf cuticle, impermeable water
 - Stomata opening at night and closed at midday when evaporation is highest
 - E.g. cactus and marram grass

- **Hydrophytes:** live in wholly or partly submerged in water. Their features are:
 - Leaves are highly divided to create large surface area for absorption and photosynthesis
 - Very little cuticle formation
 - Lack of xylem tubes, no stomata underside of leaves
 - Stomata is in the upper surface, has a thick waxy layer to repel water and to keep the stomata open and clear
 - Roots are often reduced and root hairs are often absent

18.4. Natural Selection

- The greater chance of passing on of genes by the best adapted organisms.
- Variation is natural or random changes in all living organisms.
- Variation leads to survival of the fittest since the variations in certain organisms allow that organism to have an advantage over the others in its species in that area
- The surviving organisms reproduce, since they don't get eaten up, so variation has caused the species to evolve.
- Evolution is caused by natural selection which is caused by a change in the environment

18.5. Artificial Selection

- Is breeding organisms with valued characteristics together in order to try to produce offspring which shares those useful characteristics (selective breeding).
- Can be used to produce organisms which are more economically valued
- For example: cows that produce more milk, wheat that is easier to separate from grain, dogs which have better appearance
- **Selective breeding:**
 - Selecting by humans of individuals with desirable features
 - Crossing three individuals to produce the next generation

18.6. Resistant Antibiotic

- Strains of antibiotic-resistant bacteria are developing as the use of antibiotics is increasing.
- In a group of many bacteria, one might mutate to be resistant to the antibiotic, as a result it reproduces and the others die making a new strain of bacteria, which is resistant to antibiotics.
- The susceptible (weak) bacteria are killed first
- Only resistant (strong) bacteria are left

19. Organisms and their Environment

19.1. Food Chains and Food Webs

- The sun is the principal source of energy input to biological systems.
- Energy flow is **not** a cycle; it starts from the sun and then that energy is harnessed by plants which are eaten by animals which are eaten by other animals.
- At each step, energy is lost to the environment.
- Food chain:** a chart showing the flow of energy (food) from one organism to the next beginning with a producer, for example:

Mahogany tree → caterpillar → song bird → hawk

- Food web:** showing the transfer of energy from one organism to the next, beginning with a producer
 - Energy is transferred between organisms in a food chain by ingestion
- Producer:** an organism that makes its own organic nutrients, usually using energy from sunlight, through photosynthesis
- Consumer:** an organism that gets its energy by feeding on other organisms
- Herbivore:** an animal that gets its energy by eating plants
- Carnivore:** an animal that gets its energy by eating other animals
- Decomposer:** an organism that gets its energy from dead or waste organic matter (i.e. a saprotroph)
- Ecosystem:** a unit containing all of the organisms and their environment, interacting together, in a given area e.g. decomposing log or a lake
- Trophic level:** position of an organism in a food chain, food web or pyramid of biomass, numbers or energy
 - Primary consumer:** eat vegetables
 - Secondary consumer:** eat meat/drink milk
 - Tertiary consumer:** eat a predatory fish, salmon

Food chains usually have fewer than five trophic levels, because energy transfer is inefficient:


- Sun produces light, less than 1% of the energy falls onto leaves.
- Producers 'fix' only about 5-8% of that energy, because of: transmission, reflection and incorrect wavelength.
- Primary consumer only gets between 5-10% because some parts are indigestible (e.g. cellulose) and not eating the whole plant.
- Secondary consumer gets between 10-20% because animal matter is more digestible & has higher energy value
- At each level heat is lost by respiration.

Humans eating plants is more efficient than humans eating animals because...


- We need only a couple of vegetables to have one meal, but to have meat we must feed the animal a lot of plant material in order to get far less meat.
- In the process of raising an animal, plants lose energy to environment, then animal loses energy to environment and does not use up all the plant material so it is very inefficient.

| Pyramid of Numbers | Pyramid of Biomass |
|--|---|
| | |
| Shows number of each organism in a food chain | Pyramid which shows the biomass |
| When moving up pyramid, number of individuals' decreases | (number of individuals × their individual mass) |

19.2. Carbon Cycle




19.3. Nitrogen Cycle



- Nitrogen-fixing bacteria provide usable nitrogen for plants, these may exist in the root nodules where they live in symbiosis with the plants (nitrogen fixation), or this can happen because of lightning, or microorganisms provide them through decomposition.
- Nitrifying bacteria convert nitrogen-containing substances into better nitrogen-containing substances for the plants (nitrification).
- Plants absorb these substances and convert them into proteins
- Primary consumers eat the plants and can make their own proteins, secondary consumers eat primary consumers and so on.

- Death and decay happens at each trophic level leading to stage one
- Denitrifying bacteria carry out denitrification: they convert nitrogen-containing substances into atmospheric nitrogen

19.4. Water Cycle



| Methods | O ₂ conc. in air | CO ₂ conc. in air | Why? |
|----------------------------|-----------------------------|------------------------------|---|
| Combustion of fossil fuels | Decreases | Increases | Burning uses up oxygen, and produces carbon dioxide |
| Deforestation | Decreases | Increases | Fewer trees means less photosynthesis |
| | | | Trees are usually burnt (combustion) |
| | | | Decomposition of tree trunks (respiration) |


19.5. Population

- **Population:** a group of organisms of one species, living in the same area at the same time
- **Community:** all of the populations of different species in an ecosystem
- **Ecosystem:** a unit containing the community of organisms and their environment, interacting together, e.g. a decomposing log, or a lake

19.6. Factors Affecting Rate of Population Growth

- **Food supply:** quantity and quality, for example snails need calcium to reproduce to make a shell (food quality).
- **Predation:** if predator population falls, the prey population will rise
- **Disease:** causes organisms to die so a high death rate partly cancels out birth rate meaning less population growth, especially if the organism dies before giving birth, or even population decline

19.7. Sigmoid Curve



- **Lag phase:** number of mature, reproducing individuals is low and they may be widely dispersed
- **Log phase:** exponential growth occurs, the conditions are ideal and maximum growth rate is reached. Limiting factors do not limit growth much.
- **Stationary phase:** limiting factors slow growth as population has reached "carrying capacity" of its environment; when mortality rate = birth rate; curve levels off and fluctuates around this maximum population size.

19.8. Human Population Growth

| Factors favouring growth | Factors controlling growth |
|--------------------------|----------------------------|
| Lower infant mortality | Disease |
| Higher life expectancy | famine |
| Better nutrition | War |
| Better housing | |
| Better sanitation | |
| Medicine | |
| Vaccination | |

- The human population is **becoming stable (stagnation) due to:**
 - better education (particularly for women), so they work instead of getting married and having children
 - better living conditions, fewer people die, fewer births needed
 - cities, reduced need for physical labour on farms
 - family planning
- But overall the population is still increasing.
- **Social implications** of human growth:
 - demands for roads as there is an increases number of cars
 - greater expectation for a variety of foods all year round
 - smaller families increase demand for housing
 - greater demand for leisure and recreation space

20. Biotechnology & Genetic Engineering

- Bacteria are useful in biotechnology and genetic engineering due to their rapid reproduction rate and their ability to make complex molecules
- Why are microorganisms used:
 - Bacteria and fungi are small and easy to grow in a lab
 - They reproduce quickly and don't take up much space

- No ethical issues involved
- Genetic code is the same for bacteria as it is for human
- Bacteria have loops of DNA called plasmids which are easy to transfer from one cell to another

20.2. Making Biofuel

- Use plants to make sugars which yeast then breaks down to make ethanol
- Ethanol can then be used as a fuel
 - Maize is treated with amylase enzyme (starch to glucose)
 - Add yeast (glucose used in respiration) and make it respire anaerobically
 - Ethanol that is produced is extracted by distillation
 - Mixed with gasoline to increase energy and can be used in cars

20.3. Bread Making

- Flour, sugar, water and salt are mixed with yeast to make the dough.
- Amylase breaks down some starch to make maltose and glucose. This is used by yeast in respiration
- The dough is kept in a warm, moist environment (28°C). Yeast ferments sugar making carbon dioxide which creates bubbles, so bread rises
- Cooking (at 180°C) – kills yeast, evaporates alcohol and hardens outer surface.

20.4. Uses of Enzymes

Pectinase:

- Fruit juices are extracted using pectinase (breaks down pectin)
- Pectin helps plant walls stick together
- If pectin is broke down, it's easier to squeeze juice from the fruit
- Extraction of juice from fruit, making juice clear not cloudy


Washing powders:

- Biological washing powders and liquids contain enzyme that help remove stain
- The enzymes are coated with a special wax that melts in the wash releasing the enzyme
- Once the stains have been broken down, they are easier for detergents to remove
- **Proteases** break down proteins in stains e.g. grass, blood
- **Lipases** break down stains containing fats and oil
- **Carbohydrases** break down carbohydrate-based stains, such as starch

Lactase:

- Enzyme that breaks down lactose (sugar found in milk)
- People can stop making lactase naturally, therefore can't digest lactose
- Milk can be treated with lactase to break down lactose before a person drinks it
- Also produces glucose & galactose, used in sweets

20.5. Making Penicillin



- **Penicillin:** an antibiotic produced by a fungus called penicillium.
- Stainless steel fermentation vessel is filled with medium containing sugars and ammonium salts.
- Penicillium is added to produce penicillin. They use sugar for respiration and ammonium salts to make protein and nucleic acids
- The fermentation vessel consists of **PAWS**
 - Probes monitor temperature and pH
 - Air provides oxygen for aerobic respiration in fungus
 - Water-cooled jacket removes heat to maintain temperature of 24C.
 - Stirrer keeps the microorganism suspended (allowing access to nutrients and oxygen) while maintaining an even temperature.
- Filtered to remove fungus and then can be crystallized to make capsules.

20.6. Genetic Engineering

- **Genetic engineering:** changing the genetic material of an organism by removing, changing or inserting individual genes
- Examples of genetic engineering:
 - the insertion of human genes into bacteria to produce human insulin
 - the insertion of genes into crop plants to confer resistance to herbicides
 - the insertion of genes into crop plants to confer resistance to insect pests
 - the insertion of genes into crop plants to provide additional vitamins

Human Insulin in Bacteria

- Isolation of the DNA making up a human gene using restriction enzymes, forming sticky ends
- Cutting of bacterial plasmid DNA with the same restriction enzymes, forming complementary sticky ends
- Insertion of human DNA into bacterial plasmid DNA using DNA ligase to form a recombinant plasmid – insertion of plasmid into bacteria
- Replication of bacteria containing recombinant plasmids which make human protein as they express the gene

20.7. Genetically Modified Crops

| Advantages | Disadvantages |
|--|---|
| Uniform in shape – easy to transport/ appeal consumers | Natural species may die |
| Growing season shorter | Taste often not as good |
| Drought resistant – less water | Lead to development of super weeds – stronger than GM |
| Higher yields | No one knows long term effect on humans |

21. Human Influences on Ecosystems

21.1. Food Supply

Food production has increased because:

- **Improved machinery** means less labour is needed
- **Fertilizers** help crops to grow better
- **Insecticides**: a type of pesticide that kills insects
- **Herbicides**: a type of pesticide that kills weeds
- **Artificial selection** and **genetic modification** means that yields are improved: cows produce more milk, cows are more muscular giving more meat, plant crops can resist insects and cold weather

World Food Supplies

- Not enough food available in a country to feed its people because:
 - Fast increase in population
 - Increasing use of crops for fuel
 - Decrease of farming = Climate change/Urbanization
- Famine: Wide spread scarcity of food
- The main causes of famine:
 - The rapid rate of population increase
 - Long term climatic change
 - Soil erosion and desertification
 - Economic pressure
 - Unequal distribution of food
 - Drought
 - Flood

- **Monoculture** is the continuous production of one type of crop that is often genetically uniform.
- **Negative Impacts of Monoculture**
 - If a natural disaster were to occur, the whole crop could be wiped out.
 - If pests & disease attacked crop, it could harm it easily
 - Using large fields and pesticides reduces the variety of species. This hinders biodiversity.
- **Negative Impacts of Intensive Livestock Production**
 - Welfare issues for the livestock
 - Diseases can spread easily among them
 - Waste can pollute land and waterways nearby

21.2. Habitat Destruction

- **Reason for habitat destruction**
 - Increased area for food crop growth, livestock production and housing
 - Extraction of natural resources
 - Marine pollution
- Through altering food webs, and food chains, humans can have a negative impact on habitats
- **Effects of deforestation**
 - Reduced biodiversity/destroys habitats/extinction
 - Loss of CO₂ fixation, thus increase in CO₂, thus global warming
 - **Soil erosion**: tree roots cannot retain soil, goes into rivers making the water dirty & causes blockages, soil becomes less fertile
 - **Flooding**: usually 75% of water is absorbed by foliage, root systems or evaporates. After deforestation, water accumulates in valleys

21.3. Pollution

Water and air pollution:

- Chemical waste and sewage in rivers results in water not being drinkable and eutrophication can occur
- Sulphur dioxide dissolves in rain, causing acid rain which increases acidity of lakes and leaches aluminium out of the soil causing:
 - The fishes' gills are damaged, eventually killing them. This is fixed by adding calcium hydroxide (slaked lime)
 - Destroys top of trees and aluminium damages tree roots = dead tree, important nutrients leached away
 - SO₂ poses health hazards for humans (asthma sufferers)
 - Damages limestone buildings and sculptures
 - Fewer crops can be grown on an acidic field (fixed by adding lime)

Pollution due to pesticides:

- **Insecticides (kill insects)**: meant to kill insects which eat crops, but can kill other, useful insects such as bees

which are pollinators, or by bioaccumulation (the increase in dose of toxin from one level of the food chain to the next)

- **Herbicides (kill weeds):** can be harmful to animals which eat the plants

Nuclear fall-out:

- Radioactive particles are sprayed into the atmosphere in a nuclear accident or bombing;
- These particle "rain" back to earth from clouds, sometimes far from the accident site;
- The radioactivity damages DNA and causes cancer and radiation illness at every level of the food chain.

Non-biodegradable plastics:

- Choke birds, fish and other animals
- Fill up the animals' stomachs so that they can't eat food
- Collect in rivers, and get in the way of fish

Acid rain:

- Caused by sulphur dioxide (burning fossil fuels) and nitrogen oxides (nitrogen reacting hot engines), as they dissolve and cause acid rain
- Damages trees and plants, and kills fish and other river life
- Prevention: catalytic converters, in factories slaked lime neutralizes these acidic oxides and use of flue-gas desulfurization


Global Warming:

- Increase in average temperature of the Earth
- Started at the same time as humans began burning fossil fuels
- Scientists believe fossil fuels are causing this – not proven yet
- Increase in carbon dioxide and methane concentrations in the atmosphere cause an enhanced greenhouse effect the leads to climate change
- **Negative impact of female contraceptive hormones in water courses:**
- Reduced sperm count in men and feminisation of aquatic organisms

Eutrophication: when water plants receive too many nutrients.

- Fertilisers put in soil by farmers
- Fertilisers with nitrates / detergents with phosphates leach into rivers and lakes after rain
- Water plants grow more than usual
- They block sunlight and kill plants underneath
- They die and sink to bottom

- Bacteria/fungi decompose remains using the O₂ and decreasing the O₂ concentration
- Fish and other creatures die from oxygen starvation



21.4. Conservation

- **Sustainable resource:** one which is produced as rapidly as it is removed from the environment so that it does not run out
- **Sustainable development:** development providing for the needs of an increasing human population without harming the environment
- Sustainable development requires:
 - Management of conflicting demands
 - Planning and co-operation at local, national and international levels
- Some resources can be maintained, limited to forests and fish stocks.
- They can be sustained using:
 - Education
 - Legal quotas
 - Re-stocking

Natural resources:

- **Water:** used to grow food, keep clean, provide power, control fires and to drink. We get water constantly through rainfall but we are using up planet's fresh water faster than it can be replenished.
- **Fossil fuels:** need to be conserved as they will soon run out, they should be therefore replaced with green forms of energy.

Recycling:

- **Water:** water from sewage can be returned to environment for human use by sanitation and sewage treatment
- **Paper:** sent to special centres where it is pulped to make raw materials for industry
- **Plastic:** fossil fuels, bottles → fleece clothing
- **Metal:** mining takes a lot of energy so recycling saves energy

Species and habitats: need to be conserved because:

- Organisms have value in themselves (ethical value)
- Value to medicine (new molecules from exotic plants = new drugs)
- Genetic resources are useful to humans as well and are lost when species disappear (DNA for genetic engineering)
- Each species has its role in its ecosystem; if it is removed, then the whole ecosystem could collapse

Endangered species:

- How they become endangered: climate change, habitat destruction, hunting, pollution and introduced species
- If the population size drops, variation decreases
- Endangered species can be conserved by: monitoring and protecting species and habitats, education, captive breeding programmes and seed banks

• Conservation programmes include:

- reducing extinction
- protecting vulnerable environments
- maintaining ecosystem functions, by nutrient cycling and resource provision, e.g. food, drugs, fuel and genes

CAIE IGCSE Biology (0610)

Copyright 2022 by ZNotes

These notes have been created by Maimoona Junjuna for the 2020-21 syllabus

This website and its content is copyright of ZNotes Foundation - © ZNotes Foundation 2022. All rights reserved.

The document contains images and excerpts of text from educational resources available on the internet and printed books. If you are the owner of such media, text or visual, utilized in this document and do not accept its usage then we urge you to contact us and we would immediately replace said media.

No part of this document may be copied or re-uploaded to another website without the express, written permission of the copyright owner. Under no conditions may this document be distributed under the name of false author(s) or sold for financial gain; the document is solely meant for educational purposes and it is to remain a property available to all at no cost. It is current freely available from the website www.znotes.org

This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.