

Centre Number	Candidate Number	Name
---------------	------------------	------

CAMBRIDGE INTERNATIONAL EXAMINATIONS
International General Certificate of Secondary Education

CHEMISTRY

0620/03

Paper 3

October/November 2003

1 hour 15 minutes

Candidates answer on the Question Paper.
No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number at the top of this page.
Write in dark blue or black pen in the spaces provided on the Question Paper.
You may use a pencil for any diagrams, graphs or rough working.
Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer **all** questions.


The number of marks is given in brackets [] at the end of each question or part question.
A copy of the Periodic Table is printed on page 12.

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

For Examiner's Use	
1	
2	
3	
4	
5	
TOTAL	

- 1 Ammonia contains the elements nitrogen and hydrogen. It is manufactured from these elements in the Haber process.



The forward reaction is exothermic.

- (a) (i) Nitrogen is obtained from liquid air by fractional distillation. Why does this technique separate liquid oxygen and nitrogen?

.....

- (ii) Name **two** raw materials from which hydrogen is manufactured.

.....[3]

- (b) The table shows how the percentage of ammonia in the equilibrium mixture varies with pressure at 600 °C.

percentage ammonia	8	12	15	20
pressure/atm	200	300	400	500

- (i) Explain why the percentage of ammonia increases as the pressure increases.

.....
[2]

- (ii) How would the percentage of ammonia change if the measurements had been made at a lower temperature?
 Explain your answer.

.....

[2]

- (iii) State **two** of the reaction conditions used in the Haber Process.

.....
[2]

(c) Ammonia is a base.

(i) Name a particle that an ammonia molecule can accept from an acid.

.....

(ii) Write an equation for ammonia acting as a base.

.....[3]

(d) Given aqueous solutions, 0.1 mol/dm^3 , of sodium hydroxide and ammonia, describe how you could show that ammonia is the weaker base.

.....

.....[2]

(e) Another compound that contains nitrogen and hydrogen is hydrazine, N_2H_4 .


(i) Draw the structural formula of hydrazine. Hydrogen can form only one bond per atom but nitrogen can form three.

(ii) Draw a diagram that shows the arrangement of the valency electrons in one molecule of hydrazine. Hydrazine is a covalent compound.
Use x to represent an electron from a nitrogen atom.
Use o to represent an electron from a hydrogen atom.

[3]

- 2 Some of the factors that can determine the rate of a reaction are concentration, temperature and light intensity.


- (a) A small piece of calcium carbonate was added to an excess of hydrochloric acid. The time taken for the carbonate to react completely was measured.



The experiment was repeated at the same temperature, using pieces of calcium carbonate of the same size but with acid of a different concentration. In all the experiments an excess of acid was used.

concentration of acid / mol dm ⁻³	4	2	2
number of pieces of carbonate	1	1	2	1
time / s	80	160

- (i) Complete the table (assume the rate is proportional to both the acid concentration and the number of pieces of calcium carbonate). [3]
- (ii) Explain why the reaction rate would increase if the temperature was increased.
.....
.....[2]
- (iii) Explain why the rate of this reaction increases if the piece of carbonate is crushed to a powder.
.....[1]
- (iv) Fine powders mixed with air can explode violently. Name an industrial process where there is a risk of this type of explosion.
.....
.....[1]
- (b) Sodium chlorate(I) decomposes to form oxygen and sodium chloride. This is an example of a photochemical reaction. The rate of reaction depends on the intensity of the light.



- (i) Describe how the rate of this reaction could be measured.


.....
.....[2]

(ii) How could you show that this reaction is photochemical?

.....
[1]

(c) Photosynthesis is another example of a photochemical reaction. Glucose and more complex carbohydrates are made from carbon dioxide and water.

(i) Complete the equation.



(ii) Glucose can be represented as



Draw the structure of a more complex carbohydrate that can be formed from glucose by condensation polymerisation.

[2]

3 Zinc blende is the common ore of zinc. It is usually found mixed with an ore of lead and traces of silver.

(a) (i) Describe how zinc blende is changed into zinc oxide.

.....
[2]

(ii) Write an equation for the reduction of zinc oxide by carbon.

.....[2]

(iii) The boiling point of lead is 1740 °C and that of zinc is 907 °C. Explain why, when both oxides are reduced by heating with carbon at 1400 °C, only lead remains in the furnace.

.....
[2]

(b) A major use of zinc is to make diecasting alloys. These contain about 4% of aluminium and they are stronger and less malleable than pure zinc.

(i) Give one other large scale use of zinc.

.....[1]

(ii) Describe the structure of a typical metal, such as zinc, and explain why it is malleable.

.....

.....

.....[3]


(iii) Suggest why the introduction of a different metallic atom into the structure makes the alloy stronger than the pure metal.

.....

.....[2]

(c) A solution of an impure zinc ore contained zinc, lead and silver(I) ions. The addition of zinc dust will displace both lead and silver.

(i) The ionic equation for the displacement of lead is as follows.



Which change is reduction? Explain your answer.

.....

.....[2]

(ii) Write an ionic equation for the reaction between zinc atoms and silver(I) ions.

.....[2]

- 4 Esters occur naturally in plants and animals. They are manufactured from petroleum. Ethyl ethanoate and butyl ethanoate are industrially important as solvents.

(a) (i) Explain the term *solvent*.

.....[1]

(ii) Give the formula of ethyl ethanoate.

[1]

(iii) Ethyl ethanoate can be made from ethanol and ethanoic acid. Describe how these chemicals can be made.

ethanol from ethene

.....
.....[2]


ethanoic acid from ethanol

.....
.....[2]

(iv) Name **two** chemicals from which butyl ethanoate can be made.

.....[1]

(b) The following equation represents the alkaline hydrolysis of a naturally occurring ester.




(i) Which substance in the equation is an alcohol? Underline the substance in the equation above.

[1]

(ii) What is the major use for compounds of the type $\text{C}_{17}\text{H}_{35}\text{COONa}$?

(c) A polymer has the structure shown below.



(i) What type of polymer is this?

.....[1]

(ii) Complete the following to give the structures of the two monomers from which the above polymer could be made.



[2]

(d) Esters are frequently used as solvents in chromatography. A natural macromolecule was hydrolysed to give a mixture of amino acids. These could be identified by chromatography.

(i) What type of macromolecule was hydrolysed?

.....[1]

(ii) What type of linkage was broken by hydrolysis?

.....[1]

(iii) Explain why the chromatogram must be sprayed with a locating agent before the amino acids can be identified.

.....
.....[1]

(iv) Explain how it is possible to identify the amino acids from the chromatogram.

.....
.....[2]

5 Sulphur dioxide, SO_2 , and sulphur trioxide, SO_3 , are the two oxides of sulphur.

(a) Sulphur dioxide can kill bacteria and has bleaching properties. Give a use of sulphur dioxide that depends on each of these properties.

(i) ability to kill bacteria[1]

(ii) bleaching properties[1]

(b) Sulphur trioxide can be made from sulphur dioxide.

(i) Why is this reaction important industrially?

.....[1]


(ii) Complete the word equation.

sulphur dioxide + \rightarrow sulphur trioxide [1]

(iii) What are the conditions for this reaction?

.....
.....[2]

(c) Sulphur dioxide is easily oxidised in the presence of water.




(i) What colour change would be observed when an excess of aqueous sulphur dioxide is added to an acidic solution of potassium manganate(VII)?

.....[2]

(ii) To aqueous sulphur dioxide, acidified barium chloride solution is added. The mixture remains clear. When bromine is added, a thick white precipitate forms. What is the white precipitate? Explain why it forms.

.....
.....[3]

(d) Sulphur dioxide reacts with chlorine in an addition reaction to form sulphuryl chloride.



8.0 g of sulphur dioxide was mixed with 14.2 g of chlorine. The mass of one mole of SO_2Cl_2 is 135 g.

Calculate the mass of sulphuryl chloride formed by this mixture.

Calculate the number of moles of SO_2 in the mixture =

Calculate the number of moles of Cl_2 in the mixture =

Which reagent was not in excess?

How many moles of SO_2Cl_2 were formed =

DATA SHEET
The Periodic Table of the Elements

		Group																																																																																																																	
I	II	III	IV	V	VI	VII	0																																																																																																												
7 Li Lithium	9 Be Beryllium	<table border="1" style="width: 100%; border-collapse: collapse; text-align: center;"> <tr> <td>1 H Hydrogen</td> <td colspan="10"></td> </tr> <tr> <td>11 B Boron</td> <td>12 C Carbon</td> <td>13 Al Aluminium</td> <td>14 Si Silicon</td> <td>15 P Phosphorus</td> <td>16 S Sulphur</td> <td>17 Cl Chlorine</td> <td>18 Ar Argon</td> <td>19 F Fluorine</td> <td>20 Ne Neon</td> <td>21 Sc Scandium</td> <td>22 Ti Titanium</td> <td>23 V Vanadium</td> <td>24 Cr Chromium</td> <td>25 Mn Manganese</td> <td>26 Fe Iron</td> <td>27 Co Cobalt</td> <td>28 Ni Nickel</td> <td>29 Cu Copper</td> <td>30 Zn Zinc</td> <td>31 Ga Gallium</td> <td>32 Ge Germanium</td> <td>33 As Arsenic</td> <td>34 Se Selenium</td> <td>35 Br Bromine</td> <td>36 Kr Krypton</td> <td>37 Rb Rubidium</td> <td>38 Sr Strontium</td> <td>39 Y Yttrium</td> <td>40 Zr Zirconium</td> <td>41 Nb Niobium</td> <td>42 Mo Molybdenum</td> <td>43 Tc Technetium</td> <td>44 Ru Ruthenium</td> <td>45 Rh Rhodium</td> <td>46 Pd Palladium</td> <td>47 Ag Silver</td> <td>48 Cd Cadmium</td> <td>49 In Indium</td> <td>50 Sn Tin</td> <td>51 Sb Antimony</td> <td>52 Te Tellurium</td> <td>53 I Iodine</td> <td>54 Xe Xenon</td> <td>55 Cs Caesium</td> <td>56 Ba Barium</td> <td>57 La Lanthanum</td> <td>58 Ce Cerium</td> <td>59 Pr Praseodymium</td> <td>60 Nd Neodymium</td> <td>61 Pm Promethium</td> <td>62 Sm Samarium</td> <td>63 Eu Europium</td> <td>64 Gd Gadolinium</td> <td>65 Tb Terbium</td> <td>66 Dy Dysprosium</td> <td>67 Ho Holmium</td> <td>68 Er Erbium</td> <td>69 Tm Thulium</td> <td>70 Yb Ytterbium</td> <td>71 Lu Lutetium</td> <td>72 Hf Hafnium</td> <td>73 Ta Tantalum</td> <td>74 W Tungsten</td> <td>75 Re Rhenium</td> <td>76 Os Osmium</td> <td>77 Ir Iridium</td> <td>78 Pt Platinum</td> <td>79 Au Gold</td> <td>80 Hg Mercury</td> <td>81 Tl Thallium</td> <td>82 Pb Lead</td> <td>83 Bi Bismuth</td> <td>84 Po Polonium</td> <td>85 At Astatine</td> <td>86 Rn Radon</td> <td>87 Fr Francium</td> <td>88 Ra Radium</td> <td>89 Ac Actinium</td> <td>90 Th Thorium</td> <td>91 Pa Protactinium</td> <td>92 U Uranium</td> <td>93 Np Neptunium</td> <td>94 Pu Plutonium</td> <td>95 Am Americium</td> <td>96 Cm Curium</td> <td>97 Bk Berkelium</td> <td>98 Cf Californium</td> <td>99 Es Einsteinium</td> <td>100 Fm Fermium</td> <td>101 Md Mendelevium</td> <td>102 No Nobelium</td> <td>103 Lr Lawrencium</td> </tr> </table>										1 H Hydrogen											11 B Boron	12 C Carbon	13 Al Aluminium	14 Si Silicon	15 P Phosphorus	16 S Sulphur	17 Cl Chlorine	18 Ar Argon	19 F Fluorine	20 Ne Neon	21 Sc Scandium	22 Ti Titanium	23 V Vanadium	24 Cr Chromium	25 Mn Manganese	26 Fe Iron	27 Co Cobalt	28 Ni Nickel	29 Cu Copper	30 Zn Zinc	31 Ga Gallium	32 Ge Germanium	33 As Arsenic	34 Se Selenium	35 Br Bromine	36 Kr Krypton	37 Rb Rubidium	38 Sr Strontium	39 Y Yttrium	40 Zr Zirconium	41 Nb Niobium	42 Mo Molybdenum	43 Tc Technetium	44 Ru Ruthenium	45 Rh Rhodium	46 Pd Palladium	47 Ag Silver	48 Cd Cadmium	49 In Indium	50 Sn Tin	51 Sb Antimony	52 Te Tellurium	53 I Iodine	54 Xe Xenon	55 Cs Caesium	56 Ba Barium	57 La Lanthanum	58 Ce Cerium	59 Pr Praseodymium	60 Nd Neodymium	61 Pm Promethium	62 Sm Samarium	63 Eu Europium	64 Gd Gadolinium	65 Tb Terbium	66 Dy Dysprosium	67 Ho Holmium	68 Er Erbium	69 Tm Thulium	70 Yb Ytterbium	71 Lu Lutetium	72 Hf Hafnium	73 Ta Tantalum	74 W Tungsten	75 Re Rhenium	76 Os Osmium	77 Ir Iridium	78 Pt Platinum	79 Au Gold	80 Hg Mercury	81 Tl Thallium	82 Pb Lead	83 Bi Bismuth	84 Po Polonium	85 At Astatine	86 Rn Radon	87 Fr Francium	88 Ra Radium	89 Ac Actinium	90 Th Thorium	91 Pa Protactinium	92 U Uranium	93 Np Neptunium	94 Pu Plutonium	95 Am Americium	96 Cm Curium	97 Bk Berkelium	98 Cf Californium	99 Es Einsteinium	100 Fm Fermium	101 Md Mendelevium	102 No Nobelium	103 Lr Lawrencium
1 H Hydrogen																																																																																																																			
11 B Boron	12 C Carbon	13 Al Aluminium	14 Si Silicon	15 P Phosphorus	16 S Sulphur	17 Cl Chlorine	18 Ar Argon	19 F Fluorine	20 Ne Neon	21 Sc Scandium	22 Ti Titanium	23 V Vanadium	24 Cr Chromium	25 Mn Manganese	26 Fe Iron	27 Co Cobalt	28 Ni Nickel	29 Cu Copper	30 Zn Zinc	31 Ga Gallium	32 Ge Germanium	33 As Arsenic	34 Se Selenium	35 Br Bromine	36 Kr Krypton	37 Rb Rubidium	38 Sr Strontium	39 Y Yttrium	40 Zr Zirconium	41 Nb Niobium	42 Mo Molybdenum	43 Tc Technetium	44 Ru Ruthenium	45 Rh Rhodium	46 Pd Palladium	47 Ag Silver	48 Cd Cadmium	49 In Indium	50 Sn Tin	51 Sb Antimony	52 Te Tellurium	53 I Iodine	54 Xe Xenon	55 Cs Caesium	56 Ba Barium	57 La Lanthanum	58 Ce Cerium	59 Pr Praseodymium	60 Nd Neodymium	61 Pm Promethium	62 Sm Samarium	63 Eu Europium	64 Gd Gadolinium	65 Tb Terbium	66 Dy Dysprosium	67 Ho Holmium	68 Er Erbium	69 Tm Thulium	70 Yb Ytterbium	71 Lu Lutetium	72 Hf Hafnium	73 Ta Tantalum	74 W Tungsten	75 Re Rhenium	76 Os Osmium	77 Ir Iridium	78 Pt Platinum	79 Au Gold	80 Hg Mercury	81 Tl Thallium	82 Pb Lead	83 Bi Bismuth	84 Po Polonium	85 At Astatine	86 Rn Radon	87 Fr Francium	88 Ra Radium	89 Ac Actinium	90 Th Thorium	91 Pa Protactinium	92 U Uranium	93 Np Neptunium	94 Pu Plutonium	95 Am Americium	96 Cm Curium	97 Bk Berkelium	98 Cf Californium	99 Es Einsteinium	100 Fm Fermium	101 Md Mendelevium	102 No Nobelium	103 Lr Lawrencium																							

| 23 **Na** Sodium | 24 **Mg** Magnesium | 39 **K** Potassium | 40 **Ca** Calcium | 45 **Sc** Scandium | 48 **Ti** Titanium | 51 **V** Vanadium | 52 **Cr** Chromium | 55 **Mn** Manganese | 56 **Fe** Iron | 59 **Co** Cobalt | 59 **Co** Cobalt | 64 **Cu** Copper | 65 **Zn** Zinc | 70 **Ga** Gallium | 73 **Ge** Germanium | 75 **As** Arsenic | 79 **Se** Selenium | 80 **Br** Bromine | 84 **Kr** Krypton | 85 **Rb** Rubidium | 88 **Sr** Strontium | 89 **Y** Yttrium | 90 **Zr** Zirconium | 91 **Nb** Niobium | 93 **Nb** Niobium | 94 **Mo** Molybdenum | 96 **Mo** Molybdenum | 101 **Ru** Ruthenium | 103 **Rh** Rhodium | 106 **Pd** Palladium | 108 **Ag** Silver | 112 **Cd** Cadmium | 115 **In** Indium | 119 **Sn** Tin | 122 **Sb** Antimony | 127 **I** Iodine | 131 **Xe** Xenon | 133 **Cs** Caesium | 137 **Ba** Barium | 139 **La** Lanthanum | 140 **Ce** Cerium | 141 **Pr** Praseodymium | 144 **Nd** Neodymium | 150 **Sm** Samarium | 152 **Eu** Europium | 157 **Gd** Gadolinium | 159 **Tb** Terbium | 162 **Dy** Dysprosium | 165 **Ho** Holmium | 167 **Er** Erbium | 169 **Tm** Thulium | 173 **Yb** Ytterbium | 175 **Lu** Lutetium |

3-71 Lanthanoid series
0-103 Actinoid series

a = relative atomic mass
 X = atomic symbol
 b = proton (atomic) number

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).