Name

www.Papa Cambridge.Com

# CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CHEMISTRY 0620/06

Paper 6 Alternative to Practical

OCTOBER/NOVEMBER 2003

1 hour 15 minutes

Candidates answer on the Question Paper. No additional Materials required.

#### **READ THESE INSTRUCTIONS FIRST**

Write your name, centre number and candidate number in the spaces provided at the top of this page. Write in dark blue or black pen in the spaces provided on the Question Paper. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all questions.


The number of marks is given in brackets [ ] at the end of each question or part question.

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

FOR EXAMINER'S USE			
1			
2			
3			
4			
5			
6			
TOTAL			

The apparatus below was used to separate ethanol from water.



- (a) Complete the empty boxes to name the pieces of apparatus.
- (b) Indicate by an arrow where heat is applied. [1]
- (c) Name this separation process.

A student extracted and investigated the orange colour in some sweets.

	The state of the s	
	3 A. D.	-
	tudent extracted and investigated the orange colour in some sweets. It is student followed these instructions:  Collect sweets, a watch glass, a beaker, eye protection and 100 cm³ of ethanol.  Crush the sweets.  Place the crushed sweets in the beaker containing 100 cm³ of ethanol.	Cal
1	Collect sweets, a watch glass, a beaker, eye protection and 100 cm <sup>3</sup> of ethanol.	1
2	Crush the sweets.	
3	Place the crushed sweets in the beaker containing 100 cm <sup>3</sup> of ethanol.	
4	Boil the mixture with the watch glass covering the beaker.	
5	Decant the liquid and concentrate it by evaporation until the colour is dark orange.	
6	Investigate which colours are present in the orange solution.	
(a)	Why should the sweets be crushed?	
		[2]
		.–.
(b)	Why should the experiment be carried out in a well-ventilated laboratory?	
	[	[1]
(c)	State <b>one</b> safety precaution that the student should have taken, other than carrying of the experiment in a well-ventilated laboratory and using eye protection.	out
	[	[1]
(d)	State the purpose of the watch glass.	
	г	[1]
		. ' ]
(e)	Explain the term <i>decant</i> .	
	]	[1]
(f)	Describe how the student could carry out instruction 6. You may draw a diagram in the space below to help you answer the question.	he

For Examiner's

3 A student investigated the speed of reaction between aqueous potassium brome potassium iodide solution.

A burette was filled up to the 0.0 cm<sup>3</sup> mark with aqueous potassium iodide.

To each of 5 test-tubes was added 6 cm<sup>3</sup> of aqueous potassium iodide to be used in the 5 following experiments.

## Experiment 1

By using a measuring cylinder 12 cm<sup>3</sup> of aqueous potassium bromate was poured into a small beaker. To this solution was added 4 cm<sup>3</sup> of water, 2 cm<sup>3</sup> of hydrochloric acid, 5 cm<sup>3</sup> of starch solution and 1 cm<sup>3</sup> of sodium thiosulphate solution.

The beaker was placed on a cross drawn on a piece of paper.

From one of the test-tubes 6 cm<sup>3</sup> of aqueous potassium iodide was added to the mixture in the beaker and the timer started. A dark blue colour formed. The timer was stopped when the cross on the paper could not be seen.

Use the stop clock diagram to record the time in the table.

### Experiment 2

By using a measuring cylinder  $10\,\mathrm{cm}^3$  of potassium bromate solution was poured into a beaker. The instructions were repeated exactly as given for Experiment 1, but  $6\,\mathrm{cm}^3$  of water was added to the beaker.

Use the diagram to record the time in the table.

#### Experiments 3, 4 and 5

Experiment 1 was repeated using the volumes of aqueous potassium bromate and water specified in the table of results. Record the times in the table.


## Table of results

Table of re	esults		5	MMN. P.O.	For Examiner's Use
Experiment	volu	me	clock diagram	time/s	Dridge
1	potassium bromate/cm <sup>3</sup>	water/cm <sup>3</sup>	minutes 0 seconds  0  15  15  15		0,00
2	10	6	minutes 0 seconds  0 45 15 10 30		
3	8	8	minutes 0 seconds  0 45 15 15 15		
4	6	10	minutes 0 seconds  0 45 15 5 15		
5	4	12	minutes 0 seconds  0 45 15 5 15		

(a) Plot the results on the grid. Draw a smooth line graph.

(ii)

Explain why this experiment is the quickest.




		7 State <b>two</b> possible sources of error in the experiments. 1	
		7	For Examiner's
d)	(i)	State <b>two</b> possible sources of error in the experiments.	Use
		1	Mario
			36.CC
		2	TH.
	(ii)	Suggest <b>two</b> improvements to reduce the sources of error in the experiments.	
		1	
		2	
		T 4.1	

		8 WMM. Da
(		was analysed. Substance X was an iron(N X and some of the observations are in the follotable.  Observations  dark yellow
	Tests	Observations
(a)	Colour of solution X	dark yellow
(b)	(i) Drops of aqueous sodium hydroxide were added to about $2 \text{ cm}^3$ of the solution. Excess aqueous sodium hydroxide was	
	added to the test-tube.	[3]
	(ii) The mixture was heated. The gas given off was tested with damp indicator paper.	pungent smell indicator turned blue, pH 10
c)	Experiment <b>(b)(i)</b> was repeated using aqueous ammonia instead of aqueous sodium hydroxide.	
		[2]
)	To about 2 cm <sup>3</sup> of solution <b>X</b> was added dilute sulphuric acid. Two pieces of zinc were added. The	lighted splint
	mixture was heated and the gas given off tested.	popped
	After 10 minutes the mixture was filtered and test <b>(b)(i)</b> was repeated.	green precipitate insoluble in excess
e)	A few drops of hydrochloric acid were added to about 2 cm <sup>3</sup> of solution <b>X</b> . About 1 cm <sup>3</sup> of barium chloride solution was added to the	
	mixture.	white precipitate

		Name the gas given off in (d).	For
(f)	(i)	Name the gas given off in (d).	Examiner's Use
	(ii)	What type of chemical reaction occurs in (d). Explain your answer.	Ambridge Com
			[3]
(g)	Wh	at conclusions can you draw about the anion and the other cation in substance <b>X</b>	<u>'</u> ?
	anio	on	
	cati	ion	[2]

Ammonia is produced when aqueous sodium hydroxide is warmed with am 5 sulphate. Ammonia is less dense than air and very soluble in water. The apparatus b was used to prepare a sample of dry ammonia gas.



(a)	Name substance <b>C</b> [1]
(b)	Name substance <b>D</b> [1]
(c)	What necessary piece of equipment is missing in the diagram?
	[1]
(d)	Suggest why concentrated sulphuric acid should <b>not</b> be used to dry ammonia.
	[1]
(e)	There are two other mistakes in the apparatus shown in the diagram. Identify and explain these mistakes.
	mistake 1
	explanation
	mistake 2
	explanation [4]

I	For
	Examiner's

www.PapaCambridge.com Sulphur dioxide gas is a common pollutant formed when fossil fuels burn in air. 6 dioxide can be detected by using an acidic solution of potassium dichromate(VI). dichromate solution changes colour from orange to green when a certain amount of sulph dioxide has reacted with it.

Plan an experiment to investigate which of three different samples of coal produces sulphur dioxide.	
	[6]

12

**BLANK PAGE** 

www.PapaCambridge.com