

MARK SCHEME for the May/June 2007 question paper

0620 CHEMISTRY

0620/03

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2007 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme	Syllabus
	IGCSE – May/June 2007	0620

An incorrectly written symbol, e.g. NA or CL, should be penalised once in the paper.

- 1 (a) (i) coal or coke or peat
NOT wood or charcoal
- (ii) natural gas or methane or propane or butane or petroleum gases or calor gas or refinery gas [1]
- (b) (i) petrol or gasoline
 paraffin or kerosene
 diesel
 aviation fuel or jet fuel
 fuel oil
 heavy fuel oil
 heating oil
 Any **TWO** [2]
NOT a named alkane e.g. octane
- (ii) waxes or grease or lubricants or polishes or bitumen (tar, asphalt) or naphtha [2]
 Any **TWO** from the primary or secondary distillation of petroleum
- (iii) (liquid) air or ethanol and water or alkenes (made by cracking) or Noble Gases [1]

[Total: 7]

- 2 good [1]
 named example e.g. sodium chloride [1]
ACCEPT correct formula
- silica or silicon(IV) oxide or sand or silicon oxide
 named polymer only **TWO** elements [1]
- electrons [1] and positive ions [1] [2]
 good [1]

[Total: 6]

- 3 (i) method C [1]
 sulphuric acid (allow if given in equation) [1]
 zinc oxide + sulphuric acid = zinc sulphate + water [1]
- (ii) method A [1]
 hydrochloric acid [1]
 $\text{KOH} + \text{HCl} = \text{KCl} + \text{H}_2\text{O}$ [1]
- (iii) method B [1]
 potassium iodide or any soluble iodide [1]
 $\text{Pb}^{2+} + 2\text{I}^- = \text{PbI}_2$ accept a correct equation even if soluble iodide is wrong [2]
 Not balanced - $\text{Pb}^{2+} + \text{I}^- = \text{PbI}_2$ ONLY [1]

[Total: 10]

Page 3	Mark Scheme	Syllabus
	IGCSE – May/June 2007	0620

- 4 (a) (i) BaO
- (ii) B₂O₃
- (b) (i) S²⁻ [1]
- (ii) Ga³⁺ [1]
- (c) NCl₃ [1]
COND 8e (1bp and 3nbp) around each chlorine [1]
 8e (3bp and 1nbp) around nitrogen [1]
- (d) (i) ignore a correct chemical property in (i)
 vanadium harder
 vanadium higher melting point **or** boiling point
 vanadium higher density
ANY TWO [2]
OR corresponding statements for potassium
 NB has to be comparison
- (ii) ignore a correct physical property in (ii)
 potassium more reactive or example of different reactivities-
 potassium reacts with cold water, vanadium does not.
 potassium one oxidation state, vanadium more than one
 vanadium coloured compounds, potassium white **or** colourless
 vanadium and its compounds catalysts, not potassium
ANY TWO [2]
 NB has to be comment about both elements
- (e) (i) fluorine gas [1]
 astatine solid [1]
- (ii) both have valency of one
 both can react with other elements to form halides
 both are oxidants
 or any correct Chemistry – they both form acidic hydrides
 both have diatomic molecules
 both accept one electron **or** form ion X⁻
 both have seven valency electrons
 both react with non-metals to form covalent compounds
 both react with metals to form ionic compounds
 both form acidic oxides
NOT have a valency of 7
ANY TWO [2]

[Total: 15]

Page 4	Mark Scheme	Syllabus
	IGCSE – May/June 2007	0620

- 5 (a) (i) air would react (with the magnesium **or** titanium)
OR argon would not react (with the metals)
NOT argon is inert
- (ii) any metal higher than magnesium in reactivity series [1]
- (iii) add water (to dissolve salt) [1]
 filter **or** centrifuge [1]
- (b) (i) electron loss [1]
- (ii) hydrogen [1]
- (iii) oxygen [1]
 chlorine [1]
- (iv) it cannot lose electrons (because) [1]
 it receives electrons (from the battery) [1]
- OR** reduction occurs at the cathode [1]
 oxidation at the anode (not cathode) [1]
- OR** electrons are “pushed” to rig [1]
 preventing it from being oxidised [1]
- for comments of the type – rusting needs oxygen, it is formed on titanium not iron **ONLY** [1]
NOT the idea that titanium is more reactive etc
- (v) **SET 1**
 sacrificial protection is a cell
 does not need electricity
 cathodic protection is electrolysis
 cathodic protection needs electricity
- SET 2**
 sacrificial protection needs a more reactive metal (in contact with iron or steel)
 this metal corrodes instead of steel
 cathodic protection needs an inert electrode accept unreactive or less reactive metal as
 an electrode
 has to be **ONE** comment from each set [2]
 all comments about oxide layers and coating are neutral

[Total: 12]

Page 5	Mark Scheme	Syllabus
	IGCSE – May/June 2007	0620

- 6 (a) alumina **or** aluminium oxide
sodium aluminate
iron(III) oxide
filtration **or** centrifuge NOT conditional
- (b) from left to right:
carbon cathode **or** carbon negative electrode [1]
900 to 1000°C [1]
aluminium [1]
cryolite [1]
- (c) (i) $Al^{3+} + 3e = Al$ [2]
not balanced [1]
 $Al^{3+}(aq) = 0$
- (ii) oxygen is formed **NOT** oxide [1]
reacts with carbon anode [1]
- (d) (i) low density **or** light **or** resistant to corrosion
accept strength/weight ratio **or** alloys are strong [1]
strong on its own is neutral
- (ii) not attacked **or** corroded **or** unreactive
oxide layer
easily shaped **or** malleable **or** ductile
any **TWO** [2]
- (iii) for strength **or** so it does not break **or** does not sag **or** can have pylons further apart [1]
NOT steel is a better conductor
NOT aluminium protects steel from rusting

[Total: 16]

Page 6	Mark Scheme	Syllabus
	IGCSE – May/June 2007	0620

- 7 (a) butanol
no number needed but if one is given it has to be 1
- structural formula (all bonds shown)
accept –OH **NOT** –HO
- ethanoic acid [1]
structural formula (all bonds shown) [1]
accept –OH **NOT** –HO
no conseq marking
if all bonds are not shown (CH₃–CH₂–), penalise once
- (b) (i) must have correct ester linkage [1]
COND continuation and a group on either side of the ester group [1]
Accept –COO–
- (ii) accept any sensible suggestion [1]
ropes, clothing, bottles, packaging, bags
- (c) (i) 8 [1]
- (ii) double bond becomes single and 4 bonds per carbon atom [1]
COND a bromine atom on each carbon [1]
C₂H₄Br₂ ONLY [1]
accept a structural formula with hydrogen atoms
- (iii) corn oil [1]
- (d) 100g of fat react with 86.2g of iodine
884g of fat react with **762** g of iodine [1]
limit 762 x 2
one mole of fat reacts with 762/254 moles of iodine molecules
one mole of fat reacts with **3** moles of iodine molecules [1]
- number of double bonds in one molecule of fat is **3** [1]
limit 6
consequential marking allowed provided the number of double bonds is an integer.

[Total: 14]