WWW. Palls

## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

**International General Certificate of Secondary Education** 

## MARK SCHEME for the October/November 2009 question paper for the guidance of teachers

## **0620 CHEMISTRY**

0620/32

Paper 32 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	- Per
	IGCSE – October/November 2009	0620	120

## **GENERAL INSTRUCTIONS FOR MARKING**

- Error carried forward may be allowed in calculations. This will be discussed in the scheme. This is not applied when the candidate has inserted incorrect integers or when answer is physically impossible.
- COND the award of this/these mark(s) is conditional upon a previous mark being awarded.
   Example Is the reaction exothermic or endothermic? Give a reason for your choice.
   Mark scheme exothermic [1]

**COND** a correct reason given [1]. This mark can only be awarded if the candidate has recognised that the reaction is exothermic.

- When the name of a chemical is demanded by the question, a correct formula is usually acceptable. When the formula is asked for, the name is not acceptable.
- When a word equation is required a **correct** symbol equation is usually acceptable. If an equation is requested then a word equation is not usually acceptable.
- An incorrectly written symbol, e.g. NA or CL, should be penalised once in a question.
- In the mark scheme if a word **or** phrase is underlined it (**or** an equivalent) is required for the award of the mark.

(.....) is used to denote material that is not specifically required.

- OR designates alternative and independent ways of gaining the marks for the question.
   or indicates different ways of gaining the same mark.
- Unusual responses which include correct Chemistry which answer the question should always be rewarded even if they are not mentioned in the marking scheme.

Page 3	Mark Scheme: Teachers' version	Syllabus	'A er
	IGCSE – October/November 2009	0620	100

- (a) (i) argon or krypton or helium Accept xenon and radon even though percentages are very small **NOT** hydrogen
  - (ii) water and carbon dioxide

example

- (b) (i) carbon monoxide or lead compounds or CFCs or methane or particulates or unburnt hydrocarbons or ozone [1]
  - (ii) burn a fossil fuel [1] that contains sulfur [1]
  - (iii) at high temperature or inside engine [1] nitrogen and oxygen (from the air) react [1]
- (c) liquid air [1] [1] fractional distillation

[Total: 10]

[Total: 9]

- 2 (a) pH < 7[1] [1]
  - pH > 7[1] [1] example

**NOT** amphoteric oxides Be, A*l*, Zn, Pb, Sn etc.

pH = 7[1] example H<sub>2</sub>O, CO, NO [1]

the two marks are not linked, mark each independently **NOT** amphoteric oxides Be, Al, Zn, Pb, Sn etc.

- (b) (i) shows both basic and acidic properties [1]
  - (ii) acidic reacts with sodium hydroxide only [1] amphoteric reacts with both reagents [1]
    - OR only amphoteric oxide reacts with hydrochloric acid [2]

3 (a) (i) heat/roast/burn in air [1] need both points for mark

(ii)  $ZnO + C \rightarrow Zn + CO$ [2] or  $2ZnO + C \rightarrow 2Zn + CO_2$ unbalanced ONLY [1]

Page 4		ı	Mark Scheme: Teachers' version	Syllabus	S Or	
Page 4		•	IGCSE – October/November 2009	Syllabus 0620	OB.	
	(b)	(b) zinc is more reactive it loses electrons and forms ions in preference to iron zinc corrodes not iron NOT zinc rusts		Papa Cambridge		
	OR zinc loses electrons and forms ions the electrons move on to the iron the iron cannot be oxidised or it cannot rust or it cannot lose electrons CREDIT correct Chemistry that includes the above ideas		[1] [1] [1]			
	(c)	(i)		atoms change into ions, (the zinc dissolves) $per(II)$ ions change into atoms, (becomes plated with co	opper)	[1] [1]
		(ii)	ions elec	trons		[1] [1]
						[Total: 10]
4	<ul> <li>(a) diffusion</li> <li>different M<sub>r</sub> or ozone molecules heavier than oxygen molecules</li> <li>or different densities or oxygen molecules move faster than ozone molecules</li> <li>NOT oxygen is lighter or ozone heavier</li> </ul>		[1] [1]			
				ional distillation e different boiling points		[1] [1]
	(b)	(i)		colourless (solution) rown (solution)		[1] [1]
		(ii)	I <sup>-</sup> los	ses electrons (it is oxidised)		[1]
		(iii)	-	are accepted by ozone zone is an electron acceptor		[1]
	(c)	(i)	sulfu all <b>th</b>	on dioxide ır dioxide		[2]
		(ii)	CON	ect structural skeleton  ID 4bp around both carbon atoms  and 2nbp around sulfur atom		[1] [1] [1]

[Total: 11]

Page 5	Mark Scheme: Teachers' version	Syllabus	er
	IGCSE – October/November 2009	0620	100

5	(a) (i)	strong hard light or low density high melting point or high fixed points Accept high strength to weight ratio for [2] it includes marks 1 and 3 any THREE	Cambride [3]
	(ii)	diagram 1 four silicons around one carbon diagram 2 four carbons around one silicon either diagram looks <b>or</b> stated to be tetrahedral "tetrahedral" scores mark even if diagram does not look tetrahedral independent marking of three points	[1] [1] [1]
	ead	gram to include ch germanium atom bonded 4 oxygen atoms ch oxygen to 2 germanium atoms	[1] [1]
	(c) (i)	structural formula of Ge <sub>3</sub> H <sub>8</sub> all bonds shown	[1]
	(ii)	germanium oxide water	[1] [1]
			[Total: 11]
6	(a) (i)	LISA or Toyon or Louisiana Japan	
	(a) (i)	USA or Texas or Louisiana, Japan volcanoes, natural gas, petroleum	[1]
	(a) (i) (ii)		[1]
		volcanoes, natural gas, petroleum  bleach for wood pulp/cloth/straw or preserve food or sterilising or making wine or fumigant or refrigerant	
	(ii)	volcanoes, natural gas, petroleum bleach for wood pulp/cloth/straw <b>or</b> preserve food <b>or</b> sterilising <b>or</b> making wine <b>or</b> fumigant <b>or</b> refrigerant <b>Accept</b> making paper vanadium(V) oxide or vanadium oxide or vanadium pentoxide or $V_2O_5$	[1]
	(ii)	volcanoes, natural gas, petroleum bleach for wood pulp/cloth/straw <b>or</b> preserve food <b>or</b> sterilising <b>or</b> making wine <b>or</b> fumigant <b>or</b> refrigerant <b>Accept</b> making paper vanadium(V) oxide or vanadium oxide or vanadium pentoxide or $V_2O_5$ NB oxidation state not essential but if given has to be (V)	[1]
	(ii) (iii) (iv)	volcanoes, natural gas, petroleum bleach for wood pulp/cloth/straw <b>or</b> preserve food <b>or</b> sterilising <b>or</b> making wine <b>or</b> fumigant <b>or</b> refrigerant <b>Accept</b> making paper vanadium(V) oxide or vanadium oxide or vanadium pentoxide or $V_2O_5$ NB oxidation state not essential but if given has to be (V) rate too slow <b>or</b> rate not economic	[1] [1]
	(ii) (iii) (iv) (v)	bleach for wood pulp/cloth/straw <b>or</b> preserve food <b>or</b> sterilising <b>or</b> making wine <b>or</b> fumigant <b>or</b> refrigerant <b>Accept</b> making paper  vanadium(V) oxide or vanadium oxide or vanadium pentoxide or V <sub>2</sub> O <sub>5</sub> NB oxidation state not essential but if given has to be (V)  rate too slow <b>or</b> rate not economic  reaction too violent <b>or</b> forms a mist	[1] [1] [1] [1]

		2.
Page 6	Mark Scheme: Teachers' version	Syllabus er
	IGCSE – October/November 2009	0620
		C

(c) number of moles of FeSO<sub>4</sub> used = 12.16/152 = 0.08\* number of moles of Fe<sub>2</sub>O<sub>3</sub> formed = 0.04 mass of one mole of Fe<sub>2</sub>O<sub>3</sub> =  $160 \, \mathrm{g}$  mass of iron(III) oxide formed =  $0.04 \times 160 = 6.4 \, \mathrm{g}$  number of moles of gases formed =  $0.08 \times 24 = 1.92 \, \mathrm{dm}^3$  volume of sulfur trioxide formed =  $0.08 \times 24 = 1.92 \, \mathrm{dm}^3$ 

(d) have same boiling point

[1] COM

If mass of iron(III) oxide greater than 12 g, then only marks 1 and 2 available

Apply  $\mathbf{ecf}$  to number of moles of  $\mathrm{FeSO_4}^*$  when calculating volume of sulfur trioxide. Do not apply  $\mathbf{ecf}$  to integers

[Total: 16] 7 [1] (a) (i) heat [1] catalyst (ii) equation that gives: alkene + alkane or alkene + alkene + hydrogen [1] a correct and balanced equation for the cracking of decane, C<sub>10</sub>H<sub>22</sub> but not but-1-ene [1] (iii) water or steam [1] [2] **(b) (i)**  $C_4H_9OH + 6O_2 \rightarrow 4CO_2 + 5H_2O$ If only error is balancing the oxygen atoms [1] [2] (ii) butanol + propanoic acid → butyl propanoate + water correct products or reactants ONLY [1] (c) (i) correct structural formulae [1] each [2] penalise once for CH<sub>3</sub> type diagrams For  $C_3H_8O$  [0] (ii) to conserve petroleum or reduce greenhouse effect [1]

[Total: 13]

[1]