	UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
	International General Certificate of Secondary Education
ENTRE IUMBER	CANDIDATE NUMBER
HEMISTRY	0620/61
aper 6 Alterna	tive to Practical October/November 2010
	1 hour
Candidates and	wer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

899

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Exam	iner's Use
1	
2	
3	
4	
5	
6	
7	
Total	

This document consists of 13 printed pages and 3 blank pages.





3 The following instructions were used to prepare magnesium sulfate crystals, MgSO ₄ .
The following instructions were used to prepare magnesium sulfate crystals, MgSO ₄ .
Step 1 Measure 50 cm ³ of dilute sulfuric acid into a beaker and warm the solution.
Step 2 Using a spatula, add some magnesium oxide and stir the mixture. Continuadding the magnesium oxide until excess is present.
Step 3 Separate the excess magnesium oxide from the solution of magnesium sulfate.
Step 4 Heat the solution until crystals form. Obtain the crystals and dry them.
(a) Why is the sulfuric acid warmed?
[
(b) How would you know when excess magnesium oxide is present in Step 2?
[
(c) What method is used in Step 3?
[
(d) Why must care be taken when drying the crystals in Step 4?
[
(e) Explain how the method would differ if magnesium carbonate was used instead magnesium oxide.
[:
[Total: 0

www.PapaCambridge.com 3 A student carried out an experiment to investigate the speed of reaction between thiosulfate solution and dilute nitric acid. Sulfur is formed during this reaction and the mixture turns cloudy.

Experiment 1

Using a measuring cylinder, 100 cm³ of sodium thiosulfate solution was poured into a 250 cm³ beaker. The beaker was placed on a cross drawn on a piece of paper. 20 cm³ of dilute nitric acid was added to the beaker and the timer started.



The time until the cross could not be seen was taken. The time was recorded in the table.

Experiment 1 was repeated using different volumes of sodium thiosulfate as shown in the table.

All experiments were carried out at 20 °C.

Table of results

experiment	volume of sodium thiosulfate solution/cm ³	volume of water/cm ³	time for cross to disappear/s
1	100	0	10
2	80	20	12
3	40	60	24
4	20	80	51
5	10	90	98

(a) Why was the total volume of solution kept constant?

(b) In Experiment 2, which is the last liquid to be added to the beaker?[1]

4





A student investigated the solubility of salt A in water at various temperatures. 4 Five experiments were carried out.

Experiment 1

The student was provided with a boiling tube containing 12 g of salt A.

www.papaCambridge.com A burette was filled with distilled water and 10.0 cm³ of water was added to the boiling tube. The mixture of salt **A** and water was heated until all of the solid had dissolved.

The boiling tube was removed from the heat and the solution was stirred with a thermometer and allowed to cool.

The temperature at which crystals first appeared was measured.

Use the thermometer diagram to record the temperature in the table of results.



The boiling tube and contents were kept for the next four experiments.

Experiment 2

From the burette, 1.0 cm³ more of water was added into the boiling tube and contents from Experiment 1.

The experiment was repeated exactly as before to find the temperature at which crystals first appeared.

The boiling tube was dipped for short periods of time in a beaker of cold water to speed up the cooling.

Record, in the table of results, the total volume of water in the boiling tube. Use the thermometer diagram to record the temperature at which crystals first appeared.



7

Experiment 3

From the burette 1.0 cm³ more of water was added into the boiling tube and contents from Experiment 2. The experiment was repeated exactly as before.

www.papacambridge.com Record, in the table of results, the total volume of water used. Use the thermometer diagram to record the temperature at which crystals first appeared.



This procedure was continued for Experiments 4 and 5 with two more successive additions of 1.0 cm³ of water. Note all the results in the table.





diagram for Experiment 4

diagram for Experiment 5

Table of results

experiment	total volume of water/cm ³	temperature at which crystals first appeared/°C
1	10.0	
2		
3		
4		
5		

[5]



	10 44444. Papa
(d)	Suggest, with a reason, how the results would be different if 6 g of salt A we instead of 12 g of salt A.
	[2]
(e)	Salt B is more soluble in water than salt A . Sketch on the grid the graph you would expect for salt B . Label this graph. [2]
(f)	Explain one improvement you could make to the experimental procedure to obtain more accurate results in this investigation.
	improvement
	explanation[2]
	[Total: 20]

	1	1 334. Dab
5 Two The	o different solutions, X and Y , were analy e tests on the solutions, and some of the	sed. X was copper sulfate solution. observations, are in the following table.
Cor	nplete the observations in the table.	
	tests	observations
tests or	n solution X	
(a) (i)	Appearance of solution X .	
(ii)	To a little of solution X , aqueous sodium hydroxide was added.	
(iii)	To a little of solution X , aqueous ammonia was added drop by drop and shaken.	[1]
	Excess aqueous ammonia solution was then added to the test-tube.	
tests or	n solution Y	
(b) (i)	A little of solution Y was tested with Universal Indicator paper. The pH was recorded.	pH1
(ii)	To about 3 cm^3 of solution Y a few drops of dilute hydrochloric acid and then aqueous barium chloride was added.	white precipitate

......[2]

[Total: 8]

- 6 A concentrated solution of sodium chloride was electrolysed using the apparatus belt of miners positive electrode electrode electrode aqueous sodium chloride and Universal Indicator
 One observation noted was that the Universal Indicator turned purple at the negative electrode.
 (a) What observation would be made at both electrodes?
 - (b) Why did the indicator turn purple at the negative electrode?

......[1]

(c) (i) Name the product formed at the positive electrode.
 (ii) Suggest the effect of this product on the Universal Indicator.
 [1]

[Total: 4]

		13
7	En	umbers identify chemicals which are added to foods.
-	(a)	E210 is benzoic acid. How could you show that a solution of benzoic acid is a wea acid? test
		result[2
	(b)	E211 is sodium benzoate. Name a suitable substance that would react with a solution of benzoic acid to form sodium benzoate.
		[1
	(c)	E110 is Sunset yellow. Outline a method you could use to show the presence of E110 in a food colouring. A space has been left if you want to draw a diagram to help you answer the question.
		[4
		[Total: 7



BLANK PAGE



BLANK PAGE



BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of