UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

## www.papaCambridge.com MARK SCHEME for the May/June 2010 question paper

## for the guidance of teachers

## **0607 CAMBRIDGE INTERNATIONAL MATHEMATICS**


0607/04 Paper 4 (Extended), maximum raw mark 120

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



- soi seen or implied
- www without wrong working

1	(a)	7.60	B2	[2]	If B0, SC1 for 12.4
	(b)	3.3 (33)	B2	[2]	If B0, SC1 for 3.2 (3.22 – 3.23) or M1 for $\frac{0.05}{1.5} \times 100$ oe
	(c)	0.75	B2	[2]	If B0, M1 for 0.84 ÷ 1.12 oe
	(d)	5 www 3		[3]	B2 for $4.937 - 4.938$ or $\frac{\log(\frac{4}{3})}{\log 1.06}$ or M1 for using $0.75 \times 1.06^n$ or $1 \div 1.06^n$ oe, <i>n</i> integer >1
2	(a)	$ x^{3} - x^{2} + x + x^{2} - x + 1 = x^{3} + 1 $	M1	[2]	Must see 6 correct terms, condoning one sign error Correct conclusion with no errors or omissions
	(b)	Use of $b^2 - 4ac$ (may be in formula) or sketch of U – shaped parabola or $(x - 0.5)^2 = k$	M1	[3]	
		For $\sqrt{-3}$ or for not touching $x$ – in sketch or $k = -0.75$	A1		
		Correct conclusion (e.g. not real or square root of a negative number or graph does not touch <i>x</i> -axis)	R1		Dependent on M1 A1
	(c) (i)	9	B1	[1]	
	(ii)	0	B1	[1]	
	(iii)	$\sqrt[3]{x-1}$ final answer	B3	[3]	If B0, M1 for $y - 1 = x^3$ (correct rearrangement) M1 for $\sqrt[3]{y-1} = x$ (correct cube root) M1 for interchanging <i>x</i> and <i>y</i> (this may be done first and other 2 M's follow)
	(iv)	28 cao	B1	[1]	

Page	e 3 Mark Scheme: Teac IGCSE – May/J	Syllabus of er 0607	
			aCan
3 (a)	{3, 6, 9, 12} {1, 2, 3, 5, 6, 10} {6, 7, 8, 9, 10, 11}	B1 B1 B1 [3]	Syllabus 0607 12 elements must be seen, each
(b)	2 12 3 1 5 6 10 7 8 11 4	B2ft [2]	12 elements must be seen, each only once. Correct or ft their (a) If B0, B1 for 6 or 7 regions correct o ft their (a)
(c) (i)	3, 6	B1ft [1]	Correct or ft their Venn diagram
(ii)	3, 6, 7, 8, 9, 10, 11, 12	B1ft [1]	Correct or ft their Venn diagram
(iii)	3, 6, 10	B1ft [1]	Correct or ft their Venn diagram
(iv)	4, 7, 8, 9, 11, 12	B1ft [1]	Correct or ft their Venn diagram
(v)	11 cao	B1 [1]	
4 (a)	1005 www 3	M1 M1	M1 for at least 3 correct mid-interval values seen then (985, 995, 1002.5, 1007.5, 1015, 1030) M1 for evidence of using $\Sigma fx \div 100$ with values of <i>m</i> in the intervals (3940 + 9950 + 50125 +
		A1 [3]	20150 + 8120 + 8240 = 100525)
(b) (i)	0.4, 1, 10, 4, 0.8, 0.4	B3 [3]	If B0, B2 for 4 or 5 correct, B1 for 2 or 3 correct
(ii)	Accurate histogram	M1 A1 A2ft [4]	<ul> <li>6 histogram rectangles with at least 4 correct widths</li> <li>6 correct widths</li> <li>6 correct heights, A1ft for 4 or 5 correct heights</li> </ul>

		1		32
	Page 4		Teachers' version ay/June 2010	Syllabus er 0607
_	<i>(</i> )		Di	Can
5	(a)	Three reasonable branches	B1 B1 B1 [3]	Syllabus 0607erMust be above x-axis and alway increasing roughly towards $x = -1$ Inverted U shape below x-axis and roughly between $x = -1$ and $x = 3$ Must be above x-axis and always decreasing roughly from $x = 3$
	(b)	x = -1, x = 3, y = 0	B1B1B1 [3]	
	(c)	(1, -0.25)	B1B1[2]	If B0, allow SC1 for <b>both</b> inexact values of correct answer e.g. (1.0000011, - 0.24999)
	(d)	$(x \in R), x \neq -1, x \neq 3$ oe	B1B1ft [2]	B1 for $x \neq -1$ B1 for $x \neq 3$ oe ft (b) or correct. isw limits from grid
		$y \le -0.25, y > 0$	B1B1ft [2]	B1 for $y \le -0.25$ B1 for $y > 0$ (f(x) OK for y and condone x for y ft (c) for their $-0.25$ and (b) for their 0, or correct.
	(e) (i)	2	B1 [1]	
	(ii)	4	B1 [1]	
6	(a)	250	B2 [2]	If B0, M1 for 100 ÷ 5.6 × 14
	(b) (i)	20	B2 [2]	If B0, M1 for 72 ÷ 3.6 oe
	(ii)	0.225 ft	B2ft [2]	ft 4.5 ÷ their (b)(i). If B0, M1 for 4.5 ÷ their (b)(i)
7	(a)	$\frac{x}{5}$	B1 [1]	
	(b) (i)	$\frac{x+13}{4}$	B1 [1]	
	(ii)	$\frac{x}{5} + \frac{x+13}{4} = 46$	B1ft [1]	ft their (a) and (b)(i) if fractions of the form $\frac{x}{l}, \frac{x+p}{q}, q \neq l$ neither = 1
	(iii)	95	B3ft dep [3]	ft dependent on B1 in (ii) but if $p = 0$ , SC2 for 102.(22) If B0, M1for " $4x$ "+" $5x$ "+" $65$ " o.e. (or better) then M1 (dependent) for " $4x$ "+" $5x$ "+" $65$ "="920" o.e. (or better) these two M's may still be over common denominator
	(c)	4.41 ft	B2ft [2]	ft (2 × their (b)(iii) + 13) ÷ 46 If B0, M1 for (2 × their (b)(iii) + 13) ÷ 46 oe, 4.4, 4.413 imply M1

Pa	ige 5	<b>j</b>	Mark Scheme: Teac IGCSE – May/Ju		on		Syllabus 0607
			IGCSE – May/J				0007 780
(a)		Positi	ve	B1	[1]		Syllabus 0607 0607 0607 0607 0607 0607
(b)	(i)	9.6		B1	[1]		
(	(ii)	11		B1	[1]		
(i	iii)	1.2(0)	r + 1.5(0)	B1B	1 [2]	<i>mr</i> + zero.	or $1.2(0)r (1.198) + c$ , or B1 for $1.5(0) (1.497)$ with $c, m$ not Allow $x$ for $r$ . for $1.2(0)$ and $1.5(0)$ seen
) (a)		1190	(1193) cao www3	M2 A1	[3]	$\frac{180}{\sin 8}$ or M $\frac{AK}{\sin 8}$ SC2 f	
(b)		1199	– 1200 cao www 3	M1 A2	[3]	A1 fo SC2 f	for $2^{2} + 770^{2} - 2 \times 1410 \times 770 \cos 58.$ for 1 439 990.8 for correct answer by other methor e of other triangle if working seen
(c)		56850	00 – 569400 cao www 3	M2 A1	[3]	82.5 - o.e.	for $0.5 \times 180 \times$ their <b>(a)</b> $\times \sin(180 - 8.6) + 0.5 \times 1410 \times 770 \times \sin 58$ 1 for area of one triangle
(d)		1490	– 1500 cao www 5	B2 M1 A2	[5]	angle If this metho M1 fo <i>AT</i> or e.g. 1 cos (8	or angle $TDK = 32.96 - 33.09$ or to be $DKT = 88.36 - 88.61$ is B0, then M1 for correct full od for finding one of these angles for correct explicit expression for r $AT^2$ $80^2 + 1410^2 - 2 \times 180 \times 1410 \times$ 82.5 + their angle TDK) or 2 235 000 to 2 240 000
(e)		28.2	cm	B2	[2]	If B0	, SC1 for figures 282

Page		Mark Scheme: Teachers' version						
IGCSE – May/June 2010 0607								
10 (a)	Prism	B1 [1]	Syllabus         er           0607         0607           If B0, M1 for 0.5 (3 + 4) × 1.5 oe         end           ft their (b)(i) × 154.4. If B0 then M2 for         their (b)(i) × 154.4. If B0 then M2 for					
(b) (i)	5.25	B2 [2]	If B0, M1 for $0.5 (3 + 4) \times 1.5$ oe					
(ii)	810.6 (or 811)	B3ft [3]	ft their (b)(i) $\times$ 154.4. If B0 then M2 for their (b)(i) $\times$ 8 $\times$ 19.3 or M1 for their (b)(i) $\times$ 8.					
(iii)	91.78 – 91.8 cao www 4	M3 A1 [4]	M1 for $(BC^2) = 1.5^2 + 0.5^2$ (implied by 1.58 or 2.5) may be on diagram or near to diagram M1 for 2 × their (b)(ii) M1 for (3 + 4 + 2 × their BC) × 8 (BC must be from Pythagoras or trigonometry)					
(c)	24 ft	M1 A2ft[3ft]	ft 20 000 ÷ their (b)(ii) M1 for mass of box ÷ mass of one bar (implied figs 2466 to 2467) A1 for 24.66 – 24.67 or 24.6 or 24.7 (i.e. 1 dp) ft Answers less than 1 can only have M1					
11 (a)	$\frac{1}{6}$ oe	B1 [1]	Penalty of –1 for 2 sf decimals or percentages. Do not accept ratio or worded forms.					
(b) (i)	2 then 4 branch tree diagram 3 pairs of correct probabilities $\frac{5}{6}, \frac{1}{6}$ an $\frac{7}{8}, \frac{1}{8}$ and $\frac{1}{4}, \frac{3}{4}$ on a 2 then 4 branch tree diagram	DI	B1 each correct pair with correct orientation If incorrect orientation check labelling for the three pairs					
(ii)	$\frac{37}{48}(0.7708)$ cao	B3 [3]	If B0, M2 for $\frac{5}{6} \times \frac{7}{8} + \frac{1}{6} \times \frac{1}{4}$ or M1 for one of the above products.					
(iii)	0.00063 cao	B2 [2]	If B0 M1 for $(1 - \text{their } (\mathbf{h})(\mathbf{ii}))^5$ or					
			$\left(\frac{5}{6} \times \frac{1}{8} + \frac{1}{6} \times \frac{3}{4}\right)^5 \text{ ft their tree oe}$ 0.000632 implies M1					
(iv)	148	B1ft [1]	ft their (b)(ii) × 192 Allow decimals or integers (rounded or truncated)					

Page 7	Mark Scheme: Teachers' version IGCSE – May/June 2010			Syllabus 0607 Babac
	IGUSE – May/.	June 2010	0607 732	
	$\frac{192}{3 \times 2^n}$ oe	B1 B2		.g. $6 \times 2^{n-1}$ f B0, B1 for $2^k$ , $k \in \mathbb{Z}$ in final answer
(b)	24 $n^2 - 2n$ or $(n-1)^2 - 1$ oe	B1 B3	[3] an A If	f B0, M2 for recognising form $n^2 + bn + c$ , $a \neq 0$ , b and c not both zero Allow other variables f B0 M0, M1 for reaching second ifferences of 2