PapaCambridge

	1(a)	a=13 b=21 or 8+ their (i)							
	b	z=x+y oe							
2	(a) (b)	$p = 11$ $q = 30$ $r = 60$ $s = 6$ $x = 2n + 1 \text{ oe}$ $y = n(n + 1) \text{ oe}$ $z = 2n(n + 1) \text{ oe} \sqrt{2} \times y$	all four	B2 B1 B1 B1	2 3	sc1 for 2 or 3 correct In (b) , accept any unsimplified form but -1, once, if not given explicitly			
	(c)	102		B1	1 [6]				
	3a)	24 35	1			0			
	3b)	<i>n</i> (<i>n</i> +2) oe	2	B1 fo	B1 for quadratic expression in <i>n</i>				
3	c)(i)	35	3	B2 for 35 × 37 or 35.8 to 35.9 OR M1 for <i>their</i> $n(n + 2) = 1358$ M1 for solution of <i>their</i> quadratic $\frac{-2 \pm \sqrt{2^2 - 4 \times 1 \times (-1358)}}{2 \times 1}$					
30	c)(ii)	7	2	M1F	T for 13	$358-their (c)(i) \times (their (c)(i) + 2)$			
2	l(a)	14 17		1					
2	ł(b)	3n+2 oe final answer				k oe seen 2 oe seen $j \neq 0$			
2	ł(c)	12		<i>their</i> or fo	r(3n+2) or 62 s	stitution of 20 into 2) een or for 38 seen r(3k+2) = 100 - their 62			
5	5(a)(i)	20 24	1	Both c	orrect				
5	(a)(ii)	4n + 4 oe final answer	2	B1 for	4n+k	oe seen			
5	(a)(iii)	36	2	M1 for	r <i>their</i> ((4p+4) = 150 soi			
5	5(b)(i)	44	2	M1 for	$r \frac{26-2}{4}$	$\frac{2}{2}$ or difference = [–]6			
5	(b)(ii)	50 - 6n oe final answer	2	B1 for	-6n+k	k oe seen			

6	$\frac{5n+7}{(n+3)^2}$ oe final answer				4		B2 for <i>n</i> th term for numerator sequence 5n + 7 oe final answer or B1 for $5n + k$ oe seen AND B2 for <i>n</i> th term for denominator sequence $(n + 3)^2$ oe final answer or B1 for quadratic expression in <i>n</i> seen for denominator sequence Maximum 3 marks if final answer incorrect
7(a)	48				1		
7(b)	$n^2 + 5n - 2$ oe final answer				3	0 0	32 for answer $n^2 + 5n + k$ oe r for $5n - 2$ oe seen r B1 for answer $n^2 + an + b$ r for $5n + k$ oe seen
8(a)	9 12 15 12 17 22		2	B 1	for or	ne	row correct
8(b)	5 $n-3$ oe final answer		2	B 1	for 5n	ı +	- <i>k</i> oe seen
8(c)	57		2	M1 or	for <i>th</i> B1 fo	eir r r	r (5n-3) = 92 n = 19 soi or for answer 19
9(a)	$\frac{23}{24} \frac{27}{28}$			1	5	く	
9(b)	300			1	0		
9(c)	$\frac{4n-1}{4n}$ oe	2		2	B1 fo	or	$\frac{\dots}{4n}$, or for $4n - 1$ oe
10(a)	71	5		1			
10(b)	[p =] 2 [q =] 1			1	Bot	h c	correct
10(c)	A = 2 $B = 4$ $C = 1$			2	or fe or fe (<i>n</i> +	or or • <i>tl</i>	r two correct $(n + 1)^2 = n^2 + 2n + 1$ heir $q)^2 = n^2 + 2n(their q) + (their q)^2$ A + B + C = 7 for $4A + 2B + C = 17$ 9A + 3B + C = 31
11(a)	49, 19, 30			1			
11(b)(i)	3n + 4 oe and isw			1			
11(b)(ii)	$(n+2)^2$ oe			1			
11(c)	$n^2 + n$; or $n(n + 1)$			2	both j answe empty	pai er y:	attempt at <i>their</i> (bii) – <i>their</i> (bi), provided rts are different expressions in n , and the space also contains an expression in n , or is valid method.

	12(a)	Correct pattern drawn		
	12(b)	15 21 10 15	B1 for 2 or 3	correct
	12(c)	n^2 oe final answer	e.g. $\left(\frac{1}{2}n^2 + \frac{1}{2}\right)$	$\frac{1}{2}n\left(\frac{1}{2}n^2-\frac{1}{2}n\right)$
	12(d)	465	l 	
	12(e)	$n^2 - \left(\frac{1}{2}n^2 + \frac{1}{2}n\right)$	_	
		or $\left(\frac{1}{2}(n-1)^2 + \frac{1}{2}(n-1)\right)$		
		$n^{2} - \left(\frac{1}{2}n^{2} + \frac{1}{2}n\right)$ or $\left(\frac{1}{2}(n-1)^{2} + \frac{1}{2}(n-1)\right)$ or $\left(\frac{1}{2}n^{2} + \frac{1}{2}n\right) - n$		
		leading to $\left(\frac{1}{2}n^2 - \frac{1}{2}n\right)$ without error AG		
	12(f)		M1 for $\frac{1}{2}m^2$ +	
			or B2 for [<i>m</i> =	$m = 0$ or $m^2 = 9$ m or $m - 9 = 0$ or $m + 1 = 10$ = 9] 5 m = 45 and crosses = 45 these for 5 m and the number of crosses seen for
			at least $m = 7$ After 0, SC1	and 8
13	(a) (i)		1	
	(ii)	421	1	
	(b) (i)	8	1	
	(ii)	14	1	
14	(a) (i)	-6	1	
	(ii)	15	2*	C1 for $15^2 - 5 \times 15$ or for 15, -10 OR
			0.*	M1 for $(p + 10)(p - 15) = 0$
	(b)	4	2*	B1 for $3 \times 5^2 - 5k = 55$ oe
15	(a)	43 47 cao	1	
	(b)			
	```	-)10	1	
	( <b>d</b> )			
	(e)			

16	05 (		2	D1 for (1) good
16	95 - 6n oe isw		2	<b>B1</b> for – 6 <i>n</i> seen
(ii)	16 cao		1	
(b) (i)	2 <i>n</i> – 3		2	<b>M1</b> for $(n + 1)^2 - 4(n + 1)$ seen
(ii)	39 cao		1	
<b>17 (a)</b>			1	
(b) ²	41 40 81 (all three)		1	
(c) (	$(2n+1)^2$ oe		1	
18 (a)	23 – 6 <i>n</i> cao	2	2	<b>B1</b> for <i>6n</i> soi
(b) (i)	4, 10, 18, 28	2	2	<b>B1</b> for 3 correct terms seen
(ii)	3 and 24	2		<b>M1</b> for $\frac{n^2 + 3n}{5n - 12} = 6$ or better <b>M1</b> for $n^2 - 27n + 72 = 0$ <b>B1</b> for either 3 or 24
19 (a)	$15^2 - 1^2 = 8 \times (1 + 2 + 3 + 4 + 5 + 6 + 7)$	7)		
(b)	$(2n+1)^2 - 1^2$ oe		25	ſ
(c)	$(2n+1)^2 = 4n^2 + 4n + 1$ or $(2n+1)^2 - 1^2 = 4n^2 + 4n$ , or $(2n)(2n - 1)^2 - 1^2 = 4n^2 + 4n$	+ 2)	E	31
	Division of both sides by 8 and result o correctly	btained	N	11
20 (a)	25, 21, 45	2	B1 f	or 2 correct
<b>(b)</b>	$n^2$	1		
(c)	32	2	B1 f	or $(T =)$ 1024 seen
( <b>d</b> )	$\frac{3}{2}n(n+1)$ oe	1		
(e)	360	1ft		
(f)	$\frac{1}{2}(n+1)(n+2)$ oe	2	or C	1 for $\frac{1}{2}(n-1)(n-2)$ oe
21(a)	128		1	
21(b)(i) 2	$2^{n+1} - 3$ final answer		1	
21(b)(ii) 2	$2^{n+1} + 5n + 1$ oe final answer		2	<b>B1</b> for $2^{n+1} + 5n + k$ oe
				or <b>M1</b> for 6 11 16 21 [26]

		l					
22	<b>(a)</b>		7, 21			1	
	(b)		2 <i>n</i> – 1 oe			1	
	(c)		FT 3 × <i>their</i> (b) provided this is a function of $n$ ; or $6n - 3$ oe			1 √	
	(d) (	i)	48			1	
	(i	i)	$3n^2$			2 *	M1 for a sensible method, e.g. writing terms as $3 \times 1$ , $3 \times 4$ , $3 \times 9$ , or B1 for $An^2 + Bn + C$ , $A \neq 0$ from a valid
							or <b>B1</b> for $An^2 + Bn + C$ , $A \neq 0$ from a valid method.
23	(a)	201	1			2	<b>B1</b> for $(n = )$ 223 seen
	(b)	36				1	
		(i)	9x - 9y, or $9y - 9x$ , or any equiv.			1	
			"123 is not a multiple of 9" oe			1	
24	(a)	. /	132	1	Cc	ndone -8	87
	(b)		87 f.t. 219 <b>or</b> {{their132  +  their87 }}	1 1			
				5	2	0	