Cambridge IGCSE™

| CANDIDATE NAME	

CENTRE NUMBER		CANDIDATE NUMBER	

ADDITIONAL MATHEMATICS **0606/13**

Paper 1 **May/June 2022**

2 hours

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 80.
- The number of marks for each question or part question is shown in brackets [ ].

This document has **16** pages. Any blank pages are indicated.

Mathematical Formulae

1. ALGEBRA

Quadratic Equation

For the equation $ax^2 + bx + c = 0$,

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Binomial Theorem

$$(a+b)^n = a^n + \binom{n}{1}a^{n-1}b + \binom{n}{2}a^{n-2}b^2 + \ldots + \binom{n}{r}a^{n-r}b^r + \ldots + b^n$$

where n is a positive integer and $\binom{n}{r} = \frac{n!}{(n-r)!\,r!}$

Arithmetic series $u_n = a + (n-1)d$

$$S_n = \frac{1}{2}n(a+l) = \frac{1}{2}n\{2a + (n-1)d\}$$

Geometric series $u_n = ar^{n-1}$

$$S_n = \frac{a(1-r^n)}{1-r} \quad (r \neq 1)$$

$$S_\infty = \frac{a}{1-r} \quad (|r| < 1)$$

2. TRIGONOMETRY

Identities

$$\sin^2 A + \cos^2 A = 1$$
$$\sec^2 A = 1 + \tan^2 A$$
$$\operatorname{cosec}^2 A = 1 + \cot^2 A$$

*Formulae for $\triangle ABC$*

$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc\cos A$$

$$\Delta = \frac{1}{2}bc\sin A$$

1 **(a)** Find the rational numbers a, b and c, such that the first three terms, in descending powers of x, in the expansion of $\left(3x^2 - \dfrac{1}{9x}\right)^5$ can be written in the form $ax^{10} + bx^7 + cx^4$.　　　　　　[3]

(b) Hence find the coefficient of x^4 in the expansion of $\left(3x^2 - \dfrac{1}{9x}\right)^5 \left(1 + \dfrac{1}{x^3}\right)^2$.　　　　　　[3]

2 In this question, all lengths are in centimetres and all angles are in radians.

The diagram shows a circle, centre O, radius 8. The points A, B and C lie on the circumference of the circle. The chord AB has length 10.

(a) Show that angle BOA is 1.35 correct to 2 decimal places. [2]

(b) Given that the minor arc BC has a length of 18, find angle BOC. [2]

(c) Find the area of the minor sector AOC. [3]

3 **(a)** Find the exact solution of the equation $2e^{6x} - 3e^{3x} - 5 = 0$. [3]

(b) Solve the following simultaneous equations.

$$e^{4x-7} \div e^{5x+7y} = \frac{1}{e^2}$$
$$xy + 18 = 0$$ [5]

 [Turn over

6

4 Variables x and y are such that when e^{4y} is plotted against x, a straight line of gradient $\frac{2}{5}$, passing through (10, 2), is obtained.

(a) Find y in terms of x. [3]

(b) Find the value of y when $x = 45$, giving your answer in the form $\ln p$. [2]

(c) Find the values of x for which y can be defined. [1]

5 The velocity, $v\,\text{ms}^{-1}$, of a particle moving in a straight line, t seconds after passing through a fixed point O, is given by $v = 6\sin 3t$.

(a) Find the time at which the acceleration of the particle is first equal to $-9\,\text{ms}^{-2}$. [4]

(b) Find the displacement of the particle from O when $t = 5.6$. [4]

6 **(a)** It is given that

$$f : x \rightarrow 2x^2 \text{ for } x \geqslant 0,$$

$$g : x \rightarrow 2x+1 \text{ for } x \geqslant 0.$$

Each of the expressions in the table can be written as one of the following.

$$f' \quad f'' \quad g' \quad g'' \quad fg \quad gf \quad f^2 \quad g^2 \quad f^{-1} \quad g^{-1}$$

Complete the table. The first row has been completed for you. [5]

Expression	Function notation
2	g'
0	
$4x$	
$8x^2 + 8x + 2$	
$4x + 3$	
$\dfrac{x-1}{2}$	

(b) It is given that $h(x) = (x-1)^2 + 3$ for $x \geqslant a$. The value of a is as small as possible such that h^{-1} exists.

    (i) Write down the value of a. [1]

    (ii) Write down the range of h. [1]

    (iii) Find $h^{-1}(x)$ and state its domain. [3]

7 A curve has equation $y = \dfrac{(2x+1)^{\frac{3}{2}}}{x+5}$ for $x \geqslant 0$.

(a) Show that $\dfrac{\mathrm{d}y}{\mathrm{d}x} = \dfrac{(2x+1)^{\frac{1}{2}}}{(x+5)^2}(Ax+B)$, where A and B are integers to be found. [4]

(b) Show that there are no stationary points on this curve. [1]

(c) Find the approximate change in y when x increases from 1 to $1+p$, where p is small. [2]

(d) Given that when $x = 1$ the rate of change in x is 2.5 units per second, find the corresponding rate of change in y. [2]

8 **(a)** A 6-digit number is formed from the digits 0, 1, 2, 5, 6, 7, 8, 9. A number cannot start with 0 and each digit can be used at most once in any 6-digit number.

    (i) Find how many 6-digit numbers can be formed if there are no further restrictions.   [1]

    (ii) Find how many of these 6-digit numbers are divisible by 5.   [3]

    (iii) Find how many of these 6-digit numbers are greater than 850 000.   [3]

(b) A team of 8 people is to be chosen from 12 people. Three of the people are brothers who must not be separated. Find the number of different teams that can be chosen. [3]

9 **(a)** Solve the equation $3\csc^2\left(2\phi - \dfrac{\pi}{3}\right) = 4$, for $0 < \phi < \pi$. Give your solutions in terms of π.

[4]

(b) Given that $2x - 1 = \csc^2\theta$ and $y + 1 = \tan^2\theta$, find y in terms of x. [4]

10 (a) Show that $\dfrac{6}{2+3x} + \dfrac{4}{(x+1)^2} - \dfrac{2}{x+1}$ can be written as $\dfrac{14x+10}{(2+3x)(x+1)^2}$. [2]

(b) Hence find the exact value of $\displaystyle\int_0^2 \dfrac{14x+10}{(2+3x)(x+1)^2}\,dx.$ Give your answer in the form $p + \ln q$,

where p and q are rational numbers. [6]

BLANK PAGE