

## **Cambridge International Examinations**

Cambridge International General Certificate of Secondary Education

MATHEMATICS (US)

Paper 2 (Extended)

May/June 2016

MARK SCHEME
Maximum Mark: 70

## **Published**

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2016 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.



Page 2	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2016	0444	21

## **Abbreviations**

cao correct answer only

dep dependent

FT follow through after error isw ignore subsequent working

oe or equivalent SC Special Case

nfww not from wrong working

soi seen or implied

Question	Answer	Mark	Part marks
1	8(h) 52 (min)	1	
2	12	1	
3	[0].00127	1	
4	28	2	<b>B1</b> for 24 or −3
5	540	2	<b>M1</b> for 2000 × 0.27
6	144	2	M1 for finding a correct product of prime factors or correctly listing a minimum of 3 multiples of 36 and 48 or for answer $2^4 \times 3^2$ oe or $144k$
7	11	2	M1 for $-2 \times -7 -3$ soi
8	$\frac{py}{q}$ final answer	2	M1 for 1 correct step
9	[a = ] 70 [b = ] 40	2	B1 for each
10	$ \begin{bmatrix} x = ] - 2 \\ y = ] 7 $	1 1	If 0 scored, <b>SC1</b> for two values satisfying one of the original equations
11 (a)	112	1	
(b)	56	1	
12	$2p^4$ final answer	2	<b>B1</b> for $kp^4$ or $2p^k$ as answer
13	$n > \frac{15}{4}$	2	<b>M1</b> for $7 + 8 < 5n - n$ oe
14	$2\cos\frac{1}{2}x$	3	<b>B1</b> for cos <b>B1</b> for amplitude = 2 or 2sin or 2cos <b>B1</b> for $\frac{1}{2}x$ oe

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2016	0444	21

Question	Answer	Mark	Part marks
15 (a)	-3	1	
(b)	9 – 2 <i>n</i> oe	2	<b>B1</b> for $-2n + k$ or $dn + 9$ where $d \neq 0$
16	$\frac{18}{35}$ cao	3	M2 for $\frac{6}{7} \times \frac{3}{5}$ or $\frac{18}{21} \div \frac{35}{21}$ oe  B1 for $\frac{5}{3}$ oe  or M1 for $\frac{6}{7} \times their \frac{3}{5}$
17	145	3	M2 for $(6-2) \times 180 - 5 \times 115$ or M1 for $(6-2) \times 180$ Alt method M2 for $180 - (360 - 5 \times (180 - 115))$ or M1 for $360 - 5 \times (180 - 115)$
18	2 nfww	3	M2 for $(36 + 4) \div (72 \times \frac{1000}{60 \times 60})$ oe or M1 for $72 \times \frac{1000}{60 \times 60}$ or for a distance $\div$ a speed SC2 for answer 1.8
19	2	3	M1 for $y = k\sqrt{x}$ A1 for $k = 4$ or M2 for $\frac{\sqrt{9}}{12} = \frac{\sqrt{\frac{1}{4}}}{y}$ oe
20	5 6	3	M2 for $1 - \frac{2}{3} \times \frac{1}{4}$ or $\frac{1}{3} + \frac{2}{3} \times \frac{3}{4}$ or $\frac{1}{3} \times \frac{3}{4} + \frac{1}{3} \times \frac{1}{4} + \frac{2}{3} \times \frac{3}{4}$ or M1 for $\frac{2}{3} \times \frac{1}{4}$ or $\frac{1}{3} \times \frac{1}{4} + \frac{2}{3} \times \frac{3}{4}$
21 (a)	$5\sqrt{5}$ final answer	1	
(b)	$-24-5\sqrt{5}$ final answer	2	<b>B1</b> for three terms correct from $6-9\sqrt{5}+4\sqrt{5}-6\times\sqrt{5}\times\sqrt{5}$
22	27	3	M2 for $\frac{6\pi}{\pi \times 2 \times 9} \times \pi \times 9^2$ oe or M1 for $\frac{6\pi}{\pi \times 2 \times 9}$ oe

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge IGCSE – May/June 2016	0444	21

Question	Answer	Mark	Part marks
23	30 nfww	4	M2 for height = $\sqrt{5^2 - 4^2}$ or M1 for $4^2 + h^2 = 5^2$ oe and M1 for $\frac{1}{2}(8+12) \times their$ perp height oe
24 (a)	(a+2)(2+p)	2	<b>B1</b> for $2(a+2) + p(a+2)$ or $a(2+p) + 2(2+p)$
(b)	2(9+2t)(9-2t) oe	2	<b>B1</b> for $2(81-4t^2)$ oe or $(18+4t)(9-2t)$ oe If 0 scored <b>SC1</b> for $(9+2t)(9-2t)$ final answer
25	$y = -\frac{3}{7}x + 11 \text{ oe}$	6	<b>B2</b> for gradient = $-\frac{3}{7}$ or M1 for [gradient = ] $\frac{15-1}{10-4}$ oe or for the negative reciprocal of <i>their</i> gradient and B2 for [midpoint of $AB = (7, 8)$ ] or B1 for $(7, k)$ or $(k, 8)$ ] and M1 for substitution of <i>their</i> midpoint or $(4, 1)$ or $(10, 15)$ into a linear equation
26 (a)	$6\sqrt{3}$	3	M2 for $\frac{1}{2} \times 8 \times 3 \times \frac{\sqrt{3}}{2}$ oe or M1 for $\frac{1}{2} \times 8 \times 3 \times \sin 60$ oe or B1 for $[\sin 60 = ]$ $\frac{\sqrt{3}}{2}$
(b)	7	3	M2 for $3^2 + 8^2 - 2 \times 3 \times 8 \times \frac{1}{2}$ oe or M1 for $3^2 + 8^2 - 2 \times 3 \times 8 \times \cos 60$ oe or B1 for $[\cos 60 = ] \frac{1}{2}$