

Cambridge IGCSE™

CAMBRIDGE INTERNATIONAL MATHEMATICS Paper 6 (Extended) MARK SCHEME Maximum Mark: 60 Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge International will not enter into discussions about these mark schemes.

Cambridge International is publishing the mark schemes for the October/November 2023 series for most Cambridge IGCSE, Cambridge International A and AS Level components, and some Cambridge O Level components.

Generic Marking Principles

These general marking principles must be applied by all examiners when marking candidate answers. They should be applied alongside the specific content of the mark scheme or generic level descriptors for a question. Each question paper and mark scheme will also comply with these marking principles.

GENERIC MARKING PRINCIPLE 1:

Marks must be awarded in line with:

- the specific content of the mark scheme or the generic level descriptors for the question
- the specific skills defined in the mark scheme or in the generic level descriptors for the question
- the standard of response required by a candidate as exemplified by the standardisation scripts.

GENERIC MARKING PRINCIPLE 2:

Marks awarded are always whole marks (not half marks, or other fractions).

GENERIC MARKING PRINCIPLE 3:

Marks must be awarded **positively**:

- marks are awarded for correct/valid answers, as defined in the mark scheme. However, credit is given for valid answers which go beyond the scope of the syllabus and mark scheme, referring to your Team Leader as appropriate
- marks are awarded when candidates clearly demonstrate what they know and can do
- marks are not deducted for errors
- marks are not deducted for omissions
- answers should only be judged on the quality of spelling, punctuation and grammar when these features are specifically assessed by the question as indicated by the mark scheme. The meaning, however, should be unambiguous.

GENERIC MARKING PRINCIPLE 4:

Rules must be applied consistently, e.g. in situations where candidates have not followed instructions or in the application of generic level descriptors.

GENERIC MARKING PRINCIPLE 5:

Marks should be awarded using the full range of marks defined in the mark scheme for the question (however; the use of the full mark range may be limited according to the quality of the candidate responses seen).

GENERIC MARKING PRINCIPLE 6:

Marks awarded are based solely on the requirements as defined in the mark scheme. Marks should not be awarded with grade thresholds or grade descriptors in mind.

© UCLES 2023 Page 2 of 7

Mathematics-Specific Marking Principles

- 1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.
- 2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.
- Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.
- 4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).
- Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.
- Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

MARK SCHEME NOTES

The following notes are intended to aid interpretation of mark schemes in general, but individual mark schemes may include marks awarded for specific reasons outside the scope of these notes.

Types of mark

- M Method marks, awarded for a valid method applied to the problem.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. For accuracy marks to be given, the associated Method mark must be earned or implied.
- B Mark for a correct result or statement independent of Method marks.

When a part of a question has two or more 'method' steps, the M marks are in principle independent unless the scheme specifically says otherwise; and similarly where there are several B marks allocated. The notation 'dep' is used to indicate that a particular M or B mark is dependent on an earlier mark in the scheme.

Abbreviations

awrt answers which round to cao correct answer only

dep dependent

FT follow through after error isw ignore subsequent working nfww not from wrong working

oe or equivalent

rot rounded or truncated

SC Special Case soi seen or implied

© UCLES 2023 Page 3 of 7

Question	Answer	Marks	Partial Marks
1(a)(i)	1024	1	
1(a)(ii)	4^{8} or 256×4^{4} oe or 1024×4^{3} oe or 1024 , 4096 , 16384	C1	
	65536	1	
1(a)(iii)	4^n oe	1	
1(b)(i)	3×2^0	1	
1(b)(ii)	6, 12, 24	1	
1(c)	36, 18, 9, 4.5	2	B1 for 3 terms correct
1(d)	$5 \times 3^{n-1}$	2	B1 for $[5 \times] 3^k$ for some k where k is an expression in terms of n
2	$12 \times \frac{12}{16}$ oe or $16 \times \left(\frac{12}{16}\right)^{3-1}$ oe leading to 9	2	B1 for $\frac{12}{16}$ oe or $\frac{16}{12}$ oe seen
3(a)(i)	62.5 ÷ 12.5	C1	
	12.5 ÷ their (62.5 ÷ 12.5)	C1	
	2.5 oe	1	
3(a)(ii)	their $2.5 \times (their (62.5 \div 12.5))^{n-1}$	1	FT
3(b)	$24k^3 = 81$ oe with one further correct step leading to 1.5 oe or $\sqrt[3]{\frac{81}{24}} = 1.5$	C2	C1 for $24k^3 = 81$ or $\frac{81}{24}$ oe or 1.5 used as multiplier
	$81 \times their \left(\frac{81}{24}\right)^{\frac{1}{3}}$ oe	C1	
	121.5 oe	1	
4(a)(i)	Correct comment referring to Multiplier/divisor/scale factor/common ratio and Equal to the same	1	
4(a)(ii)	Eliminating fractions e.g. $(2x+3)^2 = x(11x-6)$ oe	1	
	$(2x+3)^2 = 4x^2 + 6x + 6x + 9$	1	

© UCLES 2023 Page 4 of 7

Question	Answer	Marks	Partial Marks
	Rearrangement with one further correct step leading to $7x^2 - 18x - 9 = 0$.	1	
4(b)	(7x+3)(x-3) = 0	1	
	$x = 3 \text{ and } x = -\frac{3}{7}$	1	If 0 scored SC1 for $(7x-3)(x+3)$ and $x = \frac{3}{7}$ and $x = -3$
4(c)	Two correct sequences of three terms or nth term 3, 9, 27 or $3 \times 3^{n-1}$ and $-\frac{3}{7}$, $\frac{15}{7}$, $-\frac{75}{7}$	C2	FT their 3 and their $-\frac{3}{7}$ but not if decimal C1FT for one correct sequence of three terms
	$\frac{81}{\frac{375}{7}} \text{ or } 53\frac{4}{7}$	2	B1 for each
5(a)	$20 \times 20 \times 25$	C1	
	10 000 cm ³	1	
5(b)(i)	Full correct method shown. e.g. 20×20 [× 1] ÷ $50 = 8$ or $10\ 000 \div 25 \div 50 = 8$	2	B1 for $400 \times 1 = 50 = 8$ or B1FT their $10000 \div 25$ or their $10000 \div 50$ or their $10000 \div 1250$
5(b)(ii)	200	1	
5(c)(i)	Every 8 seconds the height goes up 1 cm oe	1	
5(c)(ii)	Substituting $t = 60$ into $8h = t$ (implied by $60 \div 8$)	C1	
	7.5 cm oe	1	
5(c)(iii)	Correct sketch	1	Ruled straight line from the origin with positive gradient reaching approx. $h = 25$
	Labelling t axis and line goes through approx. (25, 200) or (25, 200) marked on the line above $h = 25$	C1	
6(a)	$\pi \times 16 \times 16 \times 24 \div 3$ oe	C1	

© UCLES 2023 Page 5 of 7

Question	Answer	Marks	Partial Marks
	6430 cm ³	1	
6(b)(i)	$[x =] \frac{16}{24} \times 3 \text{oe}$	2	B1 for $\frac{x}{3} = \frac{16}{24}$ or $\frac{x}{16} = \frac{3}{24}$
	or $[x =] \frac{3}{24} \times 16$ oe		or scale factor = $\frac{3}{24}$ or $\frac{24}{3}$ oe
	or $[x =] 16 \div \frac{24}{3}$ oe		
	leading to $[x =] 2$		
6(b)(ii)	$\frac{1}{3} \times \pi \times 2^2 \times 3$	C1	
	12.6 cm ³	1	
6(b)(iii)	their 12.6 ÷ 20	C1	FT
	0.628 seconds	1	
6(c)(i)	$t = \frac{\frac{1}{3}\pi\left(\frac{2}{3}\right)^2 h^3}{20}$ leading to $\frac{\pi h^3}{135}$	3	B2 for $\frac{1}{3}\pi \left(\frac{2}{3}h\right)^2 h$ oe
	20 135		$\mathbf{B1} \text{ for } \frac{r}{h} = \frac{16}{24} \text{ oe}$
6(c)(ii)	$\frac{\pi 24^3}{135} \text{ or their } \mathbf{6(a)} \div 20$	C1	
	322	1	
6(c)(iii)	Correct Sketch 322 0 0 24	1	Constantly increasing curve from the origin reaching approx. $h = 24$
	Labelling t axis and curve goes through approx. (24, 322) or (24, 322) marked on the line above $h = 24$	C1	

© UCLES 2023 Page 6 of 7

Question	Answer	Marks	Partial Marks
6(d)	$\frac{\frac{1}{3}\pi\left(\frac{2}{3}\right)^2 h^3}{\text{Rate}} = \frac{\pi h^3}{189} \text{ oe}$ or $\frac{\left[\right]}{20} = \frac{189}{135} \text{ oe}$ or $\frac{\pi 24^3}{189}$ or cm ³ per second oe	C1	
	28	1	
7(a)	$\frac{\pi h^3}{135} = 8h$ or a sketch of both graphs.	C1	
	18.5 cm	1	
7(b)	148 s	1	If 0 or C1 only scored in 7(a) and 7(b) then also SC1 for reversed correct answers

© UCLES 2023 Page 7 of 7