WWW. Pals

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

MARK SCHEME for the October/November 2009 question paper for the guidance of teachers

0652 PHYSICAL SCIENCE

0652/05

Paper 5 (Practical Test), maximum raw mark 30

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	10
	IGCSE – October/November 2009	0652	100

1 (a) recording the refractive index;

(d) 5 suitable drawings;
Give ONE mark if only 4;

Table

angles of \boldsymbol{i} are about 60, 55, 40, 30 and 20 and \boldsymbol{r} values ;

ONE mark if only 4 values;

angles of **r** decrease appropriately; [1]

(e) Graph

scale; plotting;

suitable curve ; [3]

(f) correctly read from graph; [1]

(g) both sines correct, estimating if necessary; if clearly not from the table provided do not allow ratio correctly calculated; [2]

(h) suitable comment; looking for some idea of accuracy of experiment. [1]

(i) if the refractive index is greater, then angles of refraction would be smaller; or vice versa the denser the medium the greater the ray is bent; [2]

[Total: 15]

[2]

Page 3	Mark Scheme: Teachers' version	Syllabus	. O. T.
	IGCSE – October/November 2009	0652	100

2 (a) accept anything between 0.03 and 0.08;

(c) results for all three; value for B is half A within 2 drops; value for C is half B within 2 drops; value for C within 2 drops of Supervisors value;	
if A is larger than 20, 2 drops becomes 3 drops	[4]
(d) most concentrated is A; needs the largest no. of drops;	[2]
(e) brown ppt.;	[1]
(f) (i) white <u>ppt</u> .;	[1]
(ii) dirty white, grey or cloudy (ppt. not necessary here as reaction is	slow); [1]
(iii) green <u>ppt</u> .;	[1]
(g) one mark for iron(II) and one for sulfate;	[2]
(h) count drops into a measuring cylinder to suitable volume (at least 5 condivide by number of drops;	m ³) ; [2]

[Total: 15]