Cambridge International Examinations
Cambridge International General Certificate of Secondary Education

PHYSICS **0625/32**

Paper 3 Core Theory **October/November 2016**

MARK SCHEME

Maximum Mark: 80

Published

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2016 series for most Cambridge IGCSE®, Cambridge International A and AS Level components and some Cambridge O Level components.

This syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

Question	Answer	Marks
1(a)	speed **OR** velocity on y-axis **AND** time x-axis	**B1**
1(b)	A to B	**B1**
1(c)	area under graph	**C1**
	$0.5 \times 5 \times 5$ (+ (3×5))	**C1**
	27.5 (m)	**A1**
1(d)	correctly placed continuous single thin straight line from A to E drawn using a rule	**B1**
	Total:	**6**

Question	Answer	Marks
2(a)	1 rule(r) 2 balance	**B2**
2(b)	250 (cm^3)	**B1**
2(c)	D = M/V in any form	**C1**
	20/250	**C1**
	0.8 (g/cm^3)	**A1**
2(d)	freon, glycerol, sea water	**B2**
	Total:	**8**

Question	Answer	Marks
3(a)	both boxes ticked	**B1**
3(b)	moment = force × distance in any form	**C1**
	300 × 1.4	**C1**
	420 (Nm)	**A1**
3(c)	clockwise moments = anticlockwise moments	**C1**
	W x 0.6 = candidates (b) **OR** *W* = candidates (b) / 0.6	**C1**
	700 (N)	**A1**
3(d)	child 's OR left side goes down **OR** adult side goes up **OR** right side goes up **OR** child's moment is larger **OR** child's turning force larger	**B1**
	Total:	**8**

Question	Answer	Marks
4(a)	$W = m \times g$ in any form	C1
	400 (N)	A1
4(b)	pressure = force ÷ area in any form	C1
	400 **OR** candidates (a) ÷ 0.02	C1
	20 000 (N/m^2)	A1
4(c)	greater pressure **OR** wtte	B1
	(same force/weight acts on a) smaller area	B1
	Total:	7

Question	Answer	Marks
5(a)	radiation	B1
5(b)	black can has bigger rise or higher temperature	B1
	silver reflects (radiant) heat (better) **OR** poor absorber of (radiant) heat	B1
	black is (a better) absorber of thermal energy	B1
5(c)	evaporation/evaporated	B1
	more energetic or higher energy molecules	B1
	overcome force of attraction	B1
	Total:	**7**

Question	Answer	Marks
6(a)	(angle of) reflection	B1
6(b)(i)	image 'I' correctly positioned	B1
6(b)(ii)	angle of reflection incorrect **OR** object and image are not same distance from mirror owtte	C1
	angle of incidence ≠ angle of reflection owtte	A1
	Total:	**4**

Question	Answer	Marks
7(a)	speed = distance ÷ time in any form	C1
	indication of halving e.g. 450/2 **OR** 1500 × 0.15	C1
	225 (m)	A1
7(b)	more than 20 000 Hz	B1
7(c)	any wave from electromagnetic spectrum	B1
	Total:	5

Question	Answer	Mark
8(a)	30 ÷ 4	C1
	7.5 (cm)	A1
8(b)	number of waves (passing a point) in 1 second	B1
8(c)	f = 4/0.05	C1
	80	A1
	Hz	B1
	Total:	6

Question	Answer	Marks
9(a)(i)	changes higher voltage to lower voltage owtte	**B1**
9(a)(ii)	copper	**B1**
9(a)(iii)	$V_s/V_p = N_s/N_p$ in any form $(12/240) \times 10\,000 \div 20$ 500	**C1** **C1** **A1**
9(b)	any two from: thinner wires or cables less heating or less energy or power wasted or more efficient lower current in cables fewer power stations needed transmit longer distances (without drop in power)	**B2**
	Total:	**7**

Question	Answer	Marks
10(a)	heater clearly identified	**B1**
10(b)(i)	change current	**B1**
10(b)(ii)	change temperature of heater or output of heater	**B1**
10(c)	V = IR in any form or V ÷ I	**C1**
	250 ÷ 2	**C1**
	125(Ω)	**A1**
10(d)	fuse	**M1**
	(large) current melts fuse wire owtte	**A1**
	Total:	**8**

Question	Answer	Marks
11(a)(i)	(current flow of charge in) one direction owtte	**B1**
11(a)(ii)	iron forms (temporary) magnet	**B1** **B1**
11(b)	Any three from: current in coil creates electromagnet owtte (electromagnet) attracts armature contacts (on 2nd circuit) close 2nd circuit complete	**B3**
11(c)	prevent overheating of cables owtte	**B1**
	Total:	**7**

Question	Answer	Marks
12(a)	unstable atoms	**B1**
	random/spontaneous decay (of atoms)	**B1**
12(b)(i)	20 cpm = approx. 9000 **AND** 10 cpm = approx. 15 000	**B1**
12(b)(ii)	5000 – 6500	**B1**
12(c)	two half-life indicated	**B1**
	2.5 (g)	**B1**
12(d)	any sensible precaution: tongs/screening/lead apron minimise time exposure maximise distance between source and people restrict access to sources etc.	**B1**
	Total:	**7**