

## MARK SCHEME for the May/June 2013 series

# 0443 PHYSICS (US)

0443/33

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	S.	Y
	IGCSE – May/June 2013	0443	10.	

### NOTES ABOUT MARK SCHEME SYMBOLS & OTHER MATTERS

- Cambridge.com are method marks upon which further marks depend. For an M mark to be scored M marks point to which it refers **must** be seen in a candidate's answer. If a candidate fails to see a particular M mark, then none of the dependent marks can be scored.
- B marks are independent marks, which do not depend on other marks. For a B mark to be scored, the point to which it refers must be seen specifically in the candidate's answers.
- A marks In general A marks are awarded for final answers to numerical questions. If a final numerical answer, eligible for A marks, is correct, with the correct unit and an acceptable number of significant figures, all the marks for that question are normally awarded. It is very occasionally possible to arrive at a correct answer by an entirely wrong approach. In these rare circumstances, do not award the A marks, but award C marks on their merits. However, correct numerical answers with no working shown gain all the marks available.
- C marks are compensatory marks in general applicable to numerical questions. These can be scored even if the point to which they refer are not written down by the candidate, provided subsequent working gives evidence that they must have known it. For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct substitution or working which shows he knew the equation, then the C mark is scored. A C mark is not awarded if a candidate makes two points which contradict each other. Points which are wrong but irrelevant are ignored.
- brackets () around words or units in the mark scheme are intended to indicate wording used to clarify the mark scheme, but the marks do not depend on seeing the words or units in brackets, e.g. 10 (J) means that the mark is scored for 10, regardless of the unit given.
- underlining indicates that this must be seen in the answer offered, or something very similar.
- OR / or indicates alternative answers, any one of which is satisfactory for scoring the marks.
- means 'each error or omission'. e.e.o.o.
- means 'or words to that effect'. o.w.t.t.e.
- Spelling Be generous about spelling and use of English. If an answer can be understood to mean what we want, give credit. However, beware of and do not allow ambiguities, accidental or deliberate: e.g. spelling which suggests confusion between reflection / refraction / diffraction / thermistor / transistor / transformer.
- Not/NOT Indicates that an incorrect answer is not to be disregarded, but cancels another otherwise correct alternative offered by the candidate, i.e. right plus wrong penalty applies.
- Ignore Indicates that something which is not correct or irrelevant is to be disregarded and does not cause a right plus wrong penalty.

			2.
Page 3	Mark Scheme	Syllabus	S.
	IGCSE – May/June 2013	0443	10

e.c.f. meaning 'error carried forward' is mainly applicable to numerical questions, particular circumstances be applied in non-numerical questions. This indicates that if a candidate has made an earlier mistake and has carrie incorrect value forward to subsequent stages of working, marks indicated by ecf may awarded, provided the subsequent working is correct, bearing in mind the earlier mistake. This prevents a candidate being penalised more than once for a particular mistake, but **only** applies to marks annotated e.c.f.

#### Significant Figures

Answers are normally acceptable to any number of significant figures  $\dot{u}$  2. Accept answers that round to give the correct answer to 2 s.f. Any exceptions to this general rule will be specified in the mark scheme.

Units Deduct one mark for each incorrect or missing unit from a final answer that would otherwise gain all the marks available for that answer: maximum 1 per question.

#### Arithmetic errors

Deduct one mark if the **only** error in arriving at a final answer is clearly an arithmetic one.

#### Transcription errors

Deduct one mark if the only error in arriving at a final answer is because given or previously calculated data has clearly been misread but used correctly.

Fractions e.g. <sup>1</sup>/<sub>2</sub>, <sup>1</sup>/<sub>4</sub>, 1/10 etc. are only acceptable where specified.

Pa	age 4		abus 443	<u> </u>
(-)	(1)	· · · ·	abus 443 C1 A1	
(a)	(i)	constant/uniform gradient/slope OR straight line		760
	(ii)	$(a = \Delta) v \div t \text{ OR } 36 \div 48$ 0.75 m/s <sup>2</sup> (NOT 0.76)	C1 A1	
(b)	(i)	horizontal line from (48, 36) to (120, 36)	B1	
	(ii)	area <u>under</u> graph (mentioned <b>or</b> implied)	B1	
		864 OR 2592 3500/3460/3456 m	C1 A1	[7
(a)	(i)	$(m = ) \rho V \text{ OR } 1000 \times 1.8 \times 10^6$	C1	
		1.8×10 <sup>9</sup> kg	A1	
	(ii)	(g.p.e. = ) <i>mgh</i> OR 1.8×10 <sup>9</sup> ×10×350 (e.c.f. from <b>(a)(i)</b> ) 6.3 × 10 <sup>12</sup> J (e.c.f. from <b>(a)(i)</b> )	C1 A1	
	(iii)	$(P = )E/t \text{ OR } 6.3 \times 10^{12}/7 \text{ OR } 6.3 \times 10^{12}/(7 \times 60) \text{ OR } 6.3 \times 10^{12}/(7 \times 10^{12}/(7 \times 60)) \text{ OR } 6.3 \times 10^{12}/(7 \times 10^{12}$	3600) C1	
		2.5 × 10 <sup>8</sup> W (e.c.f. from <b>(a)(i)(ii)</b> )	A1	
(b)	(i)	continuously regenerated / not used up / everlasting supply		
( )		IGNORE used again / recycled / can be renewed	B1	
	(ii)	any <b>two</b> of: biomass/geothermal/solar/ tidal/wave/wind energy/wind energy/win		-
		(NOT nuclear/light)	B2	[6
(a)	velo	ocity has direction/is a vector AND speed doesn't/isn't/is a scalar	B1	
. ,				
(b)	(i)	horizontal arrow to right AND touching parachutist (when extend	,	
		arrow/line horizontal AND arrow / line vertical AND making two OR rectangle	sides of triangle B1	
	(ii)	correct diagonal (i.e. top left to bottom right)	B1	
	( )	10.4–10.5 m/s 51–55° to horizontal OR 35–39° to vertical (NOT more than 2 s	iq.fiqs.) B1	
	(iii)	<sup>1</sup> /₂ <i>mv</i> <sup>2</sup> OR 0.5×85×10.5 <sup>2</sup> (e.c.f. from (b)(ii)) 0.5×85×10.5 <sup>2</sup> (e.c.f. from (b)(ii))	C1 C1	

	ige 5	5 Mark Scheme Syllabus	No. Y
		IGCSE – May/June 2013 0443	1230
(a)	850	000N (accept 83 300N)	ambrio
(b)	(i)	( <i>P</i> = ) <i>F</i> /A OR 85000/3.4 OR 85000/3.4 × 2 OR 85000/6.8 (e.c.f. from (a 1.2/1.25/1.3 × 10 <sup>4</sup> Pa (e.c.f. from (a)(i))	)(i)) C1 A1
	(ii)	larger area smaller pressure	M1 A1
(c)	(i)	(measure of) turning effect OR $F \times x$	B1
	(ii)	no resultant/net force no resultant/net turning effect/moment	B1 B1 [8]
(a)	boil boil boil	y two of: iling throughout liquid (evaporation at surface), iling at one temperature (evaporation at any / all temperature / below boil iling not affected by draught/area (evaporation is), iling produces bubbles (evaporation does not).	ng point), B2
(b)		ermal energy) does work against intermolecular forces / breaks bonds blecules separated/moved apart OR becomes PE	B1 B1
(c)	mea two dete	paratus: e.g. kettle AND balance / scales OR steam condensing in water easuring cylinder / scales AND thermometer o masses determined OR volume/mass condensed termine energy input: e.g. <i>VIt</i> <b>or</b> <i>Pt</i> <b>or</b> <i>mc</i> ⊿ <i>T</i> = )Q/m	with B1 B1 B1 B1 [8]
(a)	(i)	any two of: (gas) molecules further apart greater PE move singly / in straight lines OR vice versa for. <u>liquid</u> molecules (allow faster)	B2
	(ii)	gases compressible OR liquids incompressible forces between gas molecules weaker OR vice versa for liquid molecul	B1 es B1
(b)	(i)	$pV$ = constant OR $p_1V_1 = p_2V_2$ OR 2.6×10 <sup>6</sup> ×0.035 OR 91000 2.6×10 <sup>6</sup> ×0.035/1.0×10 <sup>5</sup> OR 91000/1.0×10 <sup>5</sup> 0.91 m <sup>3</sup>	C1 C1 A1

Pa	ge 6	i				Μ	lark S	Schen	ne				Syll	abus		S.	X	
					IG	CSE	– Ma	ay/Jur	1e 20'	13			04	143		No.	2	
(a)	(i)	(onl	ly) or	e fre	quenc	y (ac	cept	wavel	ength	)							an	76.
	(ii)	4.7	×10 <sup>1</sup>	<sup>1</sup> Hz (	OR the	e sam	ne as	befor	e OR	uncha	anged				43NM		B1	10
(b)	(i)	( <i>n</i> = 1.5	= )c/v	OR 3	8.0×10	0 <sup>8</sup> / 2	2.0×1	0 <sup>8</sup>									M1 A1	
	(ii)	(λ = 4.3/	= ) <i>c/f</i> /4.26/	OR 2 4.25	.0×10 5319×	) <sup>8</sup> /4.7 : 10 <sup>-7</sup>	/×10 <sup>1</sup> m	14									C1 A1	[6]
(a)							-	trons /not fre	•		,						B1 B1	
(b)	èar	th/to	uch (	with h	ed nylo nand) d (and	the s	phere		sphere	e							B1 B1 B1	
(c)				•	•			Il lines Ione w		surfa	ce						M1 A1	[7]
(a)	(i)	san	ne nu	mber	of / 9	2 pro	otons	(in nu	icleus	) (IGN	IORE e	electro	ons)				B1	
	(ii)	diffe	erent	numl	per of	neuti	rons										B1	
(b)	nuc sma	leus all nu	sma umbe	l / ato r defl	om mo ected	ostly e (thro	empty ugh la	rough y spac arge a ī nucle	ce angles	s)	/charge	ed					M1 A1 M1 A1	[6]
(a)	in c	order	dow	nward	ls: 1 1	10	c.a.o.										B1	
(b)	(i)	1 A	ND 0	(e.c.	f. from	n <b>(b)(</b>	i))										B1	
	(ii)	NO	T (ga	te) ( <b>a</b>	llow	NOR	(gate	e))									B1	
(c)	R = T =		ND S	= 0 (	e.c.f.	from	(b)(i)	))									B1 B1	[5]

