CAMBRIDGE INTERNATIONAL EXAMINATIONS Cambridge International General Certificate of Secondary Education

www.papacambridge.com MARK SCHEME for the October/November 2014 series

0443 PHYSICS (US)

0443/33

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2014 series for most Cambridge IGCSE[®], Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	S
	Cambridge IGCSE – October/November 2014	

NOTES ABOUT MARK SCHEME SYMBOLS AND OTHER MATTERS

- www.papaCambridge.com B marks are independent marks, which do not depend on other marks. For a B mark B marks be scored, the point to which it refers must be seen specifically in the candidate's answer.
- M marks M marks are method marks upon which accuracy marks (A marks) later depend. For an M mark to be scored, the point to which it refers **must** be seen in a candidate's answer. If a candidate fails to score a particular M mark, then none of the dependent A marks can be scored.
- C marks C marks are compensatory marks in general applicable to numerical questions. These can be scored even if the point to which they refer are not written down by the candidate. provided subsequent working gives evidence that they must have known it. For example, if an equation carries a C mark and the candidate does not write down the actual equation but does correct substitution or working which shows he knew the equation, then the C mark is scored. A C mark is not awarded if a candidate makes two points which contradict each other. Points which are wrong but irrelevant are ignored.
- A marks A marks are accuracy or answer marks which either depend on an M mark, or which are one of the ways which allow a C mark to be scored. A marks are commonly awarded for final answers to numerical questions. If a final numerical answer, eligible for A marks, is correct, with the correct unit and an acceptable number of significant figures, all the marks for that question are normally awarded. It is very occasionally possible to arrive at a correct answer by an entirely wrong approach. In these rare circumstances, do not award the A marks, but award C marks on their merits. An A mark following an M mark is a dependent mark.
- Brackets () Brackets around words or units in the mark scheme are intended to indicate wording used to clarify the mark scheme, but the marks do not depend on seeing the words or units in brackets, e.g. 10 (J) means that the mark is scored for 10, regardless of the unit given.
- Underlining Underlining indicates that this **must** be seen in the answer offered, or something very similar.
- OR / or This indicates alternative answers, any one of which is satisfactory for scoring the marks.
- e.e.o.o. This means "each error or omission".
- This means "or words to that effect". o.w.t.t.e.
- This indicates that something which is not correct or irrelevant is to be disregarded and Ignore does not cause a right plus wrong penalty.
- Spelling Be generous about spelling and use of English. If an answer can be understood to mean what we want, give credit. However, do not allow ambiguities, e.g. spelling which suggests confusion between reflection / refraction / diffraction or thermistor / transistor / transformer.
- Not / NOT This indicates that an incorrect answer is not to be disregarded, but cancels another otherwise correct alternative offered by the candidate, i.e. right plus wrong penalty applies.

Page 3	Mark Scheme	
	Cambridge IGCSE – October/November 2014	

- www.papaCambridge.com ecf meaning "error carried forward" is mainly applicable to numerical questions, but particular circumstances be applied in non-numerical questions. This indicates the candidate has made an earlier mistake and has carried an incorrect value forward to subsequent stages of working, marks indicated by ecf may be awarded, provided the subsequent working is correct, bearing in mind the earlier mistake. This prevents a candidate from being penalised more than once for a particular mistake, but only applies to marks annotated ecf.
- Sig. figs. Answers are normally acceptable to any number of significant figures \geq 2. Any exceptions to this general rule will be specified in the mark scheme. Rounding errors in the second or third significant figure will be penalised.

Arithmetic errors

Deduct one mark if the only error in arriving at a final answer is clearly an arithmetic one. Regard a power-of-ten error as an arithmetic error.

Transcription errors

Deduct one mark if the only error in arriving at a final answer is because previously calculated data has clearly been misread but used correctly.

- Fractions Allow fractions only where specified in the mark scheme.
- Units Deduct one mark for an incorrect or missing unit, but only if the answer would otherwise have gained all the marks available for that answer. Maximum one unit penalty per question.

Page	4	Mark Scheme Syl	oer oer
		Cambridge IGCSE – October/November 2014 044	10g
(a)	(i)	$(gradient =) 10 (m/s^2)$	anno
	(ii)	any linking of gradient to acceleration of freefall OR gravitational field strength	Papa Cambrid
(b)	gra	dient decreases	B1
(c)		eed/velocity stays constant OR terminal velocity/speed resultant force OR forces cancel/balance	B1 B1
(d)	gra	ally gradient steeper ph lower in second half of BC izontal final section and lower than CD	B1 B1 B1
			[Total: 8]
(a)	(i)	180 N	B1
	(ii)	(<i>P</i> =) <i>F</i> ÷ <i>A</i> OR 180÷(0.30 × 0.04) 15 000 Pa	C1 A1
(b)	(i)	arrow (labelled W) from/to correct centre of mass	B
	(ii)	1. force \times (perpendicular) distance OR 40 \times 0.60 OR 180 \times 0.15 in 2. 24 N m	C´ A´
		2. 27 Nm e.c.f. from (a)(i)	A
	(iii)	slab topples/rotates (about point D) OR corner C lifts from ground OR falls over	B1
		<u>moment</u> of force at B becomes bigger than <u>moment</u> of weight / W OR anticlockwise <u>moment</u> becomes bigger than clockwise <u>moment</u> OR weight/centre of mass outside base	B1
			[Total: 9]
(a)	(i)	(g.p.e. =) mgh OR $0.15 \times 10 \times 1.8$ 2.7 J ignore minus sign	C1 A1
	(ii)	(k.e. OR 2.7 =) $\frac{1}{2}mv^2$ OR $\frac{1}{2} \times 0.15v^2$ (v^2 =) 36 6.0 m/s	C1 C1 A1

age 5		Mark Scheme Sy	per
<u> </u>		Cambridge IGCSE – October/November 2014 044	20
(b)	(i)	<u>initial</u> temperature (of metal) OR <u>final</u> temperature (of metal) OR temperature change (of metal)	ambrid
((ii)	Mark Scheme Sy Cambridge IGCSE – October/November 2014 044 initial temperature (of metal) OR final temperature (of metal) 044 OR temperature change (of metal) 08 thermal energy transferred to something specific e.g. air/tube/stopper/ 104 thermal energy transferred to something specific e.g. air/tube/stopper/ 104 OR small spheres lost before/after weighing 09 OR not all the spheres fall the same distance 104	
		OR not all the spheres fall the same distance	B1
(i	iii)	higher temperature increase OR calculate mean of (100) readings small measurements less accurate owtte	M1 A1
			[Total: 9]
		= constant OR $p_1V_1 = p_2V_2$ OR p_1V_1/V_2 or $1.0 \times 10^5 \times 100 \div 40 \times 10^5$ Pa	C1 A1
(b)	(i)	(the particles move) <u>randomly</u>	B1
		(the particles move) slowly OR through small distances OR disappear OR zigzag OR directions change OR erratic OR straight lines between collisions	B1
(air <u>molecules/particles</u> collide with smoke particles (at high speed) fast(er) air molecules OR move randomly OR many collisions	B1 B1
(c)	diaç	gram showing:	
		ecules touching each other ecules positioned in an ordered structure	B1 B1
-	<u>Inc.</u>		[Total: 8]
(a)	(n =	-) sin <i>i</i> /sin <i>r</i> OR sin 62/sin 36	C1
	1.5((02)	C1
	(v _g - 2.0/	=) c/n OR 3.0 × 10 ⁸ /1.5 /2.00/1.997 × 10 ⁸ m/s	C1 A1
		ra-red / light) encoded OR (sent as) pulses OR multiplexing OR many messages signal OR information OR data OR internet	B1
((opt	tical fibre transmits) light/infra-red (pulse)	B1
t	tota	I internal reflection/TIR (prevents escape)	B1
			[Total: 7]
		rk (i) and (ii) together: ntion of free electrons	B1
,		rrent is) flow/movement of free electrons	B1

Page 6	Mark Scheme Syn	Dei
	Cambridge IGCSE – October/November 2014 044	230
(b) (i)	chemical (energy) to electrical (energy) (IGNORE heat)	amb
(ii)	(energy =) <i>VIt</i> OR $120 \times 96 \times 10$ (OR $\times 60$ OR $\times 10 \times 60$) OR 11520×10 (OR $\times 60$ OR $\times 10 \times 60$) 6.9×10^{6} J	baCambrids A1
(iii)	$96 \times 120~\text{OR}~1.2/1.15(2) \times 10^4~\text{OR}~12000/11500/11520$ $1.0 \times 10^4~\text{W}$	C1 A1
		[Total: 8]
(a) 150	00 m/s underlined/indicated	B1
(b) cor	mpression: closer together AND rarefaction: further apart	B1
	mpression: particles/molecules/wavefronts closer together/low pressure ID rarefaction: particles/molecules/wavefronts further apart/high pressure	B1
(c) (i)	(<i>t</i> =) <i>d</i> / <i>v</i> used OR <i>t</i> = 2 <i>d</i> / <i>v</i> OR 12/1500 OR 0.008 (s) (t =) 2 <i>d</i> /v used OR 24/1500 0.016 s	C1 C1 A1
(ii)	amplitude: decrease pitch: no change	B1 B1
		[Total: 8]
(a) 6.0	V	B1
(b) (i)	coulomb (IGNORE C)	B1
(ii)	(Q =) I t OR $0.25 \times 12 \times 60$ OR 0.25×720 OR 0.25×12 OR 3.0 OR 0.25×60 OR 15 180(C)	C1 A1
(iii)	(<i>R</i> =) V/ <i>I</i> or 6.0/0.25 or 24.0 e.c.f. from (a) OR	
	(V =) IR OR 0.25 × 16 OR 4.0 e.c.f. from (a)	C1
	8.0 Ω	A1
• •	× l OR 8.0 OR 16/2	C1
/	$R_2/(R_1 + R_2)$ OR $1/R = 1/R_1 + 1/R_2$ OR $64/16$ OR $1/R = 1/8 + 1/8$	C1
R₁F 4.0		A1

D	age	7	Mark Scheme Svi	0 DOF
10	age	<i>'</i>	Mark SchemeSyCambridge IGCSE – October/November 2014044	2D2
9	(a)	(i)	(magnetic field) lines closer together/denser/more lines	Canto
		(ii)	(magnetic field (lines) direction reversed	inde.
	(b)	(i)	ammeter needle deflects/reading on ammeter (magnetic) field cuts coil OR changing (magnetic) field (electromagnetic) <u>induction</u>	Papacambridge B1 B1 B1
		(ii)	deflection / reading on ammeter smaller OR lasts longer slower rate of cutting field lines OR slower rate of change of field	B1 B1
				[Total: 7]
10	(a)	e.g	one specific source of background radiation . rocks, ground, building materials, radon, radiation from space, Sun, smic rays, nuclear waste	B1
	(b)	(i)	electromagnetic radiation OR photons (very) high frequency OR (very) short wavelength or high energy	B1 B1
		(ii)	(count rate) decreases	B1
			(count rate decreases but) not completely absorbed (by lead) \textbf{OR} only some $\gamma\text{-rays}$ detected	B1
	(c)	(i)	no deflection (last/fifth box ticked)	B1
		(ii)	(γ -rays) are uncharged/neutral (IGNORE not affected by magnetic fields)	B1
				[Total: 7]