UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

www.papacambridge.com MARK SCHEME for the May/June 2009 question paper

for the guidance of teachers

0653 COMBINED SCIENCE

0653/03

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.


CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2009 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Syllabus er	Mark Scheme: Teachers' version	Page
0653 236	IGCSE – May/June 2009	
ABA		a) A
1	all intestine ;	E
Syllabus 0653 Babacambrid [3]	nach/small intestine ;	C
	own/digests, fats/lipids ; cids and glycerol ;	
[max 2]	ney can be absorbed ;	
[1]	, (blood) sugar/glucose ;	c) (
ore glucose/store glycogen ; [1]	es it absorb glucose/change glucose to glycogen/st	(i
[1]	ma ;	d) (
	has valves ; revent backflow of blood ;	(i
	ry has, thicker/more muscular/more elastic, wall ; to high pressure of blood ;	
[max 2]	has larger lumen ; vs easier flow ;	
[Total: 10]		
[2]	ed =) distance/time ; 0/150 = 6 km/h ;	a) (
[1]	or 1.4 ;	(i
[2]	mass x acceleration ; x 0.1 = 800 N ;	
[2]	= mass/volume <i>or</i> (mass =) density x volume ; 300 x 9 = 7200kg ;	
[2]	500 X 0 1 200 Kg ,	11

Pa	ige 3	3 Mark Scheme: Teachers' version Syllabus	s A er
		IGCSE – May/June 2009 0653	1020
(a)		orine ; oper ; on ;	www.papacambrio
(b)	(i)	sodium atom: 11 electrons arranged 2.8.1 ; oxygen atom: 8 electrons arranged 2.6 ;	[2]
	(ii)	1 more proton than electron/11p and 10e ; (other wordings possible but reject because it has lost an electron)	[1]
(c)	(i)	hydrogen + oxygen \rightarrow water ; [reject formulae]	[1]
	(ii)	reference to exothermic reaction/it melts ; gas produced (allow hydrogen)/fizzes/bubbles ; metal, dissolves/disappears ; floats ;	[max 2]
			[Total: 9]
(a)	(i)	anther/stamen ;	[1]
	(ii)	male gametes/male nuclei/male sex cells ; [ignore sperm]	[1]
(b)	(i)	the higher the temperature, the more oxygen is used ;	[1]
	(ii)	respiration ; (respiration is) aerobic/using oxygen ; (using oxygen) to produce heat ; [not to produce 'energy'] by breaking down glucose ;	[max 2]
(c)	(i)	infrared ; light ;	[2]
	(ii)	travel at same speed/transverse waves/can travel through vacuum ;	[1]
(d)	cell	cell approx. rectangular in shape, with cell wall around the outside and vacuole inside ; cell membrane labelled immediately inside the cell wall ;	
		cleus shown and labelled in cytoplasm ; oroplasts shown and labelled in cytoplasm ;	[4]

				4744
	Pa	ge 4	Mark Scheme: Teachers' version	Syllabus er
			IGCSE – May/June 2009	0653
5	(a)	(i)	P 141	Syllabus 0653 BARCambridge.com
			Q 282 R 44 S 182 ;;	
			(4 correct 1 mark)	[1]
		(ii)	4 carbons ; suitable working ;	[2]
	(b)	(i)	heated ; vaporised ; ['boiled' gets mp 1 and 2] contacted with/passed over a catalyst ;	[2]
		(ii)	M and O ; these are alkenes/contain double bonds/are unsaturated (bromine changes) from orange to colourless (not clear)	
6	(a)	(i)	(weight of empty lift = 120 00 N) (combined weight =)12 800 N ;	[1]
		(ii)	(W =) F x D <i>or</i> mgh ; = 12 800 x 9 =115 200 J ; [allow e.c.f from (i)]	[2]
	(iii)	(Power =) work/time ; = 115 200/20 = 5760 W ; [allow e.c.f from (i)]	[2]
	(b)	= 1/	= 1/R1 + 1/R2 + 1/R3 ; /2000 + 1/1000 + 1/1000 = 5/2000 ; 400 Ω;	[3] [Total: 8]



 (ii) convection ; (iii) understand the start of the	llabus ? er		Mark Scheme: Te	ge 6	Paç
 (b) solid - particles touching, regular arrangement; liquid - most particles touching, irregular arrangement; gas - few particles touching, large spaces; (c) (i) ray(s) drawn from picture to mirror to man - straight lines - angles approx. correct; arrow(s) on line(s) towards eye; (c) (i) ray(s) drawn from picture to mirror to man - straight lines - angles approx. correct; arrow(s) on line(s) towards eye; (ii) both angles required for mark; [allow e.c.f from (i) - must be consistent with arrows] [1] (iii) cannot be projected on screen/idea that brain interprets an image that is not there; [1] (ii) add excess solid; shown by mixture remaining cloudy; add solid and keep testing with indicator (paper); until mixture neutral/not acidic/specific colour with named indicator; add solid and monitor pH with a pH meter; until reading is 7/very near 7; (c) (i) positive (copper) ions are attracted to negative cathode; ions gain 2 electrons/have their charge cancelled/are discharged; (ii) oxygen has been formed; oxygen has reacted with the anode/with carbon; (to produce) carbon dioxide; 	0653	June 2009	IGCSE – May		
 (b) solid - particles touching, regular arrangement; liquid - most particles touching, irregular arrangement; gas - few particles touching, large spaces; (c) (i) ray(s) drawn from picture to mirror to man - straight lines - angles approx. correct; arrow(s) on line(s) towards eye; (c) (i) ray(s) drawn from picture to mirror to man - straight lines - angles approx. correct; arrow(s) on line(s) towards eye; (ii) both angles required for mark; [allow e.c.f from (i) - must be consistent with arrows] [1] (iii) cannot be projected on screen/idea that brain interprets an image that is not there; [1] (ii) add excess solid; shown by mixture remaining cloudy; add solid and keep testing with indicator (paper); until mixture neutral/not acidic/specific colour with named indicator; add solid and monitor pH with a pH meter; until reading is 7/very near 7; (c) (i) positive (copper) ions are attracted to negative cathode; ions gain 2 electrons/have their charge cancelled/are discharged; (ii) oxygen has been formed; oxygen has reacted with the anode/with carbon; (to produce) carbon dioxide; 	am		luction ;	(i)	(a)
 (b) solid - particles touching, regular arrangement; liquid - most particles touching, irregular arrangement; gas - few particles touching, large spaces; (c) (i) ray(s) drawn from picture to mirror to man - straight lines - angles approx. correct; arrow(s) on line(s) towards eye; (c) (i) ray(s) drawn from picture to mirror to man - straight lines - angles approx. correct; arrow(s) on line(s) towards eye; (ii) both angles required for mark; [allow e.c.f from (i) - must be consistent with arrows] [1] (iii) cannot be projected on screen/idea that brain interprets an image that is not there; [1] (ii) add excess solid; shown by mixture remaining cloudy; add solid and keep testing with indicator (paper); until mixture neutral/not acidic/specific colour with named indicator; add solid and monitor pH with a pH meter; until reading is 7/very near 7; (c) (i) positive (copper) ions are attracted to negative cathode; ions gain 2 electrons/have their charge cancelled/are discharged; (ii) oxygen has been formed; oxygen has reacted with the anode/with carbon; (to produce) carbon dioxide; 			ection ;	(ii) (
 (b) solid - particles touching, regular arrangement; liquid - most particles touching, irregular arrangement; gas - few particles touching, large spaces; (c) (i) ray(s) drawn from picture to mirror to man - straight lines - angles approx. correct; arrow(s) on line(s) towards eye; (c) (i) ray(s) drawn from picture to mirror to man - straight lines - angles approx. correct; arrow(s) on line(s) towards eye; (ii) both angles required for mark; [allow e.c.f from (i) - must be consistent with arrows] [1] (iii) cannot be projected on screen/idea that brain interprets an image that is not there; [1] (ii) add excess solid; shown by mixture remaining cloudy; add solid and keep testing with indicator (paper); until mixture neutral/not acidic/specific colour with named indicator; add solid and monitor pH with a pH meter; until reading is 7/very near 7; (c) (i) positive (copper) ions are attracted to negative cathode; ions gain 2 electrons/have their charge cancelled/are discharged; (ii) oxygen has been formed; oxygen has reacted with the anode/with carbon; (to produce) carbon dioxide; 				(iii)	(
liquid – most particles touching, irregular arrangement ; gas – few particles, not touching, large spaces ; [3] (c) (i) ray(s) drawn from picture to mirror to man – straight lines – angles approx. correct ; arrow(s) on line(s) towards eye ; [2] (ii) both angles required for mark ; [allow e.c.f from (i) – must be consistent with arrows] [1] (iii) cannot be projected on screen/idea that brain interprets an image that is not there ; [1] (iii) cannot be projected on screen/idea that brain interprets an image that is not there ; [1] (i) to speed up the reaction/so it would dissolve more quickly/because oxide less reactive than carbonate ; [1] (ii) add excess solid ; shown by mixture remaining cloudy ; add solid and keep testing with indicator (paper) ; until mixture neutral/not acidic/specific colour with named indicator ; add solid and monitor pH with a pH meter ; until reading is 7/very near 7 ; [2] (b) (CaO) + 2HCl → (CaCl ₂) + H ₂ O ;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode) ; ions gain 2 electrons/have their charge cancelled/are discharged ; [max 2] (ii) oxygen has been formed ; oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2]			eater	Ŋ	
arrow(s) on line(s) towards eye ;[2](ii) both angles required for mark ; [allow e.c.f from (i) – must be consistent with arrows] [1](iii) cannot be projected on screen/idea that brain interprets an image that is not there ;[1][Total: 10](iii) to speed up the reaction/so it would dissolve more quickly/because oxide less reactive than carbonate ;(i) to speed up the reaction/so it would dissolve more quickly/because oxide less reactive than carbonate ;(ii) add excess solid ;shown by mixture remaining cloudy ;add solid and keep testing with indicator (paper) ;until mixture neutral/not acidic/specific colour with named indicator ;add solid and monitor pH with a pH meter ;until reading is 7/very near 7 ;(b) (CaO) + 2HCl \rightarrow (CaCl ₂) + H ₂ O ;; (formulae and balanced)(c) (i) positive (copper) ions are attracted to negative cathode ;ions gain electrons (from cathode) ;ions gain 2 electrons/have their charge cancelled/are discharged ;(ii) oxygen has been formed ;oxygen has reacted with the anode/with carbon ;(to produce) carbon dioxide ;(max 2]		gular arrangement ;	nost particles touching, irre	liquio	. ,
(iii) cannot be projected on screen/idea that brain interprets an image that is not there ; [1] [[Total: 10] (a) (i) to speed up the reaction/so it would dissolve more quickly/because oxide less reactive than carbonate ; [1] (ii) add excess solid ; shown by mixture remaining cloudy ; add solid and keep testing with indicator (paper) ; until mixture neutral/not acidic/specific colour with named indicator ; add solid and monitor pH with a pH meter ; until reading is 7/very near 7 ; [2] (b) (CaO) + 2HCl \rightarrow (CaCl ₂) + H ₂ O ;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode) ; ions gain 2 electrons/have their charge cancelled/are discharged ; [max 2] (ii) oxygen has been formed ; oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2]	lles approx. correct ;	-	, .	• •	(c)
(a) (i) to speed up the reaction/so it would dissolve more quickly/because oxide less reactive than carbonate ; [1] (ii) add excess solid ; [1] (iii) caco + 2HCl \rightarrow (CaCl ₂) + H ₂ O ;; (formulae and balanced) [2] (b) (CaO) + 2HCl \rightarrow (CaCl ₂) + H ₂ O ;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; [1] ions gain electrons (from cathode) ; [1] (ii) oxygen has been formed ; [1] oxygen has reacted with the anode/with carbon ; [1] (iv) produce) carbon dioxide ; [1]	onsistent with arrows]	[allow e.c.f from (i) - must be	angles required for mark	(ii)	
(a) (i) to speed up the reaction/so it would dissolve more quickly/because oxide less reactive than carbonate ; [1] (ii) add excess solid ; [1] (iii) caco + 2HCl \rightarrow (CaCl ₂) + H ₂ O ;; (formulae and balanced) [2] (b) (CaO) + 2HCl \rightarrow (CaCl ₂) + H ₂ O ;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; [1] ions gain electrons (from cathode) ; [1] (ii) oxygen has been formed ; [1] oxygen has reacted with the anode/with carbon ; [1] (iv) produce) carbon dioxide ; [1]	ge that is not there :	dea that brain interprets an ir	ot be projected on screen	(iii)	(
(a) (i) to speed up the reaction/so it would dissolve more quickly/because oxide less reactive than carbonate ; [1] (ii) add excess solid ; shown by mixture remaining cloudy ; add solid and keep testing with indicator (paper) ; until mixture neutral/not acidic/specific colour with named indicator ; add solid and monitor pH with a pH meter ; until reading is 7/very near 7 ; [2] (b) (CaO) + 2HCl \rightarrow (CaCl ₂) + H ₂ O ;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode) ; ions gain 2 electrons/have their charge cancelled/are discharged ; [max 2] (ii) oxygen has been formed ; oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2]				,	ſ
than carbonate ;[1](ii) add excess solid ; shown by mixture remaining cloudy ; add solid and keep testing with indicator (paper) ; until mixture neutral/not acidic/specific colour with named indicator ; add solid and monitor pH with a pH meter ; until reading is 7/very near 7 ;[2](b) $(CaO) + 2HCl \rightarrow (CaCl_2) + H_2O$;; (formulae and balanced)[2](c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode) ; ions gain 2 electrons/have their charge cancelled/are discharged ;[max 2](ii) oxygen has been formed ; oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ;[max 2]					
shown by mixture remaining cloudy ; add solid and keep testing with indicator (paper) ; until mixture neutral/not acidic/specific colour with named indicator ; add solid and monitor pH with a pH meter ; until reading is 7/very near 7 ; [2] (b) (CaO) + 2HCl \rightarrow (CaCl ₂) + H ₂ O ;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode) ; ions gain 2 electrons/have their charge cancelled/are discharged ; [max 2] (ii) oxygen has been formed ; oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2]	cause oxide less react	would dissolve more quickly/			(a)
until mixture neutral/not acidic/specific colour with named indicator ; add solid and monitor pH with a pH meter ; until reading is 7/very near 7 ; [2] (b) $(CaO) + 2HCl \rightarrow (CaCl_2) + H_2O$;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode) ; ions gain 2 electrons/have their charge cancelled/are discharged ; [max 2] (ii) oxygen has been formed ; oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2]		pudy ;			
until reading is 7/very near 7; [2] (b) $(CaO) + 2HCl \rightarrow (CaCl_2) + H_2O$;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode); ions gain 2 electrons/have their charge cancelled/are discharged; [max 2] (ii) oxygen has been formed; oxygen has reacted with the anode/with carbon; (to produce) carbon dioxide; [max 2]	ator ;				
(b) $(CaO) + 2HCl \rightarrow (CaCl_2) + H_2O$;; (formulae and balanced) [2] (c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode) ; ions gain 2 electrons/have their charge cancelled/are discharged ; [max 2] (ii) oxygen has been formed ; oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2]		ı pH meter ;	•		
 (c) (i) positive (copper) ions are attracted to negative cathode ; ions gain electrons (from cathode) ; ions gain 2 electrons/have their charge cancelled/are discharged ; [max 2] (ii) oxygen has been formed ; oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2] 			reading is 7/very near 7;	I	
 ions gain electrons (from cathode); ions gain 2 electrons/have their charge cancelled/are discharged; [max 2] (ii) oxygen has been formed; oxygen has reacted with the anode/with carbon; (to produce) carbon dioxide; [max 2] 		H_2O ;; (formulae and balance)	· 2HC $l \rightarrow$ (CaC l_2) +	(CaC	(b)
oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2]	ed ; [max	de);	gain electrons (from catho	i	(c)
oxygen has reacted with the anode/with carbon ; (to produce) carbon dioxide ; [max 2]			en has been formed ·	(ii) (
	_	ode/with carbon ;	en has reacted with the a	(
(iii) 2 pairs of shared electrons and two lone pairs on each atom ; [1]	[max		roduce) carbon dioxide ;		
					,