

	UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINAT International General Certificate of Secondary Education	tions data and the com
CANDIDATE NAME		5 ¹⁷⁷
CENTRE NUMBER	CANDIDATE	
COMBINED S	CIENCE	0653/63
Paper 6 Alterna	tive to Practical	May/June 2013

1 hour

Candidates answer on the Question paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Exam	iner's Use
1	
2	
3	
4	
5	
6	
Total	
3 4 5 6	

This document consists of 19 printed pages and 1 blank page.



www.papaCambridge.com 1 A student did an experiment to investigate the effect of temperature on the respiration of yeast. This was done by measuring the rate at which the yeast produ bubbles of carbon dioxide. Yeast is able to respire, and produce carbon dioxide, even in the absence of oxygen.

Apparatus was set up as shown in Fig. 1.1.




Fig. 1.1

The student made sure that the temperature of the water bath was 20°C, and then waited for three minutes. After this time, the number of bubbles produced was counted each minute for three minutes.


The procedure was then repeated at 30 °C, at 40 °C, at 60 °C, and finally at 80 °C.

The results are shown in Table 1.1.

tomporaturo /°C	number of bubbles in one minute						
temperature/°C	1 st minute	2 nd minute	3 rd minute	average			
20	3	5	4	4			
30	10	7	7				
40	15	13	19				
60	15	16	14				
80	0	0	0	0			

Table	1	.1
-------	---	----

(a) Complete Table 1.1, by working out the average numbers of bubbles produced per minute at 30 °C, 40 °C, and 60 °C giving each of your answers to the nearest whole number. [1] (b) On the grid below, plot the average number of bubbles per minute against temper Join your points with straight lines.



		4	Camprilige Com
(d)	Exp	plain why	For For
	(i)	the student waited three minutes before taking readings at each temperature,	nonida en s
			 1]
	(ii)	three readings were taken at each temperature,	
			I
	(iii)	the experiment at 80 °C was done last.	
			[1]
(e)		ggest how this experiment could be modified to show that the bubbles of gas giv contain carbon dioxide.	en
	•••••		[1]
(f)	Sug	ggest a suitable control experiment for this investigation.	
	•••••		[1]

2 A student is investigating the characteristics of a converging lens by two different me

www.papaCambridge.com A converging lens is a lens that is thicker in the middle. It bends rays of light together that they can form an image on a screen, as shown in Fig. 2.1.

Method 1

She sets up a converging lens so that light from a distant object produces a sharp image on a screen as shown in Fig. 2.1.




Fig. 2.1

(a) The distance f, from the screen to the lens, is called the focal length. Use a ruler to measure the distance f.

f = _____ cm [1]

Method 2

The student sets up the apparatus shown in Fig. 2.2. The light shines through the hole and the screen can be moved until a sharp image of the illuminated object is formed.




Fig. 2.2

She moves the lens so that distance, *u*, is 50.0 cm. She moves the screen until a sharp image is seen. She measures distance, v, and records the value in Table 2.1. She repeats this for different values of **u**.

	6 Table 2		43.41	n Papar
u /cm	v /cm	<i>u+v</i> /cm	uv/cm^2	
20.0	59.8	78.8	1196	
30.0	30.1	60.1	903	
40.0				
45.0	22.7	67.7	1021	
50.0	21.5	71.5	1075	

Fig. 2.3 shows a diagram of the apparatus when distance u = 40.0 cm.

The diagram is drawn to the scale of 1 to 10. That means that 1 cm in the diagram is actually 10 cm.





Fig. 2.3

(b) (i) Use a ruler to measure the distance for v in Fig. 2.3, to the nearest 0.1 cm.

distance = _____ cm [1]

- (ii) Convert this measurement to the actual value of v and complete the second column of Table 2.1. [1]
- (iii) Complete column 3 of Table 2.1 by adding *u* and *v*. [1]
- (iv) Complete the final column of Table 2.1 by multiplying *u* and *v*. [1]



3 In this experiment a student is investigating how the rate of the reaction b magnesium and acid is affected by the surface area of the solid. She sets up the appart as in Fig. 3.1.




Fig. 3.1


- She removes the bung from the large test-tube and places 1 cm of magnesium ribbon • in the bottom of the test-tube.
- She measures out 10 cm^3 acid. •
- She starts the stopclock. •
- When the time on the stopclock reaches 10 seconds, she adds the acid to the test-tube and quickly replaces the bung as tightly as possible.
- She stops the clock when the volume of gas collected in the inverted measuring . cylinder is 10 cm³.
- She subtracts 10 seconds from the time displayed on the clock, then records the result in Table 3.1.

length of magnesium/cm	time/s	rate/cm ³ per s
1	37.5	
2	19.0	
3		
4		

Table 3.1

She repeats the above steps using 2 cm of magnesium ribbon. She subtracts 10 seconds from the time shown on the clock and records the result in Table 3.1.

www.papaCambridge.com (a) Fig. 3.2 shows the stopclock readings for lengths of magnesium 3 cm and 4 cm. Subtract 10 seconds from each and record the result in Table 3.1.





(b) (i) For each length of magnesium, complete column 3 of Table 3.1 by calculating the rate of the reaction using the formula

rate =
$$\frac{10}{\text{time}}$$

[1]

[2]

(ii) How does the rate of the reaction between magnesium and acid vary with the length of magnesium ribbon?

......[1]

(iii) Since the magnesium ribbon is thin, doubling the length of the ribbon approximately doubles the surface area.

How does the rate of the reaction between magnesium and acid vary with the surface area of the magnesium ribbon?

.....[1]

		424
	10	1. D
	(iv) The teacher said 'The rate of reaction doubles when the length ribbon is doubled'.	of the magn
	State whether the results show that the statement is correct or i show which results support your statement.	n of the magnine Cannin
		[3]
(c)) Suggest why she waited 10 seconds before adding the acid.	
		[1]
(d)) The student suspects that the gas collected in the reaction is hydrog	len.
	Suggest how the student should test the gas collected in this experinities is hydrogen.	ment to show that it

......[1]

4 The enzyme pectinase is used in industry for preparing fruit juices. Pectinase speeds release of fruit juice from plant cells by breaking down the cell walls.


www.papaCambridge.com A student used apples as the source of the plant cells. He wanted to produce a clear fruit juice.

- He placed some apple pieces and a little water in a blender and used this to produce a smooth apple paste.
- He then placed equal volumes of apple paste in each tube.
- He then placed pectinase and water in the tubes as shown in Table 4.1.
- He incubated the tubes at the temperatures indicated for ten minutes.

tube number	apple paste/cm ³	pectinase/cm ³	water/cm ³	temperature/°C
1	5	1	0	20
2	5	0	1	20
3	5	1	0	40
4	5	0	1	40

Table 4.1

Results






(a) (i) Pectinase was not included in tubes 2 and 4.

Explain why tubes 2 and 4 were included in the experiment.

.....[1]




13
(iii) The student modified his original experiment to make the fruit juice as of the trut provide the first possible.
Describe in detail what he did and how he could prove that the enzyme named in
(b)(ii) was effective.
[]]
[]]
[]]
[]]
[]]

© UCLES 2013

5 (a) A motor manufacturer is testing his new electric car.

> The driver is given instructions on how to drive over a set distance on a special te track, as shown in Fig. 5.1.

www.papaCambridge.com Poles are placed 10 m apart and a photograph of the position of the car is taken every second.





The distances for one test run are recorded in Table 5.1.

Table 5.1

time/s	0	1	2	3	4	5	6	7	8	9	10
distance/m	0	8	18	34	52		99		161	199	239

(i) Use Fig. 5.2 to record in Table 5.1 the distances travelled after 5 and 7 seconds.

[2]

Take your measurement from the front of the car.






Fig. 5.2







Please turn over for Question 6.

(a) Fig. 6.1 shows words and phrases about different anion tests cut from a page 6 student's note book.

www.papaCambridge.com Use a ruler to construct a table, showing each anion, test reagent and result which identifies the anion.

barium chloride

chLoride

carbonate

bubbles


white precipitate

silver nitrate

yellow precipitate

dilute hydrochloric acid

Fig. 6.1





BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.