## **CAMBRIDGE INTERNATIONAL EXAMINATIONS**

**Cambridge International General Certificate of Secondary Education** 

## MARK SCHEME for the May/June 2015 series

## 0653 COMBINED SCIENCE

0653/51

Paper 5 (Practical Test), maximum raw mark 30

www.PapaCambridge.com

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2015 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

Page 2	Mark Scheme	Sy. Sper
	Cambridge IGCSE – May/June 2015	065

1 (a) outline concave on one side;

projections on the other side;

(b) (i)

test solution	observation	
Benedict's reagent	green/yellow/orange/red;	
biuret solution	blue/no change/colour stays the same ;	
iodine solution	brown/orange/no change/colour stays the same;	

[3]


(ii) reducing sugar / glucose;(NOT sugar. DO NOT ALLOW additional food groups)

[1]

(c) (i) several small circles labelled 'stained' or 'coloured' or (c)(i) or red;

[1]

[1]



(ii) water transport; (ALLOW water and any idea of movement, 'absorbs water' is not enough)

(d) different temperatures in separate experiments;

time for coloured water to appear at top of cut stem/set time and measure distance moved ;

all other conditions/named condition kept constant;

(if one experiment proposed with gradual increase in temperature then can only score 2nd marking point)

[max 2]

(a) (i) white ppt/milky/cloudy white; (ii) blue/purple AND pH value in range 8–14; (iii) calcium oxide/CaO; (ALLOW quicklime/limewater/calcium hydroxide/Ca(OH)₂ note: accept answer if seen in (iv)  (iv) base/basic/alkali/alkaline;  [iii) blue (and white) ppt.; (some) ppt dissolves soluble in excess (ammonia); to form darker blue (solution);  (iii) Cu²²-(Cu(II)/copper (not Cu); copper oxide/CuO; note: both marks depend on 'blue' being reported in (b)(i) or (b)(ii)  (a) all t values present and increasing; (ALLOW 0:12 format)  T values correct minimum 2 sig. fig.; (if 0:12 format used for t then T = 12/20 NOT 0.12/20) (ALLOW ecf from T)  T² values correct AND to 2 sig. fig.; (ALLOW ecf from T)  (b) (i) suitable choice of scales from (0.0) using at least half of each axis (m likely to be 0.1 per 2cm); at least three plots correct to ± ½ small square; good best-fit straight line judgement; (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)  (iii) indication on graph of how data obtained AND ⇒ half the line used; correct calculation using data from graph; (iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;	Page	3	Mark Scheme	Sy. Sy. per
(ALLOW quicklime/limewater/calcium hydroxide/Ca(OH) <sub>2</sub> note: accept answer if seen in (iv)  (iv) base/basic/alkali/alkaline;  (ii) blue (and white) ppt.; (some) ppt dissolves soluble in excess (ammonia); to form darker blue (solution);  (iii) Cu²*/Cu(II)/copper (not Cu); copper oxide/CuO; note: both marks depend on 'blue' being reported in (b)(i) or (b)(ii)  (a) all t values present and increasing; (ALLOW 0:12 format)  T values correct minimum 2 sig. fig.; (if 0:12 format used for t then T = 12/20 NOT 0.12/20) (ALLOW ecf from T)  T² values correct AND to 2 sig. fig.; (ALLOW ecf from T)  (b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm); at least three plots correct to ± ½ small square; good best-fit straight line judgement; (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)  (ii) indication on graph of how data obtained AND ⇒ half the line used; correct calculation using data from graph; (iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;			Cambridge IGCSE – May/June 2015	065
(ALLOW quicklime/limewater/calcium hydroxide/Ca(OH) <sub>2</sub> note: accept answer if seen in (iv)  (iv) base/basic/alkali/alkaline;  (ii) blue (and white) ppt.; (some) ppt dissolves soluble in excess (ammonia); to form darker blue (solution);  (iii) Cu²*/Cu(II)/copper (not Cu); copper oxide/CuO; note: both marks depend on 'blue' being reported in (b)(i) or (b)(ii)  (a) all t values present and increasing; (ALLOW 0:12 format)  T values correct minimum 2 sig. fig.; (if 0:12 format used for t then T = 12/20 NOT 0.12/20) (ALLOW ecf from T)  T² values correct AND to 2 sig. fig.; (ALLOW ecf from T)  (b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm); at least three plots correct to ± ½ small square; good best-fit straight line judgement; (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)  (ii) indication on graph of how data obtained AND ⇒ half the line used; correct calculation using data from graph; (iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;	(a)	) (i)	white ppt/milky/cloudy white;	and
(ALLOW quicklime/limewater/calcium hydroxide/Ca(OH) <sub>2</sub> note: accept answer if seen in (iv)  (iv) base/basic/alkali/alkaline;  (ii) blue (and white) ppt.; (some) ppt dissolves soluble in excess (ammonia); to form darker blue (solution);  (iii) Cu²*/Cu(II)/copper (not Cu); copper oxide/CuO; note: both marks depend on 'blue' being reported in (b)(i) or (b)(ii)  (a) all t values present and increasing; (ALLOW 0:12 format)  T values correct minimum 2 sig. fig.; (if 0:12 format used for t then T = 12/20 NOT 0.12/20) (ALLOW ecf from T)  T² values correct AND to 2 sig. fig.; (ALLOW ecf from T)  (b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm); at least three plots correct to ± ½ small square; good best-fit straight line judgement; (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)  (ii) indication on graph of how data obtained AND ⇒ half the line used; correct calculation using data from graph; (iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;		(ii)	blue/purple <b>AND</b> pH value in range 8–14;	Tig
<ul> <li>(b) (i) blue (and white) ppt.;</li> <li>(ii) blue (and white) ppt.;</li> <li>(some) ppt dissolves soluble in excess (ammonia);</li> <li>to form darker blue (solution);</li> <li>(iii) Cu²*/Cu(II)/copper (not Cu);</li> <li>copper oxide/CuO;</li> <li>note: both marks depend on 'blue' being reported in (b)(i) or (b)(ii)</li> <li>(a) all t values present and increasing;</li> <li>(ALLOW 0:12 format)</li> <li>T values correct minimum 2 sig. fig.;</li> <li>(if 0:12 format used for t then T = 12/20 NOT 0.12/20)</li> <li>(ALLOW ecf from T)</li> <li>T² values correct AND to 2 sig. fig.;</li> <li>(ALLOW ecf from T)</li> <li>(b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm);</li> <li>at least three plots correct to ± ½ small square;</li> <li>good best-fit straight line judgement;</li> <li>(if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(iii) indication on graph of how data obtained AND &gt;&gt; half the line used;</li> <li>correct calculation using data from graph;</li> <li>(iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;</li> </ul>		(iii)	(ALLOW quicklime/limewater/calcium hydroxide/Ca(OH) <sub>2</sub>	[1]
<ul> <li>(ii) blue (and white) ppt.; (some) ppt dissolves soluble in excess (ammonia); to form darker blue (solution);</li> <li>(iii) Cu²²/Cu(II) / copper (not Cu); copper oxide / CuO; note: both marks depend on 'blue' being reported in (b)(i) or (b)(ii)</li> <li>(a) all t values present and increasing; (ALLOW 0:12 format)</li> <li>T values correct minimum 2 sig. fig.; (if 0:12 format used for t then T = 12/20 NOT 0.12/20) (ALLOW ecf from T)</li> <li>T² values correct AND to 2 sig. fig.; (ALLOW ecf from T)</li> <li>(b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm); at least three plots correct to ± ½ small square; good best-fit straight line judgement; (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ≥ half the line used; correct calculation using data from graph; [iii) correct calculation of k to 2/3 sig. fig. and correctly rounded; [iiii)</li> </ul>		(iv)	base/basic/alkali/alkaline;	[1]
<ul> <li>(some) ppt dissolves soluble in excess (ammonia); to form darker blue (solution);</li> <li>(iii) Cu²+/Cu(II) / copper (not Cu); copper oxide / CuO; note: both marks depend on 'blue' being reported in (b)(i) or (b)(ii)</li> <li>(a) all t values present and increasing; (ALLOW 0:12 format)  T values correct minimum 2 sig. fig.; (if 0:12 format used for t then T = 12/20 NOT 0.12/20) (ALLOW ecf from T)  T² values correct AND to 2 sig. fig.; (ALLOW ecf from T)</li> <li>(b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm); at least three plots correct to ± ½ small square; good best-fit straight line judgement; (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ⇒ half the line used; correct calculation using data from graph;</li> <li>(iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;</li> </ul>	(b)	) (i)	blue (and white) ppt.;	[1]
copper oxide / CuO; note: both marks depend on 'blue' being reported in (b)(i) or (b)(ii)  (a) all t values present and increasing;  (ALLOW 0:12 format)  T values correct minimum 2 sig. fig.;  (if 0:12 format used for t then T = 12/20 NOT 0.12/20)  (ALLOW ecf from T)  T² values correct AND to 2 sig. fig.;  (ALLOW ecf from T)  (b) (i) suitable choice of scales from (0,0) using at least half of each axis  (m likely to be 0.1 per 2cm);  at least three plots correct to ± ½ small square;  good best-fit straight line judgement;  (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)  (ii) indication on graph of how data obtained AND ⇒ half the line used;  correct calculation using data from graph;  (iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;		(ii)	(some) ppt dissolves soluble in excess (ammonia);	[3]
(ALLOW 0:12 format)  T values correct minimum 2 sig. fig.;  (if 0:12 format used for t then T = 12/20 NOT 0.12/20)  (ALLOW ecf from T)  T² values correct AND to 2 sig. fig.;  (ALLOW ecf from T)  (b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm);  at least three plots correct to ± ½ small square;  good best-fit straight line judgement;  (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)  (ii) indication on graph of how data obtained AND ≥ half the line used;  correct calculation using data from graph;  (iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;		(iii)	copper oxide/CuO;	[2]
<ul> <li>T values correct minimum 2 sig. fig.;</li> <li>(if 0:12 format used for t then T = 12/20 NOT 0.12/20)</li> <li>(ALLOW ecf from T)</li> <li>T² values correct AND to 2 sig. fig.;</li> <li>(ALLOW ecf from T)</li> <li>(b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm);</li> <li>at least three plots correct to ± ½ small square;</li> <li>good best-fit straight line judgement;</li> <li>(if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ≥ half the line used;</li> <li>correct calculation using data from graph;</li> <li>(iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;</li> </ul>	(a)	) all t	values present and increasing ;	
<ul> <li>(if 0:12 format used for t then T = 12/20 NOT 0.12/20)</li> <li>(ALLOW ecf from T)</li> <li>T² values correct AND to 2 sig. fig.;</li> <li>(ALLOW ecf from T)</li> <li>(b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm);</li> <li>at least three plots correct to ± ½ small square;</li> <li>good best-fit straight line judgement;</li> <li>(if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ≥ half the line used;</li> <li>correct calculation using data from graph;</li> <li>(iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;</li> </ul>			(ALLOW 0:12 format)	
<ul> <li>(ALLOW ecf from T)</li> <li>T² values correct AND to 2 sig. fig.;</li> <li>(ALLOW ecf from T)</li> <li>(b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm);</li> <li>at least three plots correct to ± ½ small square;</li> <li>good best-fit straight line judgement;</li> <li>(if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ≥ half the line used;</li> <li>correct calculation using data from graph;</li> <li>(iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;</li> </ul>		Tv	alues correct minimum 2 sig. fig. ;	
<ul> <li>(ALLOW ecf from T)</li> <li>(b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm); at least three plots correct to ± ½ small square; good best-fit straight line judgement; (if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ≥ half the line used; correct calculation using data from graph; (iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;</li> </ul>			· ·	
<ul> <li>(b) (i) suitable choice of scales from (0,0) using at least half of each axis (m likely to be 0.1 per 2cm); at least three plots correct to ± ½ small square; good best-fit straight line judgement; [if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ⇒ half the line used; correct calculation using data from graph; [iii) correct calculation of k to 2/3 sig. fig. and correctly rounded; [iiii)</li> </ul>		$T^2$ \	values correct AND to 2 sig. fig. ;	[3]
<ul> <li>(m likely to be 0.1 per 2cm);</li> <li>at least three plots correct to ± ½ small square;</li> <li>good best-fit straight line judgement;</li> <li>(if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ≥ half the line used;</li> <li>correct calculation using data from graph;</li> <li>(iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;</li> </ul>			(ALLOW ecf from T)	
<ul> <li>good best-fit straight line judgement;</li> <li>(if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ≥ half the line used;</li> <li>correct calculation using data from graph;</li> <li>(iii) correct calculation of k to 2/3 sig. fig. and correctly rounded;</li> </ul>	(b)	) (i)	` ,	
<ul> <li>(if non-linear then do not award scale, plot or line marks EXCEPT if non-linear region is just between 0 and 0.2 kg then do not award scale mark)</li> <li>(ii) indication on graph of how data obtained AND ⇒ half the line used; correct calculation using data from graph; [</li> <li>(iii) correct calculation of k to 2/3 sig. fig. and correctly rounded; [</li> </ul>			at least three plots correct to ± ½ small square ;	
region is just between 0 and 0.2 kg then do not award scale mark)  (ii) indication on graph of how data obtained AND ⇒ half the line used;  correct calculation using data from graph;  [iii) correct calculation of <i>k</i> to 2/3 sig. fig. and correctly rounded;  [iii) region is just between 0 and 0.2 kg then do not award scale mark)			good best-fit straight line judgement;	[3]
correct calculation using data from graph;  [iii) correct calculation of $k$ to 2/3 sig. fig. and correctly rounded;			· ·	if non-linear
(iii) correct calculation of $k$ to 2/3 sig. fig. and correctly rounded;		(ii)	indication on graph of how data obtained AND ≥ half the line used	;
			correct calculation using data from graph;	[2]
(c) eye level with top/bottom/middle of oscillations/equivalent; [		(iii)	correct calculation of $k$ to 2/3 sig. fig. and correctly rounded;	[1]
	(c)	) eye	e level with <b>top/bottom/middle</b> of oscillations/equivalent;	[1]