



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

Original Constitution

| * | |
|---|--|
| 6 | |
| ∞ | |
| J | |
| 9 | |
| 6 | |
| 6 | |
| J | |
| 6 | |
| J | |
| ū | |

| CANDIDATE NAME | | | | | | |
|-------------------|--|--|-----------------|--|--|--|
| CENTRE NUMBER | | | CANDIE NUMBE | | | |

CO-ORDINATED SCIENCES

0654/63

Paper 6 Alternative to Practical

May/June 2010

1 hour

Candidates answer on the Question paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs, tables or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

| For Examiner's Use | |
|--------------------|--|
| 1 | |
| 2 | |
| 3 | |
| 4 | |
| 5 | |
| 6 | |
| Total | |

This document consists of 18 printed pages and 2 blank pages.



- 1 This question is about variation in leaves.
- www.PapaCambridge.com (a) A student was provided with the 20 leaves shown in Fig. 1.2. Measure the length \vec{l} each leaf in millimetres as shown in Fig. 1.1a. If the lamina does not meet the petiole evenly on either side of the leaf use the longer measurement. See Fig. 1.1b.

Enter your measurements in Table 1.1.




Fig. 1.1a




Fig. 1.1b

Table 1.1

| | length of leaf // mm | | |
|----|----------------------|--|--|
| 1 | 11 | | |
| 2 | 12 | | |
| 3 | 13 | | |
| 4 | 14 | | |
| 5 | 15 | | |
| 6 | 16 | | |
| 7 | 17 | | |
| 8 | 18 | | |
| 9 | 19 | | |
| 10 | 20 | | |




Fig. 1.2

(b) Calculate the average (mean) length of the 20 leaves. Show your working.

average = _____ mm


[2]

| range / mm | number of leaves in range |
|------------|---------------------------|
| 30 - 34 | |
| 35 - 39 | |
| 40 - 44 | |
| 45 - 49 | |
| 50 - 54 | |
| 55 - 59 | |

[2]

www.PapaCambridge.com

(ii) Using the information you have entered in Table 1.2 draw a bar chart on the grid provided. Use the **number of leaves in range** as the vertical axis and the **range / mm** as the horizontal axis. Choose suitable scales for your data.



[3]

| (d) | The leaves were all of the same species yet they showed variation in length. Suggest reason for this. | t a |
|-----|---|-----|
| | | |
| | | [1] |

2 Some science students are making ammonia gas, NH₃. They are mixing solid chydroxide, Ca(OH)₂ and ammonium chloride, NH₄C*I*. Then they are heating the mixtuke the apparatus shown in Fig. 2.1.






Fig. 2.1

(a) (i) State the colour of ammonia gas.

[1]

(ii) Water, ammonia and a solid substance are formed when calcium hydroxide reacts with ammonium chloride.

Name the solid substance that is formed. Do not write its formula.

[1]

(b) The students have to decide how to collect the gas that will come out of the delivery tube. The teacher gives them the information shown in Table 2.1.

Table 2.1

| name of gas | air | ammonia |
|---------------------------------------|-----|---------|
| density of gas in g / dm ³ | 1.2 | 0.7 |

Three methods of collecting a gas are shown in Fig. 2.2.




Fig. 2.2

| (i) | Study methods A , B and C in Fig. 2.2. Choose the best method of commonia gas. Use the information in Table 2.1 and your knowledge of properties of ammonia to help you to choose. |
|------|---|
| | The best way to collect ammonia gas is method[1] |
| (ii) | Choose one of the other two methods and complete the sentence below. |
| | Method is not a good way to collect ammonia gas |
| | because |
| | |
| | [1] |
| | |

(c) The students have collected some gas-jars of ammonia. They are testing the gas using the tests shown in Table 2.2.

Complete Table 2.2.

Table 2.2

| | test | result |
|-------|--|---|
| (i) | A few cm ³ of aqueous [1] sulfate is added to the gas. | A white precipitate is formed. After shaking, the precipitate re-dissolves forming a colourless solution. |
| (ii) | A few cm ³ of aqueous copper(II) sulfate is added to the gas. | A precipitate is formed which has a colour. After shaking, the precipitate re-dissolves forming a [2] solution. |
| (iii) | A few drops of litmus is added. | The litmus turns |
| | | [1] |

For iner's

www.PapaCambridge.com (d) The teacher carries out an experiment with the gas-jar of ammonia. He soaks a paper in concentrated hydrochloric acid. Then he holds the filter paper at the mouth. the gas-jar of ammonia. This is shown in Fig 2.3.




Fig. 2.3

| [2] |
|-----|

www.papaCambridge.com A science student is trying to find out how much a liquid expands when it is turned 3 gas. He is using water as the liquid. The apparatus is shown in Fig 3.1.





Fig. 3.1

- He weighs a clean, dry plastic 2 dm³ bottle with its lid on and records the mass, m₁.
- He removes the lid and places the bottle in position as in Fig. 3.1.
- He boils the water and passes steam into the bottle for three or four minutes until the bottle is at 100 °C and no drops of water remain in it. The bottle is now full of steam.
- He quickly removes the bottle from its stand, places it upright and loosely replaces the lid.
- When the bottle has cooled to room temperature he weighs it again and records the mass, m₂.

The balance windows for the masses, m_1 and m_2 are shown in Fig. 3.2.






Fig. 3.2

(a) (i) Read and record the masses of the bottle, m_1 and m_2 , before and after passing steam into it.

mass
$$m_1$$
= _____ g

mass
$$m_2$$
= _____g [2]

| ĺ | 'ii' | Find the | increase | in | mass | ∩f | the | hottle | |
|---|------|---------------|-----------|-----|------|----|-----|--------|---|
| ١ | ш | , i ilia ilie | IIICICasc | 111 | mass | Oi | uic | DOLLIG | ٠ |

| C. Tan |
|------------|
| For iner's |
| OJ. |
| 8 |

| | | | increase in mass o | f the bottle= | | g | [1] |
|-----|-----------------|---------------------|--|---------------------------|-------------------|-----------------|------|
| (b) | Cha | anging water into s | team is a reversible | process. | | | |
| | | _ | process | A | → | | |
| | | water | process | s B | steam | | |
| | (i) | Name process A | · | | | | [1] |
| | (ii) | Name process E | | | | | [1] |
| | e ma: a)(ii) | | ed when the 2 dm ³ | of steam co | oled down is give | n in your ans | wer |
| 1 g | of wa | ater has a volume | of 1 cm ³ . | | | | |
| (c) | (i) | What volume of v | ater was produced | | | | |
| | | | volum | e of water = | | cm ³ | [1] |
| | (ii) | | ume produced, in o | | etres, when 1 cu | bic centimetre | e of |
| | | | | | | | |
| | | volume | of steam from 1 cm | າ ³ of water = | | cm³ | [2] |
| | (d) | | e, water is heated to powerful force is pro | | | f this experim | ent |
| | | | | | | | |
| | | | | | | | |
| | | | | | | | [2] |

- 4 This question compares the content of inhaled air with exhaled air. The experime done using gas-jars of inhaled and exhaled air.
- www.papaCambridge.com (a) Gas-jars that had been standing open in the laboratory were used for samples of inhaled air.

In the space below draw a labelled diagram to show how you would collect a sample of exhaled air in a gas-jar.

[2]

(b) To compare the oxygen content of the air a student lit a candle and placed it in the gas-jar. The length of time that the flame burned was noted and the results were entered in Table 4.1. The experiment was repeated three times with each type of air.

Table 4.1

| source of air | time of burning/s | | | | | |
|---------------|-------------------|--------|--------|---------|--|--|
| sample | test 1 | test 2 | test 3 | average | | |
| inhaled air | | 7.0 | 6.5 | | | |
| exhaled air | 5.0 | | 4.5 | | | |

(i) Read the stop watches in Fig. 4.1 and enter the results in Table 4.1.



inhaled air, test 1 time/s


55

10

15

50

45



55

50

45



[2]

| | | | The state of the s | | | | |
|-------|---|---|--|--|--|--|--|
| | | 11 | To a | | | | |
| (ii) | Calculate the average time of bu your results in Table 4.1. | irning of the candle in each sample | e of air an | | | | |
| | | | [2] | | | | |
| | compare the carbon dioxide c tinuously shook a fixed volume of | ontent of inhaled and exhaled limewater in the gas-jar of air. | air the student | | | | |
| (i) | _ | nhaled air after shaking the samp ewater changed appearance with | | | | | |
| | What was observed in the gas-ja | ar of exhaled air? | | | | | |
| | | | [1] | | | | |
| (ii) | The carbon dioxide content of air is increased by the lungs. What process in the body produces carbon dioxide? | | | | | | |
| | [1] | | | | | | |
| (iii) | The student then shook fresh limewater with more samples of exhaled air. One sample was collected before exercise and the other after exercise . The student forgot which sample was which. The times taken for the limewater to change were 3.2s and 8.4s. | | | | | | |
| | Enter the times in the correct pla | ce in Table 4.2. | | | | | |
| | Table 4.2 | | | | | | |
| | exhaled air sampl | e time taken for change/s | | | | | |
| | before exercise | | | | | | |
| | after exercise | | | | | | |
| | | | [1] | | | | |
| (iv) | Explain why exercise changes the | ne carbon dioxide content of exhal | ed air. | | | | |
| | | | | | | | |
| | | | | | | | |
| | | | [41 | | | | |

5 A student is investigating the rate of reaction between magnesium and dilute hydro acid. She does several experiments using different concentrations of the acid with the sa lengths of magnesium ribbon. Fig. 5.1 shows the apparatus she is using.




Fig. 5.1

- She measures out the volume of dilute hydrochloric acid for experiment 1, shown in Table 5.1, and places it in the test-tube.
- She adds water to make the total volume of liquid equal to 20 cm³.
- She cuts a 6 cm length of magnesium ribbon, puts it in the test-tube and quickly replaces the stopper and the delivery tube.
- After 40 seconds she measures the volume of hydrogen in the measuring cylinder and records it in Table 5.1.
- She carries out experiments 2, 3, 4 and 5 in the same way.

| | | 1 Table 5 | 3 | MMN. PARA | For iner's |
|----------------|--------------------------------|---------------------------------|--|---|---------------|
| experiment no. | volume of acid/cm ³ | volume of water/cm ³ | concentration of the acid in the test-tube/mol/dm³ | volume of hydrogen after 40 s / cm ³ | Oridge Com |
| 1 | 20 | 0 | 2.0 | 80 | |
| 2 | 16 | 4 | 1.6 | | ` |
| 3 | 12 | 8 | | | |
| 4 | 8 | 12 | | 20 | |

(a) Fig. 5.2 shows the measuring cylinder readings for experiments 2, 3 and 5.

16

5

4

Read the volumes of hydrogen and record the readings in Table 5.1.





Fig. 5.2

(b) Calculate the concentration of the acid in the test-tube used in experiments 3, 4, and 5.


Complete Table 5.1. [1]

[3]



| (d) (i) | To make the experiment fair, it is important for the student to use the same of magnesium ribbon in all of these experiments. |
|---------|---|
| | Suggest two reasons for this. |
| | reason 1 |
| | reason 2 [2] |
| (ii) | What does the shape of the graph show about the relationship between the volume of hydrogen given off in 40 seconds and the concentration of the acid used? |
| | [1] |

6 The science teacher has asked the students to find the heat capacity of a metal for They heat water and then pour it into the weighed metal food can. Then they find the temperature of the water.



- The food can is weighed to find its mass, **m**.
- The room temperature, \mathbf{t}_1 , is recorded in Table 6.1.
- Some water is heated and its temperature, t₂, found.
- The water is poured into the can, it is stirred and the new temperature of the water and the metal food can \mathbf{t}_3 , is found.
- A measuring cylinder is used to find the volume of water, \mathbf{v} cm³.
- (a) Study the diagram, Fig. 6.2, and read the balance window, the measuring cylinder and the thermometers. Record these values in Table 6.1.




Fig. 6.2

Table 6.1

| | 17 able 6.1 | | g g |
|-----------------------------|------------------|------|-----------------|
| mass of metal food can | m = | | g |
| room temperature | t ₁ = | 25°C | |
| first temperature of water | t ₂ = | | °C |
| second temperature of water | t ₃ = | | °C |
| volume of water | v = | | cm ³ |

[4]

(b) (i) The room temperature was 25 °C. Find p, the increase in the temperature of the metal food can, (t₃ - 25) °C.

(ii) Find **q**, the decrease in the temperature of the water.

(iii) Find the specific heat of the metal of the food can using the formula below.

specific heat =
$$\frac{\mathbf{q} \times \text{mass of water } \times 4.2}{\mathbf{p} \times \mathbf{m}}$$
 J g⁻¹ 0 C⁻¹

specific heat of the metal =
$$Jg^{-1} \circ C^{-1}$$
 [2]

www.PapaCambridge.com (c) The teacher says that the specific heat of the metal of the food can may be found an electric heater. He sets up the apparatus shown in Fig. 6.3.




Fig. 6.3

E, the amount of energy supplied by the heater is equal to the energy absorbed by the water and the food can. To calculate E, the total energy supplied by the heater, three different variables must be measured and recorded.

Complete the list of these three values and their correct units. One of them has been done for you.

| 4 | Tl | | • | | 11 _ |
|----|------|--------|----|------|--------|
| 1. | INA | e.m. | ΤI | ın \ | mite |
| | 1110 | O.111. | | | , 0110 |

| 2. | in | |
|----|----|-----|
| 3. | in | [2] |

BLANK PAGE

www.PapaCambridge.com

20

BLANK PAGE

www.PapaCambridge.com

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.