



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

+	
_	
•	
•	
۰ د	
v	
_	
_	
`	
ว ง	
v	

CANDIDATE NAME				
CENTRE NUMBER		CANDIDATE NUMBER		

CO-ORDINATED SCIENCES

0654/52

Paper 5 Practical Test

May/June 2011

2 hours

Candidates answer on the Question Paper.

Additional Materials:

As listed in Instructions to Supervisors

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

Chemistry practical notes for this paper are printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
1	
2	
3	
Total	

This document consists of 11 printed pages and 1 blank page.



1 (a) You are going to investigate the energy content of bread.




Fig. 1.1

- Clamp a large test-tube as shown in Fig. 1.1.
- Measure 20 cm³ of water. Put this into the large test-tube.
- Clamp the thermometer in the large test-tube as shown in Fig. 1.1. Ensure the thermometer is in the water.
- Read the temperature of the water, **t**_s, and record it in Table 1.1.
- Record the mass of the bread, m_b, in Table 1.1.
- Using a mounted needle pick up the piece of bread (m_b) . Ignite the bread by placing it into a flame.
- Quickly place the burning bread under the test-tube so that it heats the water.
- If the bread stops burning, re-ignite it by placing it back into the flame.
- Record the maximum temperature, t_m , of the water.

Table 1.1

starting temperature t _s /°C	maximum temperature t _m /°C	temperature rise T/°C	mass of bread m _b /g	mass of water m _w /g
				20

[3]

www.PapaCambridge.com

(i) Calculate T, the temperature rise of the water, using the formula $T = (t_m - t_s)$ and record it in Table 1.1.

(iii) The energy content of the bread can be determined by the following formula: For iner's $E = \underbrace{0.084 \times T}_{m_b}$

Show your working.

	energy content of the bread	kJ/g	[2]
(iii)	Is the energy content that you calculated likely to be accurate and close energy content?	e to the t	true
	Explain your answer.		
			[2]
(iv)	Suggest one way in which the accuracy of the experiment could be important	roved.	
			•••••
			[1]

- ube rack. For iner's
- (b) Label three hard glass test-tubes A, B, and C and put into the test-tube rack.
 - Place a piece of fresh bread into each test-tube.
 - Add about 5 cm³ of water and gently break the bread up using a glass rod.
 - Add four drops of iodine solution to tube A. Record the colour change of the solution, if any, in Table 1.2.
 - Add about 5 cm³ of Benedict's solution to **tube B** and place in a hot water bath for 5 min. Record the colour change of the solution, if any, in Table 1.2.
 - Add about 5 cm³ of biuret solution to **tube C**. Record the colour change of the solution, if any, in Table 1.2.

Table 1.2

	iodine (tube A)	Benedict's (tube B)	biuret (tube C)
colour change	from	from	from
	to	to	to
conclusion			

Use the information you have recorded in Table 1.2 to draw conclusions about the chemical composition of bread. [6]

- 2 You are going to find the mass of a metre rule using the principle of moments.
 - Push the pin into the cork.
 - Clamp the cork so that the pin protrudes horizontally as shown in Fig. 2.1.
- www.papaCambridge.com Suspend the rule from the pin at the 100 mm mark as shown in Fig. 2.2. Ensure the rule is free to pivot about the pin.
 - Attach the other end of the rule to the newton meter, at the 900 mm mark, as shown in Fig. 2.2.
 - Suspend the newton meter from the clamp and stand as shown in Fig. 2.2.
 - Attach the 500 g mass at the 200 mm mark on the rule.
 - Make the rule horizontal by adjusting the height of the clamp holding the newton meter.




Fig. 2.1

Fig. 2.2 shows the apparatus when set up.




Fig. 2.2

- in Fig. For iner's
- (a) (i) Record the distance d, from the pivot to the mass, as shown in Fig. Table 2.1.
 - (ii) Read the force shown on the scale of the newton meter.

Record this value in Table 2.1.

(b) (i) Move the 500 g mass to the 250 mm mark on the rule. Read the newton meter again.

Record the value of **d** and the force in Table 2.1.


(ii) Repeat **three** more times, moving the mass 50 mm nearer the centre of the rule each time.

Table 2.1

d/mm	force/N

[4]

(c) (i) Plot a graph of force (vertical axis) against distance.



(ii) Extend the line so that it cuts the vertical axis.

[1]

(d) (i) Read off the value of the force when the distance d = 0.

value of force = _____N [1]

(ii) Convert the force into a mass, m, in grams. Remember that the force of gravity on 100g is 1.0 N.

> mass **m** = _____ g [1]

HANNA BARBIC AND For iner's

(iii) The mass of the rule is equal to $2\,\text{m}$.

Using your value from **(d)(ii)**, calculate the mass of the rule.

	mass of the rule = g [2]
(e)	The accuracy could be improved by making sure the rule is exactly horizontal befor taking readings.	е
	Suggest one way by which you could make sure the rule is horizontal.	
		•••
		2]

		the transfer of the transfer o	
		9	1
		nixture of two compounds each containing different ions. Carry out the foliatentify all the ions in the compounds. at is the colour of solid X ? Place about 10 cm ³ of water into the large test-tube. Add all of the solid X a stopper the tube. Shake the contents for about a minute. Filter the contents of the solid X and the solid X a	Cann
(a)	Wha	at is the colour of solid X ?	[1]
	(i)	Place about 10 cm ³ of water into the large test-tube. Add all of the solid X a stopper the tube. Shake the contents for about a minute. Filter the contents of test-tube.	ind the
		Keep the filtrate for testing in part (b) . Keep the residue in the filter paper in t funnel for testing in part (c) .	the
	(ii)	What is the colour of the	
		residue on the filter paper,	
		filtrate?	[2]
(b)	Car	ry out the following tests on the filtrate from (a), recording all your observations.	
	(i)	Place about 1 cm ³ of the filtrate into a clean test-tube. Gradually add aqueo sodium hydroxide until there is no further change.	us
		observation	[2]
	(ii)	Place another 1 cm³ portion of the filtrate into a clean test-tube. Add aqued ammonia, a little at a time until there is no further change.	us
		observations	
			[3]
	(iii)	To another 1cm³ portion of the filtrate, add a few drops of dilute hydrochloric a followed by aqueous barium chloride.	cid
		observation	[1]
	(iv)	Name the two ions in the filtrate.	
		cation	
		anion	[2]

		Why.	
		10	
(c)		I about 2 cm ³ of dilute hydrochloric acid to the residue in the filter paper freecting the filtrate for use in test (d).	Cann
	Red	cord any observations and name the anion in the residue.	
	obs	ervations	[1]
	nan	ne of anion	[1]
(d)	(i)	Slowly add aqueous sodium hydroxide to the filtrate from (c) , to make the solut alkaline.	tion
		Record your observation.	
		observation	[1]
	(ii)	Suggest the name of the cation in the filtrate.	
		name of cation	[1]

BLANK PAGE

www.PapaCambridge.com

CHEMISTRY PRACTICAL NOTES

Test for anions

Test for anions	12 CHEMISTRY PRACTICAL NO	TES test result
anion	test	test result
carbonate (CO ₃ ² -)	add dilute acid	effervescence, carbon dioxide produced
chloride (Cl ⁻) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
nitrate (NO ₃ -) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO ₄ ²⁻) [in solution]	acidify then add aqueous barium chloride <i>or</i> aqueous barium nitrate	white ppt.

Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
ammonium (NH ₄ ⁺)	ammonia produced on warming	-
copper(II) (Cu ²⁺)	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe ²⁺)	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe ³⁺)	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn ²⁺)	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

Test for gases

gas	test and test results
ammonia (NH ₃)	turns damp red litmus paper blue
carbon dioxide (CO ₂)	turns limewater milky
chlorine (Cl ₂)	bleaches damp litmus paper
hydrogen (H ₂)	"pops" with a lighted splint
oxygen (O ₂)	relights a glowing splint

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.