



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS  
General Certificate of Education Ordinary Level

CANDIDATE  
NAME

CENTRE  
NUMBER

--	--	--	--	--

CANDIDATE  
NUMBER

--	--	--	--



**CHEMISTRY**

**5070/21**

Paper 2 Theory

**October/November 2012**

**1 hour 30 minutes**

Candidates answer on the Question Paper.

No Additional Materials are required.

**READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES.

**Section A**

Answer **all** questions.

Write your answers in the spaces provided in the Question Paper.

**Section B**

Answer any **three** questions.

Write your answers in the spaces provided in the Question Paper.

A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use	
<b>Section A</b>	
<b>B7</b>	
<b>B8</b>	
<b>B9</b>	
<b>B10</b>	
<b>Total</b>	

This document consists of **18** printed pages and **2** blank pages.



## Section A

Answer **all** the questions in this section in the spaces provided.

The total mark for this section is 45.

A1 (a) Define the term *element*.

..... [1]

(b) Choose from the following elements to answer the questions below.

**aluminium**

**argon**

**bromine**

**gallium**

**helium**

**hydrogen**

**magnesium**

**nitrogen**

**oxygen**

**sodium**

Each element can be used once, more than once or not at all.

Which element

(i) is in Group III and Period 4 of the Periodic Table, ..... [1]

(ii) has atoms with 8 electrons in their outer shell, ..... [1]

(iii) is a liquid at room temperature, ..... [1]

(iv) reduces unsaturated vegetable oils to form a solid product, ..... [1]

(v) forms an ionic chloride with the formula  $XCl_2$ , ..... [1]

(vi) is used in light bulbs? ..... [1]

(c) Draw the electronic structure of an aluminium atom.

[1]

[Total: 8]

**A2** Steel is more resistant to corrosion than iron.

**(a)** What are the essential conditions for the corrosion of iron?

.....  
 ..... [1]

**(b)** Ships' hulls can be prevented from corroding by attaching pieces of magnesium to them. Explain why this prevents the hulls from corroding.

.....  
 ..... [2]

**(c)** Steel is an alloy. Explain the meaning of the term *alloy*.

.....  
 ..... [1]

**(d)** Samples of iron were placed in aqueous solutions having different pH values. The table shows how the speed of corrosion of iron varies with the pH of the solution.

speed of corrosion/cm per year	0.043	0.029	0.012	0.010	0.010	0.010	0.009	0.006
pH	2	3	4	5	6	8	10	12

Describe how pH affects the speed of corrosion of iron.

.....  
 .....  
 ..... [2]

[Total: 6]

**A3** The table below shows both the formulae and boiling points of the first five members of an alcohol homologous series.

alcohol	formula	boiling point /°C
methanol	CH <sub>3</sub> OH	65
ethanol	C <sub>2</sub> H <sub>5</sub> OH	79
propanol	C <sub>3</sub> H <sub>7</sub> OH	98
butanol	C <sub>4</sub> H <sub>9</sub> OH	117
pentanol	C <sub>5</sub> H <sub>11</sub> OH	138

**(a) (i)** Deduce the formula of the sixth member of the alcohol homologous series.

..... [1]

**(ii)** Predict the boiling point of this alcohol.

..... [1]

**(b)** Ethanol can be made industrially by fermentation.

Describe one other method of making ethanol industrially, stating the conditions required for the reaction.

.....  
 .....  
 .....  
 ..... [3]

**(c) (i)** Ethanol can be oxidised to ethanoic acid by atmospheric oxygen. Name one other suitable oxidising agent which can be used.

..... [1]

**(ii)** Propanol can be oxidised to propanoic acid. Draw the structure for propanoic acid.

[1]

[Total: 7]

**A4** Water from natural sources, such as lakes and rivers, contains many dissolved substances.

**(a)** Name two dissolved substances that occur naturally in unpolluted water from lakes and rivers.

..... [1]

**(b)** Pollution in lakes and rivers can be caused by leaching of fertilisers from farmland. This can cause eutrophication.

**(i)** Name two ions present in fertilisers which cause eutrophication.


..... [2]

**(ii)** Describe the essential stages in eutrophication.

.....  
.....  
.....  
.....  
.....  
.....  
..... [4]


[Total: 7]

- A5** The diagram below shows the apparatus used to electrolyse aqueous sodium hydroxide in the laboratory.



Electrolysis of the aqueous sodium hydroxide, results in the formation of hydrogen at the cathode (negative electrode) and oxygen at the anode (positive electrode).

- (a)** Complete the equation for the formation of oxygen at the anode.



- (b) (i)** When the power pack is replaced by a voltmeter, the apparatus acts like a fuel cell. The left hand electrode in the diagram becomes the negative pole of the cell and the right hand electrode becomes the positive pole.

State the direction of the electron flow in the external circuit.  
Give a reason for your answer.

.....  
..... [1]

- (ii)** In this fuel cell, hydrogen reacts with aqueous hydroxide ions to form water. Construct an equation for this reaction.

[1]


- (c) (i)** Suggest two advantages of using a fuel cell rather than petrol to power a car.

.....  
.....  
..... [2]

- (ii)** Suggest one disadvantage of fuel cells.

..... [1]


**A6** A flow diagram for the manufacture of sulfuric acid is shown below.



- (a)** In the sulfur burner, a spray of molten sulfur is burned in a furnace. Construct an equation for this reaction. Include state symbols.

[1]

- (b)** In the converter, the following reaction occurs:



The yield of  $\text{SO}_3$  is 95% at  $450^\circ\text{C}$  and atmospheric pressure.

- (i)** Name the catalyst used in this reaction.

..... [1]

- (ii)** Explain why increasing the pressure shifts the position of equilibrium further to the right.

..... [1]

- (iii)** Explain why the reaction is carried out at atmospheric pressure even though an increase in pressure shifts the position of equilibrium further to the right.


..... [1]

- (iv)** Explain why the reaction is carried out at  $450^\circ\text{C}$  and not at a higher or lower temperature.

.....  
 .....  
 ..... [3]




- (c) Sulfuric acid is formed from sulfur trioxide in two stages.  
Firstly, the sulfur trioxide,  $\text{SO}_3$ , is absorbed in concentrated sulfuric acid to form oleum,  $\text{H}_2\text{S}_2\text{O}_7$ .



The oleum is then mixed with water to form sulfuric acid.  
Construct an equation for this reaction.

[1]

- (d) Aqueous sulfuric acid is titrated with aqueous sodium hydroxide.



It requires  $28.0 \text{ cm}^3$  of  $0.100 \text{ mol/dm}^3$  aqueous sodium hydroxide to neutralise  $9.50 \text{ cm}^3$  of sulfuric acid.

Calculate the concentration, in  $\text{mol/dm}^3$ , of the aqueous sulfuric acid.

Give your answer to 3 significant figures.

concentration of the aqueous sulfuric acid .....  $\text{mol/dm}^3$  [3]

[Total: 11]

## Section B

Answer **three** questions from this section in the spaces provided.

The total mark for this section is 30.

**B7** Tin is a metal in Group IV of the Periodic Table.

(a) Draw a labelled diagram to show the structure of a metal.

[2]

(b) Explain why metals

(i) conduct electricity, .....

(ii) are malleable. ....

..... [2]

(c) At high temperatures, tin reacts with steam to form tin(II) oxide, SnO, and one other product.

This reaction is reversible.

The other product is a gas which gives a 'pop' with a lighted splint.

(i) Construct an equation for this reaction.

[1]


(ii) Tin(II) oxide is an amphoteric oxide.  
Explain the meaning of the term *amphoteric oxide*.

..... [1]

(d) (i) Concentrated nitric acid reacts with tin to form tin(IV) oxide, SnO<sub>2</sub>, nitrogen dioxide and water.

Construct an equation for this reaction.

[1]



- (ii) Nitric acid contains nitrate ions.  
Describe a test for nitrate ions.  
Give the result of a positive test.

.....

.....

..... [3]

[Total: 10]

**B8** Petroleum is separated into fractions by fractional distillation.

**(a)** Explain how fractional distillation separates petroleum into different fractions.

.....  
.....  
.....  
.....  
..... [3]

**(b)** The refinery gas fraction contains the first four members of the alkane homologous series.

**(i)** Explain the meaning of the term *homologous series*.

.....  
.....  
..... [2]

**(ii)** Draw the structure, showing all atoms and bonds, of the two isomers of butane, the fourth member of the alkane homologous series.

[2]

**(c)** Construct an equation for the complete combustion of hexane,  $C_6H_{14}$ .

[1]

(d) When long-chained alkanes are cracked in an oil refinery, shorter-chained alkanes and alkenes are formed.

(i) Explain why the process of cracking needs to be carried out.

.....  
..... [1]

(ii) Describe a chemical test to distinguish between an alkane and an alkene.

test .....


result ..... [1]

[Total: 10]

**B9 (a)** Define the term *relative atomic mass*.

.....  
 ..... [1]

**(b)** The relative atomic mass of magnesium can be determined in the laboratory by finding the volume of hydrogen given off when magnesium reacts with hydrochloric acid.




0.036 g of magnesium reacts at room temperature and pressure with excess hydrochloric acid to produce 36 cm<sup>3</sup> of hydrogen.

1 mole of any gas at room temperature and pressure occupies 24 dm<sup>3</sup>.

Show by calculation that the relative atomic mass of magnesium is 24.

[3]

**(c)** Magnesium reacts with oxygen in the air to form magnesium oxide.



**(i)** If the yield of the reaction is 75% calculate the mass of magnesium oxide formed when 12 kg of magnesium burns in excess air.

[2]

**(ii)** Magnesium nitride is also formed when magnesium burns in air. Magnesium nitride is an ionic compound. Deduce the formula for magnesium nitride.

..... [1]

(d) When magnesium is heated with silicon, magnesium silicide,  $Mg_2Si$ , is formed. Magnesium silicide reacts with water to form silane,  $SiH_4$ , and magnesium oxide.

(i) Construct an equation for the reaction of magnesium silicide with water.

[1]

(ii) Silane has a structure similar to methane. Draw a 'dot-and-cross' diagram for silane. Show only the outer shell electrons.

[1]

(iii) Silane reacts with oxygen to form silicon dioxide and water. Construct an equation for this reaction.

[1]

[Total: 10]

**B10** Limestone consists mainly of the compound calcium carbonate.

- (a) Explain why limestone is used in the blast furnace for the extraction of iron. Include any relevant equations in your answer.

.....

.....

.....

.....

.....


..... [3]

- (b) Group II carbonates decompose on heating. The temperatures at which some Group II carbonates decompose are given in the table below.

Group II carbonate	decomposition temperature /°C
barium carbonate	1360
calcium carbonate	900
magnesium carbonate	540
strontium carbonate	1280

- (i) Which one of these carbonates is least likely to decompose on heating?  
..... [1]
- (ii) Describe how the thermal stability of these carbonates changes with the reactivity of the metal.  
..... [1]





(c) The speed of reaction of calcium carbonate with hydrochloric acid can be calculated by measuring the volume of gas given off at various time intervals.

(i) Draw a labelled diagram of the apparatus you could use to follow the course of this reaction.

[2]

(ii) State and explain the effect of the following on the volume of a fixed mass of gas

- increasing the pressure,
- increasing the temperature.

.....

.....

.....

..... [3]

[Total: 10]





**DATA SHEET**  
**The Periodic Table of the Elements**

		Group																																																																																											
I	II	III	IV	V	VI	VII	0																																																																																						
7 <b>Li</b> Lithium 4	9 <b>Be</b> Beryllium 4	1 <b>H</b> Hydrogen 1	11 <b>B</b> Boron 5	12 <b>C</b> Carbon 6	13 <b>Al</b> Aluminium 13	14 <b>Si</b> Silicon 14	15 <b>P</b> Phosphorus 15	16 <b>S</b> Sulfur 16	17 <b>Cl</b> Chlorine 17	18 <b>Ar</b> Argon 18	19 <b>F</b> Fluorine 9	20 <b>Ne</b> Neon 10	2 <b>He</b> Helium 2																																																																																
23 <b>Na</b> Sodium 12	24 <b>Mg</b> Magnesium 12	27 <b>Co</b> Cobalt 27	28 <b>Ni</b> Nickel 28	29 <b>Cu</b> Copper 29	30 <b>Zn</b> Zinc 30	31 <b>Ga</b> Gallium 31	32 <b>Ge</b> Germanium 32	33 <b>As</b> Arsenic 33	34 <b>Se</b> Selenium 34	35 <b>Br</b> Bromine 35	36 <b>Kr</b> Krypton 36	37 <b>Rb</b> Rubidium 37	38 <b>Sr</b> Strontium 38	39 <b>Y</b> Yttrium 39	40 <b>Ca</b> Calcium 20	41 <b>K</b> Potassium 19	42 <b>V</b> Vanadium 23	43 <b>Cr</b> Chromium 24	44 <b>Mn</b> Manganese 25	45 <b>Fe</b> Iron 26	46 <b>Ru</b> Ruthenium 44	47 <b>Rh</b> Rhodium 45	48 <b>Pd</b> Palladium 46	49 <b>Cd</b> Cadmium 48	50 <b>In</b> Indium 49	51 <b>Tl</b> Thallium 81	52 <b>Hg</b> Mercury 80	53 <b>Au</b> Gold 79	54 <b>Pt</b> Platinum 78	55 <b>Ir</b> Iridium 77	56 <b>Os</b> Osmium 76	57 <b>Rf</b> Rutherfordium 104	58 <b>Ce</b> Cerium 58	59 <b>Pr</b> Praseodymium 59	60 <b>Nd</b> Neodymium 60	61 <b>Pm</b> Promethium 61	62 <b>Sm</b> Samarium 62	63 <b>Eu</b> Europium 63	64 <b>Gd</b> Gadolinium 64	65 <b>Tb</b> Terbium 65	66 <b>Dy</b> Dysprosium 66	67 <b>Ho</b> Holmium 67	68 <b>Er</b> Erbium 68	69 <b>Tm</b> Thulium 69	70 <b>Yb</b> Ytterbium 70	71 <b>Lu</b> Lutetium 71	72 <b>Hf</b> Hafnium 72	73 <b>Ta</b> Tantalum 73	74 <b>W</b> Tungsten 74	75 <b>Re</b> Rhenium 75	76 <b>Os</b> Osmium 76	77 <b>Ir</b> Iridium 77	78 <b>Pt</b> Platinum 78	79 <b>Au</b> Gold 79	80 <b>Hg</b> Mercury 80	81 <b>Tl</b> Thallium 81	82 <b>Pb</b> Lead 82	83 <b>Bi</b> Bismuth 83	84 <b>Po</b> Polonium 84	85 <b>At</b> Astatine 85	86 <b>Rn</b> Radon 86	87 <b>Fr</b> Francium 87	88 <b>Ra</b> Radium 88	89 <b>Ac</b> Actinium 89	90 <b>Th</b> Thorium 90	91 <b>Pa</b> Protactinium 91	92 <b>U</b> Uranium 92	93 <b>Np</b> Neptunium 93	94 <b>Pu</b> Plutonium 94	95 <b>Am</b> Americium 95	96 <b>Cm</b> Curium 96	97 <b>Bk</b> Berkelium 97	98 <b>Cf</b> Californium 98	99 <b>Es</b> Einsteinium 99	100 <b>Fm</b> Fermium 100	101 <b>Md</b> Mendelevium 101	102 <b>No</b> Nobelium 102	103 <b>Lr</b> Lawrencium 103	104 <b>Rf</b> Rutherfordium 104	105 <b>Db</b> Dubnium 105	106 <b>Sg</b> Seaborgium 106	107 <b>Bh</b> Bohrium 107	108 <b>Hs</b> Hassium 108	109 <b>Mt</b> Meitnerium 109	110 <b>Ds</b> Darmstadtium 110	111 <b>Rg</b> Roentgenium 111	112 <b>Cn</b> Copernicium 112	113 <b>Nh</b> Nihonium 113	114 <b>Fl</b> Flerovium 114	115 <b>Mc</b> Moscovium 115	116 <b>Lv</b> Livermorium 116	117 <b>Ts</b> Tennessine 117	118 <b>Og</b> Oganesson 118

8–71 Lanthanoid series  
90–103 Actinoid series

a = relative atomic mass  
X = atomic symbol  
b = atomic (proton) number

The volume of one mole of any gas is 24dm<sup>3</sup> at room temperature and pressure (r.t.p.).