

**CAMBRIDGE INTERNATIONAL EXAMINATIONS**

Cambridge Ordinary Level

## **MARK SCHEME for the October/November 2015 series**

### **4037 ADDITIONAL MATHEMATICS**

**4037/13**

Paper 1, maximum raw mark 80

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.




Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2015 series for most Cambridge IGCSE<sup>®</sup>, Cambridge International A and AS Level components and some Cambridge O Level components.

® IGCSE is the registered trademark of Cambridge International Examinations.

### Abbreviations


Awrt	answers which round to
Cao	correct answer only
dep	dependent
FT	follow through after error
isw	ignore subsequent working
oe	or equivalent
rot	rounded or truncated
SC	Special Case
soi	seen or implied
www	without wrong working

<b>1</b>	<b>(i)</b>		B1	
	<b>(ii)</b>		B1	
	<b>(iii)</b>		B1	
<b>2</b>	$\cos\left(3x - \frac{\pi}{4}\right) = (\pm)\frac{1}{\sqrt{2}} \text{ oe}$ $3x - \frac{\pi}{4} = -\frac{\pi}{4}, \frac{\pi}{4}, \frac{3\pi}{4}$ $x = \left(-\frac{\pi}{4} + \frac{\pi}{4}\right) \div 3, \left(\frac{\pi}{4} + \frac{\pi}{4}\right) \div 3, \left(\frac{3\pi}{4} + \frac{\pi}{4}\right) \div 3 \text{ oe}$ $x = 0 \text{ and } \frac{\pi}{6} \text{ (or 0 and 0.524)}$ $x = \frac{\pi}{3} \text{ (or 1.05)}$	M1   DM1  A2/1/0	division by 2 and square root   correct order of operations in order to obtain a solution  A2 for 3 solutions and no extras in the range A1 for 2 solutions A0 for one solution or no solutions	

Page 3	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2015	4037	13

<p>3 (a)</p> $\begin{pmatrix} 12 & 16 & 4 \\ 30 & 32 & 10 \end{pmatrix}$ <p>(b)</p> $\begin{pmatrix} 28 & -24 \\ -8 & 76 \end{pmatrix} = m \begin{pmatrix} 4 & 6 \\ 2 & -8 \end{pmatrix} + n \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ <p><math>-24 = 6m</math> or <math>-8 = 2m</math> giving <math>m = -4</math></p> <p><math>28 = 4m + n</math> or <math>76 = -8m + n</math>  <math>n = 44</math></p> <p>(c)</p> $a^2 - 6 = 0$ <p>so <math>a = \pm\sqrt{6}</math></p>		<p>B2,1,0</p> <p>B2,1,0</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B2,1,0</p>	<p>B2 for 6 elements correct, B1 for 5 elements correct</p> <p>B2 for 4 correct elements in <math>\mathbf{X}^2</math> B1 for 3 correct elements in <math>\mathbf{X}^2</math></p> <p>For <math>m = -4</math> using correct <b>I</b></p> <p>complete method to obtain <math>n</math></p> <p>or B1 for <math>a = \pm\sqrt{6}</math> or <math>a = \pm 2.45</math>, with no incorrect statements seen or B1 for <math>a = \pm\sqrt{6}</math> or <math>a = \pm 2.45</math> seen or B1 for <math>a = \sqrt{6}</math> and no incorrect working</p>
<p>4 (i)</p> $\frac{1}{2}(4\sqrt{3}+1) \times BC = \frac{47}{2}$ $BC = \frac{47}{(4\sqrt{3}+1)} \times \frac{(4\sqrt{3}-1)}{(4\sqrt{3}-1)}$ $BC = 4\sqrt{3}-1$ <p><b>Alternative method</b></p> $\frac{1}{2}(4\sqrt{3}+1) \times BC = \frac{47}{2}$ $(4\sqrt{3}+1)(a\sqrt{3}+b) = 47$ <p>Leading to <math>12a + b = 47</math> and <math>a + 4b = 0</math> Solution of simultaneous equations</p> $BC = 4\sqrt{3}-1$ <p>(ii)</p> $(4\sqrt{3}+1)^2 + (4\sqrt{3}-1)^2$ $= (48 + 8\sqrt{3} + 1) + (48 - 8\sqrt{3} + 1)$ $AC^2 = 98$ $AC = 7\sqrt{2} \text{ or } p = 7$		<p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B1FT</p> <p>B1cao</p>	<p>correct use of the area</p> <p>correct rationalisation</p> <p>Dependent on all method being seen</p> <p>Dependent on all method seen including solution of simultaneous equations</p> <p>6 correct FT terms seen</p> <p>98 and <math>7\sqrt{2}</math> or 98 and <math>p = 7</math></p>

Page 4	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2015	4037	13

5	<p>When <math>x = \frac{\pi}{4}</math>, <math>y = 2</math></p> $\frac{dy}{dx} = 5\sec^2 x$ <p>When <math>x = \frac{\pi}{4}</math>, <math>\frac{dy}{dx} = 10</math></p> <p>Equation of normal <math>y - 2 = -\frac{1}{10}\left(x - \frac{\pi}{4}\right)</math></p> $10y + x - 20 - \frac{\pi}{4} = 0 \quad \text{or} \quad 10y + x - 20.8 = 0 \quad \text{oe}$	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p>	<p><math>y = 2</math></p> <p><math>5\sec^2 x</math></p> <p>10 from differentiation</p> $y - \text{their } 2 = -\frac{1}{\text{their } 10}\left(x - \frac{\pi}{4}\right)$ <p>allow unsimplified</p>
6	<p>(i)</p>  <p>(ii)</p> <p>(2, 16)</p> <p>(iii)</p> <p><math>k = 0</math></p> <p><math>k &gt; 16</math></p>	<p>B1</p> <p>B1</p> <p>B1</p> <p>M1</p> <p>A1</p> <p>B1</p> <p>B1</p>	<p>shape</p> <p>intercepts on <math>x</math>-axis</p> <p>intercept on <math>y</math>-axis for a curve with a maximum and two arms</p> <p><math>(2, \pm 16)</math> seen or <math>(2, k)</math> where <math>k &gt; 0</math></p> <p><math>(2, 16)</math> or <math>x = 2</math> <b>and</b> <math>y = 16</math> only</p>

Page 5	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2015	4037	13

7	$\frac{dy}{dx} = 2 \sin 3x \quad (+c)$ $4\sqrt{3} = 2 \frac{\sqrt{3}}{2} + c$ $\frac{dy}{dx} = 2 \sin 3x + 3\sqrt{3}$ $y = -\frac{2}{3} \cos 3x + 3\sqrt{3}x \quad (+d)$ $-\frac{1}{3} = -\frac{2}{3} \cos \frac{\pi}{3} + 3\sqrt{3} \left( \frac{\pi}{9} \right) + d$ $y = -\frac{2}{3} \cos 3x + 3\sqrt{3}x - \frac{\sqrt{3}}{3} \pi$	<p>B1</p> <p>M1</p> <p>A1</p> <p>B1FT</p> <p>M1</p> <p>A1</p>	<p>2 sin 3x</p> <p>finding constant using <math>\frac{dy}{dx} = k \sin 3x + c</math> making use of <math>\frac{dy}{dx} = 4\sqrt{3}</math> and <math>x = \frac{\pi}{9}</math></p> <p>Allow with <math>c = 5.20</math> or <math>\sqrt{27}</math></p> <p>FT integration of <i>their</i> <math>k \sin 3x</math></p> <p>finding constant <math>d</math> for <math>k \cos 3x + cx + d</math></p> <p>Allow  <math>y = -0.667 \cos 3x + 5.20x - 0.577\pi</math>  or better</p>
8	<p>(a)</p> $(2 + kx)^8 = 256 + 1024kx + 1792k^2x^2 + 1792k^3x^3$ $k = \frac{1}{4}$ $p = 112$ $q = 28$ <p>(b)</p> ${}^9C_3 x^6 \left( -\frac{2}{x^2} \right)^3$ $84x^6 \left( -\frac{8}{x^6} \right) \text{ leading to } -672$	<p>B1</p> <p>B1FT</p> <p>B1FT</p> <p>M1</p> <p>DM1</p> <p>A1</p>	<p>FT 1792 multiplied by <i>their</i> <math>k^2</math></p> <p>FT 1792 multiplied by <i>their</i> <math>k^3</math></p> <p>correct term seen</p> <p>Term selected and <math>2^3</math> and <math>{}^9C_3</math> correctly evaluated</p>

Page 6	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2015	4037	13

9	(a) (i)	Number of arrangements with Maths books as one item = $4!$ or $4 \times 3!$	M1	$4!(\times 2)$ or $4 \times 3!(\times 2)$ oe
		or Maths books can be arranged 2! ways and History 3! ways = $2! \times 3!$		$2! \times 3!(\times 4)$ or $2 \times 3!(\times 4)$ oe
		$2 \times 4!$ or $2 \times 4 \times 3!$ or $4 \times 2 \times 3! = 48$	A1	A1 for 48
	(ii)	$5! - 48$ or $6 \times 2 \times 3!$	M1	$5!$ – <i>their</i> answer to (i)
		72	A1	or for $6 \times 2 \times 3$
	(b) (i)	3003	B1	
	(ii)	$3003 - 6 - 135$	M1	<i>their</i> answer to (i) – $6 - {}^6C_4 \times 9$
		2862	B1	135 subtracted
		or	A1	
		$2M\ 3W = 720$	M1	complete correct method using 4 cases, may be implied by working. Must have at least one correct
	$3M\ 2W = 1260$			
	$4M\ 1W = 756$			
	$5M = 126$	B1	any 3 correct	
	2862	A1		

Page 7	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2015	4037	13

10	(i)	$10^2 = 6^2 + 6^2 - 2 \times 6 \times 6 \times \cos ABC$ <p>or</p> $\sin\left(\frac{ABC}{2}\right) = \frac{5}{6}$ <p>or</p> $ABC = \pi - \sin^{-1} \frac{10\sqrt{11}}{36}$ $ABC = 1.9702$	M1	correct cosine rule statement or correct statement for $\sin \frac{ABC}{2}$ or equating areas oe
	(ii)	$XY = 2$ <p>Arc length <math>6\left(\frac{\pi - 1.970}{2}\right)</math> oe</p> <p>Perimeter = <math>2 + 2\left(6\left(\frac{\pi - 1.970}{2}\right)\right)</math> = 9.03</p>	B1 B1 M1 A1	1.9702 or better for $XY$ ( may be implied by later work, allow on diagram) correct arc length (unsimplified) <i>their</i> $2 + 2 \times 6 \times$ <i>their</i> angle $C$
	(iii)	$\left(\frac{1}{2} \times 6^2 \left(\frac{\pi - 1.970}{2}\right) - \frac{1}{2} \times 5 \times \sqrt{11}\right) \times 2$ <p>= 4.50 or 4.51 or better</p>	M1 M1 A1	sector area using <i>their</i> $C$ area of $\triangle ABM$ where $M$ is the midpoint of $AC$ , or ( $\triangle s$ $ABY$ and $BXY$ ) or $\triangle ABC$ Answers to 3sf or better





Page 9	Mark Scheme	Syllabus	Paper
	Cambridge O Level – October/November 2015	4037	13

12 (a)	$2^{2x-1} \times 2^{2(x+y)} = 2^7$ and $\frac{3^{2(2y-x)}}{3^{3(y-4)}} = 1$  $2x - 1 + 2(x + y) = 7$ oe $2(2y - x) = 3(y - 4)$ oe leading to $x = 4, y = -4$  <u>Example of Alternative method</u> Method mark as above $2x - 1 + 2(x + y) = 7$  leading to $y = \frac{(8 - 4x)}{2}$  Correctly substituted in $\frac{3^{2(2y-x)}}{3^{3(y-4)}} = 1$  Leading to $2\left(\frac{2(8 - 4x)}{2} - x\right) = 3\left(\frac{(8 - 4x)}{2} - 4\right)$ Leading to $x = 4$ and $y = -4$	M1	expressing $4^{x+y}$ , 128 as powers of 2 and $9^{2y-x}$ , $27^{y-4}$ as powers of 3
		A1 A1 A1	Correct equation from correct working Correct equation from correct working for both
(b)	$(2(5^z) - 1)(5^z + 1) = 0$ leading to $2.5^z = 1$ ( $5^z = -1$ )  $5^z = 0.5$  $z = \frac{\log 0.5}{\log 5}$ or $z = -0.431$ or better	M1 A1 DM1 A1	As before One of the correct equations in $x$ and $y$  Correct, unsimplified, equation in $x$ or $y$ only Both answers  solution of quadratic correct solution  correct attempt to solve $2.5^z = k$ , where $k$ is positive  must have one solution only